

# Microbiology



## OpenStax

Rice University  
6100 Main Street MS-375  
Houston, Texas 77005

To learn more about OpenStax, visit <http://openstaxcollege.org>.  
Individual print copies and bulk orders can be purchased through our website.

© 2016 Rice University. Textbook content produced by OpenStax is licensed under a Creative Commons Attribution 4.0 International License. Under this license, any user of this textbook or the textbook contents herein must provide proper attribution as follows:

- If you redistribute this textbook in a digital format (including but not limited to EPUB, PDF, and HTML), then you must retain on every page the following attribution:  
"Download for free at <http://cnx.org/content/col12063/latest/>."
- If you redistribute this textbook in a print format, then you must include on every physical page the following attribution:  
"Download for free at <http://cnx.org/content/col12063/latest/>."
- If you redistribute part of this textbook, then you must retain in every digital format page view (including but not limited to EPUB, PDF, and HTML) and on every physical printed page the following attribution:  
"Download for free at <http://cnx.org/content/col12063/latest/>."
- If you use this textbook as a bibliographic reference, then you should cite it as follows: OpenStax, *Microbiology*. OpenStax. 26 09 2016. <<http://cnx.org/content/col12063/latest/>>.

For questions regarding this licensing, please contact [partners@openstaxcollege.org](mailto:partners@openstaxcollege.org).

## Trademarks

The OpenStax name, OpenStax logo, OpenStax book covers, OpenStax CNX name, OpenStax CNX logo, Connexions name, and Connexions logo are not subject to the license and may not be reproduced without the prior and express written consent of Rice University.

**ISBN-10**      **1 -938168 -14 -3**  
**ISBN-13**      **978 -1 -938168 -14 -7**  
**Revision**     **MB -2016 -001(09/16) -BKB**

## OpenStax

OpenStax is a non-profit organization committed to improving student access to quality learning materials. Our free textbooks are developed and peer-reviewed by educators to ensure they are readable, accurate, and meet the scope and sequence requirements of modern college courses. Through our partnerships with companies and foundations committed to reducing costs for students, OpenStax is working to improve access to higher education for all.

## OpenStax CNX

The technology platform supporting OpenStax is OpenStax CNX (<http://cnx.org>), one of the world's first and largest open-education projects. OpenStax CNX provides students with free online and low-cost print editions of the OpenStax library and provides instructors with tools to customize the content so that they can have the perfect book for their course.

## Rice University

OpenStax and OpenStax CNX are initiatives of Rice University. As a leading research university with a distinctive commitment to undergraduate education, Rice University aspires to path-breaking research, unsurpassed teaching, and contributions to the betterment of our world. It seeks to fulfill this mission by cultivating a diverse community of learning and discovery that produces leaders across the spectrum of human endeavor.



---

## Foundation Support

OpenStax is grateful for the tremendous support of our sponsors. Without their strong engagement, the goal of free access to high-quality textbooks would remain just a dream.



Laura and John Arnold Foundation (LJAF) actively seeks opportunities to invest in organizations and thought leaders that have a sincere interest in implementing fundamental changes that not only yield immediate gains, but also repair broken systems for future generations. LJAF currently focuses its strategic investments on education, criminal justice, research integrity, and public accountability.



The William and Flora Hewlett Foundation has been making grants since 1967 to help solve social and environmental problems at home and around the world. The Foundation concentrates its resources on activities in education, the environment, global development and population, performing arts, and philanthropy, and makes grants to support disadvantaged communities in the San Francisco Bay Area.



Guided by the belief that every life has equal value, the Bill & Melinda Gates Foundation works to help all people lead healthy, productive lives. In developing countries, it focuses on improving people's health with vaccines and other life-saving tools and giving them the chance to lift themselves out of hunger and extreme poverty. In the United States, it seeks to significantly improve education so that all young people have the opportunity to reach their full potential. Based in Seattle, Washington, the foundation is led by CEO Jeff Raikes and Co-chair William H. Gates Sr., under the direction of Bill and Melinda Gates and Warren Buffett.



The Maxfield Foundation supports projects with potential for high impact in science, education, sustainability, and other areas of social importance.



Our mission at the Twenty Million Minds Foundation is to grow access and success by eliminating unnecessary hurdles to affordability. We support the creation, sharing, and proliferation of more effective, more affordable educational content by leveraging disruptive technologies, open educational resources, and new models for collaboration between for-profit, nonprofit, and public entities.





# Table of Contents

---

Chapter 1: An Invisible World . . . . .	5
1.1 What Our Ancestors Knew . . . . .	6
1.2 A Systematic Approach . . . . .	13
1.3 Types of Microorganisms . . . . .	19
Chapter 2: How We See the Invisible World . . . . .	33
2.1 The Properties of Light . . . . .	34
2.2 Peering Into the Invisible World . . . . .	40
2.3 Instruments of Microscopy . . . . .	43
2.4 Staining Microscopic Specimens . . . . .	61
Chapter 3: The Cell . . . . .	79
3.1 Spontaneous Generation . . . . .	80
3.2 Foundations of Modern Cell Theory . . . . .	84
3.3 Unique Characteristics of Prokaryotic Cells . . . . .	92
3.4 Unique Characteristics of Eukaryotic Cells . . . . .	113
Chapter 4: Prokaryotic Diversity . . . . .	139
4.1 Prokaryote Habitats, Relationships, and Microbiomes . . . . .	140
4.2 Proteobacteria . . . . .	147
4.3 Nonproteobacteria Gram-Negative Bacteria and Phototrophic Bacteria . . . . .	157
4.4 Gram-Positive Bacteria . . . . .	163
4.5 Deeply Branching Bacteria . . . . .	172
4.6 Archaea . . . . .	174
Chapter 5: The Eukaryotes of Microbiology . . . . .	183
5.1 Unicellular Eukaryotic Parasites . . . . .	184
5.2 The Parasitic Helminths . . . . .	200
5.3 Fungi . . . . .	207
5.4 Algae . . . . .	218
5.5 Lichens . . . . .	222
Chapter 6: Acellular Pathogens . . . . .	229
6.1 Viruses . . . . .	230
6.2 The Viral Life Cycle . . . . .	239
6.3 Isolation, Culture, and Identification of Viruses . . . . .	250
6.4 Viroids, Virusoids, and Prions . . . . .	261
Index . . . . .	275



## Chapter 1

---

# An Invisible World



**Figure 1.1** A veterinarian gets ready to clean a sea turtle covered in oil following the Deepwater Horizon oil spill in the Gulf of Mexico in 2010. After the spill, the population of a naturally occurring oil-eating marine bacterium called *Alcanivorax borkumensis* skyrocketed, helping to get rid of the oil. Scientists are working on ways to genetically engineer this bacterium to be more efficient in cleaning up future spills. (credit: modification of work by NOAA's National Ocean Service)

### Chapter Outline

- 1.1 What Our Ancestors Knew
- 1.2 A Systematic Approach
- 1.3 Types of Microorganisms

## Introduction

---

From boiling thermal hot springs to deep beneath the Antarctic ice, microorganisms can be found almost everywhere on earth in great quantities. Microorganisms (or microbes, as they are also called) are small organisms. Most are so small that they cannot be seen without a microscope.

Most microorganisms are harmless to humans and, in fact, many are helpful. They play fundamental roles in ecosystems everywhere on earth, forming the backbone of many food webs. People use them to make biofuels, medicines, and even foods. Without microbes, there would be no bread, cheese, or beer. Our bodies are filled with microbes, and our skin alone is home to trillions of them.<sup>[1]</sup> Some of them we can't live without; others cause diseases that can make us sick or even kill us.

Although much more is known today about microbial life than ever before, the vast majority of this invisible world remains unexplored. Microbiologists continue to identify new ways that microbes benefit and threaten humans.

---

1. J. Hulcr et al. "A Jungle in There: Bacteria in Belly Buttons are Highly Diverse, but Predictable." *PLoS ONE* 7 no. 11 (2012): e47712. doi:10.1371/journal.pone.0047712.

## 1.1 What Our Ancestors Knew

### Learning Objectives

- Describe how our ancestors improved food with the use of invisible microbes
- Describe how the causes of sickness and disease were explained in ancient times, prior to the invention of the microscope
- Describe key historical events associated with the birth of microbiology


Most people today, even those who know very little about microbiology, are familiar with the concept of microbes, or “germs,” and their role in human health. Schoolchildren learn about bacteria, viruses, and other microorganisms, and many even view specimens under a microscope. But a few hundred years ago, before the invention of the microscope, the existence of many types of microbes was impossible to prove. By definition, **microorganisms**, or **microbes**, are very small organisms; many types of microbes are too small to see without a microscope, although some parasites and fungi are visible to the naked eye.

Humans have been living with—and using—microorganisms for much longer than they have been able to see them. Historical evidence suggests that humans have had some notion of microbial life since prehistoric times and have used that knowledge to develop foods as well as prevent and treat disease. In this section, we will explore some of the historical applications of microbiology as well as the early beginnings of microbiology as a science.

### Fermented Foods and Beverages

People across the world have enjoyed fermented foods and beverages like beer, wine, bread, yogurt, cheese, and pickled vegetables for all of recorded history. Discoveries from several archeological sites suggest that even prehistoric people took advantage of fermentation to preserve and enhance the taste of food. Archaeologists studying pottery jars from a Neolithic village in China found that people were making a fermented beverage from rice, honey, and fruit as early as 7000 BC.<sup>[2]</sup>

Production of these foods and beverages requires microbial fermentation, a process that uses bacteria, mold, or yeast to convert sugars (carbohydrates) to alcohol, gases, and organic acids (**Figure 1.3**). While it is likely that people first learned about fermentation by accident—perhaps by drinking old milk that had curdled or old grape juice that had fermented—they later learned to harness the power of fermentation to make products like bread, cheese, and wine.



Yeast fermentation yields ethanol and CO<sub>2</sub>.

**Figure 1.3** A microscopic view of *Saccharomyces cerevisiae*, the yeast responsible for making bread rise (left). Yeast is a microorganism. Its cells metabolize the carbohydrates in flour (middle) and produce carbon dioxide, which causes the bread to rise (right). (credit middle: modification of work by Janus Sandsgaard; credit right: modification of work by “MDreibelbis”/Flickr)

2. P.E. McGovern et al. “Fermented Beverages of Pre- and Proto-Historic China.” *Proceedings of the National Academy of Sciences of the United States of America* 1 no. 51 (2004):17593–17598. doi:10.1073/pnas.0407921102.

## Clinical Focus

### Part 1


Cora, a 41-year-old lawyer and mother of two, has recently been experiencing severe headaches, a high fever, and a stiff neck. Her husband, who has accompanied Cora to see a doctor, reports that Cora also seems confused at times and unusually drowsy. Based on these symptoms, the doctor suspects that Cora may have meningitis, a potentially life-threatening infection of the tissue that surrounds the brain and spinal cord.

Meningitis has several potential causes. It can be brought on by bacteria, fungi, viruses, or even a reaction to medication or exposure to heavy metals. Although people with viral meningitis usually heal on their own, bacterial and fungal meningitis are quite serious and require treatment.

Cora's doctor orders a lumbar puncture (spinal tap) to take three samples of cerebrospinal fluid (CSF) from around the spinal cord (**Figure 1.2**). The samples will be sent to laboratories in three different departments for testing: clinical chemistry, microbiology, and hematology. The samples will first be visually examined to determine whether the CSF is abnormally colored or cloudy; then the CSF will be examined under a microscope to see if it contains a normal number of red and white blood cells and to check for any abnormal cell types. In the microbiology lab, the specimen will be centrifuged to concentrate any cells in a sediment; this sediment will be smeared on a slide and stained with a Gram stain. Gram staining is a procedure used to differentiate between two different types of bacteria (gram-positive and gram-negative).

About 80% of patients with bacterial meningitis will show bacteria in their CSF with a Gram stain.<sup>[3]</sup> Cora's Gram stain did not show any bacteria, but her doctor decides to prescribe her antibiotics just in case. Part of the CSF sample will be cultured—put in special dishes to see if bacteria or fungi will grow. It takes some time for most microorganisms to reproduce in sufficient quantities to be detected and analyzed.

- What types of microorganisms would be killed by antibiotic treatment?



**Figure 1.2** (a) A lumbar puncture is used to take a sample of a patient's cerebrospinal fluid (CSF) for testing. A needle is inserted between two vertebrae of the lower back, called the lumbar region. (b) CSF should be clear, as in this sample. Abnormally cloudy CSF may indicate an infection but must be tested further to confirm the presence of microorganisms. (credit b: modification of work by James Heilman)

Jump to the [next](http://cnx.org/content/m58782/1.2/#) (<http://cnx.org/content/m58782/1.2/#>) Clinical Focus box.

## The Iceman Treateth

Prehistoric humans had a very limited understanding of the causes of disease, and various cultures developed different beliefs and explanations. While many believed that illness was punishment for angering the gods or was simply the result of fate, archaeological evidence suggests that prehistoric people attempted to treat illnesses and infections. One example of this is Ötzi the Iceman, a 5300-year-old mummy found frozen in the ice of the Ötztal Alps on the Austrian-Italian border in 1991. Because Ötzi was so well preserved by the ice, researchers discovered that he was infected with the eggs of the parasite *Trichuris trichiura*, which may have caused him to have abdominal pain and anemia. Researchers also found evidence of *Borrelia burgdorferi*, a bacterium that causes Lyme disease.<sup>[4]</sup> Some researchers think Ötzi may have been trying to treat his infections with the woody fruit of the *Piptoporus betulinus* fungus, which was discovered tied to his belongings.<sup>[5]</sup> This fungus has both laxative and antibiotic properties. Ötzi was also covered in tattoos that were made by cutting incisions into his skin, filling them with herbs, and then burning the herbs.<sup>[6]</sup> There is speculation that this may have been another attempt to treat his health ailments.

## Early Notions of Disease, Contagion, and Containment

Several ancient civilizations appear to have had some understanding that disease could be transmitted by things they could not see. This is especially evident in historical attempts to contain the spread of disease. For example, the Bible refers to the practice of quarantining people with leprosy and other diseases, suggesting that people understood that diseases could be communicable. Ironically, while leprosy is communicable, it is also a disease that progresses slowly. This means that people were likely quarantined after they had already spread the disease to others.

The ancient Greeks attributed disease to bad air, *mal'aria*, which they called “miasmatic odors.” They developed hygiene practices that built on this idea. The Romans also believed in the miasma hypothesis and created a complex sanitation infrastructure to deal with sewage. In Rome, they built aqueducts, which brought fresh water into the city, and a giant sewer, the *Cloaca Maxima*, which carried waste away and into the river Tiber (**Figure 1.4**). Some researchers believe that this infrastructure helped protect the Romans from epidemics of waterborne illnesses.

---


3. Rebecca Buxton. “Examination of Gram Stains of Spinal Fluid—Bacterial Meningitis.” *American Society for Microbiology*. 2007. <http://www.microbelibrary.org/library/gram-stain/3065-examination-of-gram-stains-of-spinal-fluid-bacterial-meningitis>

4. A. Keller et al. “New Insights into the Tyrolean Iceman's Origin and Phenotype as Inferred by Whole-Genome Sequencing.” *Nature Communications*, 3 (2012): 698. doi:10.1038/ncomms1701.

5. L. Capasso. “5300 Years Ago, the Ice Man Used Natural Laxatives and Antibiotics.” *The Lancet*, 352 (1998) 9143: 1864. doi: 10.1016/s0140-6736(05)79939-6.

6. L. Capasso, L. “5300 Years Ago, the Ice Man Used Natural Laxatives and Antibiotics.” *The Lancet*, 352 no. 9143 (1998): 1864. doi: 10.1016/s0140-6736(05)79939-6.





**Figure 1.4** (a) The *Cloaca Maxima*, or “Greatest Sewer” (shown in red), ran through ancient Rome. It was an engineering marvel that carried waste away from the city and into the river Tiber. (b) These ancient latrines emptied into the *Cloaca Maxima*.


Even before the invention of the microscope, some doctors, philosophers, and scientists made great strides in understanding the invisible forces—what we now know as microbes—that can cause infection, disease, and death.

The Greek physician Hippocrates (460–370 BC) is considered the “father of Western medicine” (Figure 1.5). Unlike many of his ancestors and contemporaries, he dismissed the idea that disease was caused by supernatural forces. Instead, he posited that diseases had natural causes from within patients or their environments. Hippocrates and his heirs are believed to have written the *Hippocratic Corpus*, a collection of texts that make up some of the oldest surviving medical books.<sup>[7]</sup> Hippocrates is also often credited as the author of the Hippocratic Oath, taken by new physicians to pledge their dedication to diagnosing and treating patients without causing harm.

7. G. Pappas et al. “Insights Into Infectious Disease in the Era of Hippocrates.” *International Journal of Infectious Diseases* 12 (2008) 4:347–350. doi: <http://dx.doi.org/10.1016/j.ijid.2007.11.003>.

While Hippocrates is considered the father of Western medicine, the Greek philosopher and historian Thucydides (460–395 BC) is considered the father of scientific history because he advocated for evidence-based analysis of cause-and-effect reasoning (Figure 1.5). Among his most important contributions are his observations regarding the Athenian plague that killed one-third of the population of Athens between 430 and 410 BC. Having survived the epidemic himself, Thucydides made the important observation that survivors did not get re-infected with the disease, even when taking care of actively sick people.<sup>[8]</sup> This observation shows an early understanding of the concept of immunity.

Marcus Terentius Varro (116–27 BC) was a prolific Roman writer who was one of the first people to propose the concept that things we cannot see (what we now call microorganisms) can cause disease (Figure 1.5). In *Res Rusticae* (*On Farming*), published in 36 BC, he said that “precautions must also be taken in neighborhood swamps . . . because certain minute creatures [*animalia minuta*] grow there which cannot be seen by the eye, which float in the air and enter the body through the mouth and nose and there cause serious diseases.”<sup>[9]</sup>



**Figure 1.5** (a) Hippocrates, the “father of Western medicine,” believed that diseases had natural, not supernatural, causes. (b) The historian Thucydides observed that survivors of the Athenian plague were subsequently immune to the infection. (c) Marcus Terentius Varro proposed that disease could be caused by “certain minute creatures . . . which cannot be seen by the eye.” (credit c: modification of work by Alessandro Antonelli)



### Check Your Understanding

- Give two examples of foods that have historically been produced by humans with the aid of microbes.
- Explain how historical understandings of disease contributed to attempts to treat and contain disease.

## The Birth of Microbiology

While the ancients may have suspected the existence of invisible “minute creatures,” it wasn’t until the invention of the microscope that their existence was definitively confirmed. While it is unclear who exactly invented the microscope, a Dutch cloth merchant named Antonie van Leeuwenhoek (1632–1723) was the first to develop a lens powerful enough to view microbes. In 1675, using a simple but powerful microscope, Leeuwenhoek was able to observe single-celled organisms, which he described as “animalcules” or “wee little beasties,” swimming in a drop

8. Thucydides. *The History of the Peloponnesian War. The Second Book*. 431 BC. Translated by Richard Crawley. <http://classics.mit.edu/Thucydides/pelopwar.2.second.html>.

9. Plinio Prioreschi. *A History of Medicine: Roman Medicine*. Lewiston, NY: Edwin Mellen Press, 1998: p. 215.




of rain water. From his drawings of these little organisms, we now know he was looking at bacteria and protists. (We will explore Leeuwenhoek's contributions to microscopy further in **Chapter 2**.)

Nearly 200 years after van Leeuwenhoek got his first glimpse of microbes, the “Golden Age of Microbiology” spawned a host of new discoveries between 1857 and 1914. Two famous microbiologists, Louis Pasteur and Robert Koch, were especially active in advancing our understanding of the unseen world of microbes (**Figure 1.6**). Pasteur, a French chemist, showed that individual microbial strains had unique properties and demonstrated that fermentation is caused by microorganisms. He also invented pasteurization, a process used to kill microorganisms responsible for spoilage, and developed vaccines for the treatment of diseases, including rabies, in animals and humans. Koch, a German physician, was the first to demonstrate the connection between a single, isolated microbe and a known human disease. For example, he discovered the bacteria that cause anthrax (*Bacillus anthracis*), cholera (*Vibrio cholera*), and tuberculosis (*Mycobacterium tuberculosis*).<sup>[10]</sup> We will discuss these famous microbiologists, and others, in later chapters.



(a)



(b)

**Figure 1.6** (a) Louis Pasteur (1822–1895) is credited with numerous innovations that advanced the fields of microbiology and immunology. (b) Robert Koch (1843–1910) identified the specific microbes that cause anthrax, cholera, and tuberculosis.

As microbiology has developed, it has allowed the broader discipline of biology to grow and flourish in previously unimagined ways. Much of what we know about human cells comes from our understanding of microbes, and many of the tools we use today to study cells and their genetics derive from work with microbes.



### Check Your Understanding


- How did the discovery of microbes change human understanding of disease?

## Micro Connections

### Microbiology Toolbox

Because individual microbes are generally too small to be seen with the naked eye, the science of microbiology is dependent on technology that can artificially enhance the capacity of our natural senses of perception. Early microbiologists like Pasteur and Koch had fewer tools at their disposal than are found in modern laboratories, making their discoveries and innovations that much more impressive. Later chapters of this text will explore many applications of technology in depth, but for now, here is a brief overview of some of the fundamental tools of the microbiology lab.

- **Microscopes** produce magnified images of microorganisms, human cells and tissues, and many other types of specimens too small to be observed with the naked eye.
- **Stains and dyes** are used to add color to microbes so they can be better observed under a microscope. Some dyes can be used on living microbes, whereas others require that the specimens be fixed with chemicals or heat before staining. Some stains only work on certain types of microbes because of differences in their cellular chemical composition.
- **Growth media** are used to grow microorganisms in a lab setting. Some media are liquids; others are more solid or gel-like. A growth medium provides nutrients, including water, various salts, a source of carbon (like glucose), and a source of nitrogen and amino acids (like yeast extract) so microorganisms can grow and reproduce. Ingredients in a growth medium can be modified to grow unique types of microorganisms.
- A **Petri dish** is a flat-lidded dish that is typically 10–11 centimeters (cm) in diameter and 1–1.5 cm high. Petri dishes made out of either plastic or glass are used to hold growth media (**Figure 1.7**).
- **Test tubes** are cylindrical plastic or glass tubes with rounded bottoms and open tops. They can be used to grow microbes in broth, or semisolid or solid growth media.
- A **Bunsen burner** is a metal apparatus that creates a flame that can be used to sterilize pieces of equipment. A rubber tube carries gas (fuel) to the burner. In many labs, Bunsen burners are being phased out in favor of infrared **microincinerators**, which serve a similar purpose without the safety risks of an open flame.
- An **inoculation loop** is a handheld tool that ends in a small wire loop (**Figure 1.7**). The loop can be used to streak microorganisms on agar in a Petri dish or to transfer them from one test tube to another. Before each use, the inoculation loop must be sterilized so cultures do not become contaminated.



**Figure 1.7** (a) This Petri dish filled with agar has been streaked with *Legionella*, the bacterium responsible for causing Legionnaire's disease. (b) An inoculation loop like this one can be used to streak bacteria on agar in a Petri dish. (credit a: modification of work by Centers for Disease Control and Prevention; credit b: modification of work by Jeffrey M. Vinocur)

## 1.2 A Systematic Approach

### Learning Objectives

- Describe how microorganisms are classified and distinguished as unique species
- Compare historical and current systems of taxonomy used to classify microorganisms

Once microbes became visible to humans with the help of microscopes, scientists began to realize their enormous diversity. Microorganisms vary in all sorts of ways, including their size, their appearance, and their rates of reproduction. To study this incredibly diverse new array of organisms, researchers needed a way to systematically organize them.

### The Science of Taxonomy

**Taxonomy** is the classification, description, identification, and naming of living organisms. Classification is the practice of organizing organisms into different groups based on their shared characteristics. The most famous early taxonomist was a Swedish botanist, zoologist, and physician named Carolus Linnaeus (1701–1778). In 1735, Linnaeus published *Systema Naturae*, an 11-page booklet in which he proposed the Linnaean taxonomy, a system of categorizing and naming organisms using a standard format so scientists could discuss organisms using consistent terminology. He continued to revise and add to the book, which grew into multiple volumes (**Figure 1.8**).




**Figure 1.8** Swedish botanist, zoologist, and physician Carolus Linnaeus developed a new system for categorizing plants and animals. In this 1853 portrait by Hendrik Hollander, Linnaeus is holding a twinflower, named *Linnaea borealis* in his honor.

In his taxonomy, Linnaeus divided the natural world into three kingdoms: animal, plant, and mineral (the mineral kingdom was later abandoned). Within the animal and plant kingdoms, he grouped organisms using a hierarchy of increasingly specific levels and sublevels based on their similarities. The names of the levels in Linnaeus's original taxonomy were kingdom, class, order, family, genus (plural: genera), and species. Species was, and continues to be, the most specific and basic taxonomic unit.

### Evolving Trees of Life (Phylogenies)

With advances in technology, other scientists gradually made refinements to the Linnaean system and eventually created new systems for classifying organisms. In the 1800s, there was a growing interest in developing taxonomies that took into account the evolutionary relationships, or **phylogenies**, of all different species of organisms on earth. One way to depict these relationships is via a diagram called a phylogenetic tree (or tree of life). In these diagrams, groups of organisms are arranged by how closely related they are thought to be. In early phylogenetic trees, the relatedness of organisms was inferred by their visible similarities, such as the presence or absence of hair or the number of limbs. Now, the analysis is more complicated. Today, phylogenetic analyses include genetic, biochemical, and embryological comparisons, as will be discussed later in this chapter.

Linnaeus's tree of life contained just two main branches for all living things: the animal and plant kingdoms. In 1866, Ernst Haeckel, a German biologist, philosopher, and physician, proposed another kingdom, Protista, for unicellular organisms (**Figure 1.9**). He later proposed a fourth kingdom, Monera, for unicellular organisms whose cells lack nuclei, like bacteria.




**Figure 1.9** Ernst Haeckel's rendering of the tree of life, from his 1866 book *General Morphology of Organisms*, contained three kingdoms: Plantae, Protista, and Animalia. He later added a fourth kingdom, Monera, for unicellular organisms lacking a nucleus.

Nearly 100 years later, in 1969, American ecologist Robert Whittaker (1920–1980) proposed adding another kingdom—Fungi—in his tree of life. Whittaker's tree also contained a level of categorization above the kingdom level—the empire or superkingdom level—to distinguish between organisms that have membrane-bound nuclei in their cells (**eukaryotes**) and those that do not (**prokaryotes**). Empire Prokaryota contained just the Kingdom Monera. The Empire Eukaryota contained the other four kingdoms: Fungi, Protista, Plantae, and Animalia. Whittaker's five-kingdom tree was considered the standard phylogeny for many years.

**Figure 1.10** shows how the tree of life has changed over time. Note that viruses are not found in any of these trees. That is because they are not made up of cells and thus it is difficult to determine where they would fit into a tree of life.





**Figure 1.10** This timeline shows how the shape of the tree of life has changed over the centuries. Even today, the taxonomy of living organisms is continually being reevaluated and refined with advances in technology.



## Check Your Understanding

- Briefly summarize how our evolving understanding of microorganisms has contributed to changes in the way that organisms are classified.

## Clinical Focus

### Part 2

Antibiotic drugs are specifically designed to kill or inhibit the growth of bacteria. But after a couple of days on antibiotics, Cora shows no signs of improvement. Also, her CSF cultures came back from the lab negative. Since bacteria or fungi were not isolated from Cora's CSF sample, her doctor rules out bacterial and fungal meningitis. Viral meningitis is still a possibility.

However, Cora now reports some troubling new symptoms. She is starting to have difficulty walking. Her muscle stiffness has spread from her neck to the rest of her body, and her limbs sometimes jerk involuntarily. In addition, Cora's cognitive symptoms are worsening. At this point, Cora's doctor becomes very concerned and orders more tests on the CSF samples.


- What types of microorganisms could be causing Cora's symptoms?

Jump to the **next** Clinical Focus box. Go back to the **previous** Clinical Focus box.

## The Role of Genetics in Modern Taxonomy

Haeckel's and Whittaker's trees presented hypotheses about the phylogeny of different organisms based on readily observable characteristics. But the advent of molecular genetics in the late 20th century revealed other ways to organize phylogenetic trees. Genetic methods allow for a standardized way to compare all living organisms without relying on observable characteristics that can often be subjective. Modern taxonomy relies heavily on comparing the nucleic acids (deoxyribonucleic acid [DNA] or ribonucleic acid [RNA]) or proteins from different organisms. The more similar the nucleic acids and proteins are between two organisms, the more closely related they are considered to be.

In the 1970s, American microbiologist Carl Woese discovered what appeared to be a “living record” of the evolution of organisms. He and his collaborator George Fox created a genetics-based tree of life based on similarities and differences they observed in the small subunit ribosomal RNA (rRNA) of different organisms. In the process, they discovered that a certain type of bacteria, called archaebacteria (now known simply as archaea), were significantly different from other bacteria and eukaryotes in terms of the sequence of small subunit rRNA. To accommodate this difference, they created a tree with three Domains above the level of Kingdom: Archaea, Bacteria, and Eukarya (**Figure 1.11**). Genetic analysis of the small subunit rRNA suggests archaea, bacteria, and eukaryotes all evolved from a common ancestral cell type. The tree is skewed to show a closer evolutionary relationship between Archaea and Eukarya than they have to Bacteria.



**Figure 1.11** Woese and Fox's phylogenetic tree contains three domains: Bacteria, Archaea, and Eukarya. Domains Archaea and Bacteria contain all prokaryotic organisms, and Eukarya contains all eukaryotic organisms.

Scientists continue to use analysis of RNA, DNA, and proteins to determine how organisms are related. One interesting, and complicating, discovery is that of horizontal gene transfer—when a gene of one species is absorbed into another organism's genome. Horizontal gene transfer is especially common in microorganisms and can make it difficult to determine how organisms are evolutionarily related. Consequently, some scientists now think in terms of “webs of life” rather than “trees of life.”



## Check Your Understanding

- In modern taxonomy, how do scientists determine how closely two organisms are related?
- Explain why the branches on the “tree of life” all originate from a single “trunk.”

### Naming Microbes

In developing his taxonomy, Linnaeus used a system of **binomial nomenclature**, a two-word naming system for identifying organisms by genus and species. For example, modern humans are in the genus *Homo* and have the species name *sapiens*, so their scientific name in binomial nomenclature is *Homo sapiens*. In binomial nomenclature, the genus part of the name is always capitalized; it is followed by the species name, which is not capitalized. Both names are italicized.

Taxonomic names in the 18th through 20th centuries were typically derived from Latin, since that was the common language used by scientists when taxonomic systems were first created. Today, newly discovered organisms can be given names derived from Latin, Greek, or English. Sometimes these names reflect some distinctive trait of the organism; in other cases, microorganisms are named after the scientists who discovered them. The archaeon *Haloquadratum walsbyi* is an example of both of these naming schemes. The genus, *Haloquadratum*, describes the microorganism’s saltwater habitat (*halo* is derived from the Greek word for “salt”) as well as the arrangement of its square cells, which are arranged in square clusters of four cells (*quadratum* is Latin for “foursquare”). The species, *walsbyi*, is named after Anthony Edward Walsby, the microbiologist who discovered *Haloquadratum walsbyi* in 1980. While it might seem easier to give an organism a common descriptive name—like a red-headed woodpecker—we can imagine how that could become problematic. What happens when another species of woodpecker with red head coloring is discovered? The systematic nomenclature scientists use eliminates this potential problem by assigning each organism a single, unique two-word name that is recognized by scientists all over the world.

In this text, we will typically abbreviate an organism’s genus and species after its first mention. The abbreviated form is simply the first initial of the genus, followed by a period and the full name of the species. For example, the bacterium *Escherichia coli* is shortened to *E. coli* in its abbreviated form. You will encounter this same convention in other scientific texts as well.

### Bergey’s Manuals

Whether in a tree or a web, microbes can be difficult to identify and classify. Without easily observable macroscopic features like feathers, feet, or fur, scientists must capture, grow, and devise ways to study their biochemical properties to differentiate and classify microbes. Despite these hurdles, a group of microbiologists created and updated a set of manuals for identifying and classifying microorganisms. First published in 1923 and since updated many times, *Bergey’s Manual of Determinative Bacteriology* and *Bergey’s Manual of Systematic Bacteriology* are the standard references for identifying and classifying different prokaryotes. (**Appendix D** (<http://cnx.org/content/m58949/latest/>) of this textbook is partly based on Bergey’s manuals; it shows how the organisms that appear in this textbook are classified.) Because so many bacteria look identical, methods based on nonvisual characteristics must be used to identify them. For example, biochemical tests can be used to identify chemicals unique to certain species. Likewise, serological tests can be used to identify specific antibodies that will react against the proteins found in certain species. Ultimately, DNA and rRNA sequencing can be used both for identifying a particular bacterial species and for classifying newly discovered species.





## Check Your Understanding

- What is binomial nomenclature and why is it a useful tool for naming organisms?
- Explain why a resource like one of Bergey's manuals would be helpful in identifying a microorganism in a sample.

## Micro Connections

### Same Name, Different Strain

Within one species of microorganism, there can be several subtypes called strains. While different strains may be nearly identical genetically, they can have very different attributes. The bacterium *Escherichia coli* is infamous for causing food poisoning and traveler's diarrhea. However, there are actually many different strains of *E. coli*, and they vary in their ability to cause disease.

One pathogenic (disease-causing) *E. coli* strain that you may have heard of is *E. coli* O157:H7. In humans, infection from *E. coli* O157:H7 can cause abdominal cramps and diarrhea. Infection usually originates from contaminated water or food, particularly raw vegetables and undercooked meat. In the 1990s, there were several large outbreaks of *E. coli* O157:H7 thought to have originated in undercooked hamburgers.

While *E. coli* O157:H7 and some other strains have given *E. coli* a bad name, most *E. coli* strains do not cause disease. In fact, some can be helpful. Different strains of *E. coli* found naturally in our gut help us digest our food, provide us with some needed chemicals, and fight against pathogenic microbes.

## Link to Learning




Learn more about phylogenetic trees by exploring the Wellcome Trust's interactive Tree of Life. The [website \(http://www.openstaxcollege.org//22wellcome\)](http://www.openstaxcollege.org//22wellcome) contains information, photos, and animations about many different organisms. Select two organisms to see how they are evolutionarily related.

## 1.3 Types of Microorganisms

### Learning Objectives

- List the various types of microorganisms and describe their defining characteristics
- Give examples of different types of cellular and viral microorganisms and infectious agents
- Describe the similarities and differences between archaea and bacteria
- Provide an overview of the field of microbiology

Most microbes are unicellular and small enough that they require artificial magnification to be seen. However, there are some unicellular microbes that are visible to the naked eye, and some multicellular organisms that are microscopic. An object must measure about 100 micrometers ( $\mu\text{m}$ ) to be visible without a microscope, but most microorganisms are many times smaller than that. For some perspective, consider that a typical animal cell measures roughly  $10\ \mu\text{m}$  across but is still microscopic. Bacterial cells are typically about  $1\ \mu\text{m}$ , and viruses can be 10 times smaller than bacteria (**Figure 1.12**). See **Table 1.1** for units of length used in microbiology.



**Figure 1.12** The relative sizes of various microscopic and nonmicroscopic objects. Note that a typical virus measures about 100 nm, 10 times smaller than a typical bacterium (~1 μm), which is at least 10 times smaller than a typical plant or animal cell (~10–100 μm). An object must measure about 100 μm to be visible without a microscope.

### Units of Length Commonly Used in Microbiology

Metric Unit	Meaning of Prefix	Metric Equivalent
meter (m)	—	1 m = 10 <sup>0</sup> m
decimeter (dm)	1/10	1 dm = 0.1 m = 10 <sup>-1</sup> m
centimeter (cm)	1/100	1 cm = 0.01 m = 10 <sup>-2</sup> m
millimeter (mm)	1/1000	1 mm = 0.001 m = 10 <sup>-3</sup> m
micrometer (μm)	1/1,000,000	1 μm = 0.000001 m = 10 <sup>-6</sup> m
nanometer (nm)	1/1,000,000,000	1 nm = 0.000000001 m = 10 <sup>-9</sup> m

**Table 1.1**

Microorganisms differ from each other not only in size, but also in structure, habitat, metabolism, and many other characteristics. While we typically think of microorganisms as being unicellular, there are also many multicellular organisms that are too small to be seen without a microscope. Some microbes, such as viruses, are even **acellular** (not composed of cells).

Microorganisms are found in each of the three domains of life: Archaea, Bacteria, and Eukarya. Microbes within the domains Bacteria and Archaea are all prokaryotes (their cells lack a nucleus), whereas microbes in the domain Eukarya are eukaryotes (their cells have a nucleus). Some microorganisms, such as viruses, do not fall within any of the three domains of life. In this section, we will briefly introduce each of the broad groups of microbes. Later chapters will go into greater depth about the diverse species within each group.

## Link to Learning




How big is a bacterium or a virus compared to other objects? Check out this [interactive website \(http://www.openstaxcollege.org//22relsizes\)](http://www.openstaxcollege.org//22relsizes) to get a feel for the scale of different microorganisms.

## Prokaryotic Microorganisms

**Bacteria** are found in nearly every habitat on earth, including within and on humans. Most bacteria are harmless or helpful, but some are **pathogens**, causing disease in humans and other animals. Bacteria are prokaryotic because their genetic material (DNA) is not housed within a true nucleus. Most bacteria have cell walls that contain peptidoglycan.

Bacteria are often described in terms of their general shape. Common shapes include spherical (coccus), rod-shaped (bacillus), or curved (spirillum, spirochete, or vibrio). **Figure 1.13** shows examples of these shapes.



**Figure 1.13** Common bacterial shapes. Note how coccobacillus is a combination of spherical (coccus) and rod-shaped (bacillus). (credit “Coccus”: modification of work by Janice Haney Carr, Dr. Richard Facklam, Centers for Disease Control and Prevention; credit “Bacillus”: modification of work by “Elapied”/Wikimedia Commons)

They have a wide range of metabolic capabilities and can grow in a variety of environments, using different combinations of nutrients. Some bacteria are photosynthetic, such as oxygenic cyanobacteria and anoxygenic green sulfur and green nonsulfur bacteria; these bacteria use energy derived from sunlight, and fix carbon dioxide for growth. Other types of bacteria are nonphotosynthetic, obtaining their energy from organic or inorganic compounds in their environment.

**Archaea** are also unicellular prokaryotic organisms. Archaea and bacteria have different evolutionary histories, as well as significant differences in genetics, metabolic pathways, and the composition of their cell walls and membranes. Unlike most bacteria, archaeal cell walls do not contain peptidoglycan, but their cell walls are often composed of a similar substance called pseudopeptidoglycan. Like bacteria, archaea are found in nearly every habitat on earth, even extreme environments that are very cold, very hot, very basic, or very acidic (**Figure 1.14**). Some archaea live in the human body, but none have been shown to be human pathogens.



**Figure 1.14** Some archaea live in extreme environments, such as the Morning Glory pool, a hot spring in Yellowstone National Park. The color differences in the pool result from the different communities of microbes that are able to thrive at various water temperatures.



### Check Your Understanding

- What are the two main types of prokaryotic organisms?
- Name some of the defining characteristics of each type.


## Eukaryotic Microorganisms

The domain Eukarya contains all eukaryotes, including uni- or multicellular eukaryotes such as protists, fungi, plants, and animals. The major defining characteristic of eukaryotes is that their cells contain a nucleus.

### Protists

**Protists** are unicellular eukaryotes that are not plants, animals, or fungi. Algae and protozoa are examples of protists.

**Algae** (singular: alga) are plant-like protists that can be either unicellular or multicellular (**Figure 1.15**). Their cells are surrounded by cell walls made of cellulose, a type of carbohydrate. Algae are photosynthetic organisms that extract energy from the sun and release oxygen and carbohydrates into their environment. Because other organisms can use their waste products for energy, algae are important parts of many ecosystems. Many consumer products contain ingredients derived from algae, such as carrageenan or alginic acid, which are found in some brands of ice cream, salad dressing, beverages, lipstick, and toothpaste. A derivative of algae also plays a prominent role in the microbiology laboratory. Agar, a gel derived from algae, can be mixed with various nutrients and used to grow microorganisms in a Petri dish. Algae are also being developed as a possible source for biofuels.



**Figure 1.15** Assorted diatoms, a kind of algae, live in annual sea ice in McMurdo Sound, Antarctica. Diatoms range in size from 2  $\mu\text{m}$  to 200  $\mu\text{m}$  and are visualized here using light microscopy. (credit: National Oceanic and Atmospheric Administration)

**Protozoa** (singular: protozoan) are protists that make up the backbone of many food webs by providing nutrients for other organisms. Protozoa are very diverse. Some protozoa move with help from hair-like structures called cilia or whip-like structures called flagella. Others extend part of their cell membrane and cytoplasm to propel themselves forward. These cytoplasmic extensions are called pseudopods (“false feet”). Some protozoa are photosynthetic; others feed on organic material. Some are free-living, whereas others are parasitic, only able to survive by extracting nutrients from a host organism. Most protozoa are harmless, but some are pathogens that can cause disease in animals or humans (**Figure 1.16**).



**Figure 1.16** *Giardia lamblia*, an intestinal protozoan parasite that infects humans and other mammals, causing severe diarrhea. (credit: modification of work by Centers for Disease Control and Prevention)


## Fungi

**Fungi** (singular: fungus) are also eukaryotes. Some multicellular fungi, such as mushrooms, resemble plants, but they are actually quite different. Fungi are not photosynthetic, and their cell walls are usually made out of chitin rather than cellulose.

Unicellular fungi—yeasts—are included within the study of microbiology. There are more than 1000 known species. Yeasts are found in many different environments, from the deep sea to the human navel. Some yeasts have beneficial



uses, such as causing bread to rise and beverages to ferment; but yeasts can also cause food to spoil. Some even cause diseases, such as vaginal yeast infections and oral thrush (**Figure 1.17**).



**Figure 1.17** *Candida albicans* is a unicellular fungus, or yeast. It is the causative agent of vaginal yeast infections as well as oral thrush, a yeast infection of the mouth that commonly afflicts infants. *C. albicans* has a morphology similar to that of coccus bacteria; however, yeast is a eukaryotic organism (note the nuclei) and is much larger. (credit: modification of work by Centers for Disease Control and Prevention)

Other fungi of interest to microbiologists are multicellular organisms called **molds**. Molds are made up of long filaments that form visible colonies (**Figure 1.18**). Molds are found in many different environments, from soil to rotting food to dank bathroom corners. Molds play a critical role in the decomposition of dead plants and animals. Some molds can cause allergies, and others produce disease-causing metabolites called mycotoxins. Molds have been used to make pharmaceuticals, including penicillin, which is one of the most commonly prescribed antibiotics, and cyclosporine, used to prevent organ rejection following a transplant.



**Figure 1.18** Large colonies of microscopic fungi can often be observed with the naked eye, as seen on the surface of these moldy oranges.




### Check Your Understanding

- Name two types of protists and two types of fungi.

- Name some of the defining characteristics of each type.

## Helminths

Multicellular parasitic worms called **helminths** are not technically microorganisms, as most are large enough to see without a microscope. However, these worms fall within the field of microbiology because diseases caused by helminths involve microscopic eggs and larvae. One example of a helminth is the guinea worm, or *Dracunculus medinensis*, which causes dizziness, vomiting, diarrhea, and painful ulcers on the legs and feet when the worm works its way out of the skin (**Figure 1.19**). Infection typically occurs after a person drinks water containing water fleas infected by guinea-worm larvae. In the mid-1980s, there were an estimated 3.5 million cases of guinea-worm disease, but the disease has been largely eradicated. In 2014, there were only 126 cases reported, thanks to the coordinated efforts of the World Health Organization (WHO) and other groups committed to improvements in drinking water sanitation.<sup>[11][12]</sup>



**Figure 1.19** (a) The beef tapeworm, *Taenia saginata*, infects both cattle and humans. *T. saginata* eggs are microscopic (around 50  $\mu\text{m}$ ), but adult worms like the one shown here can reach 4–10 m, taking up residence in the digestive system. (b) An adult guinea worm, *Dracunculus medinensis*, is removed through a lesion in the patient's skin by winding it around a matchstick. (credit b: modification of work by Centers for Disease Control and Prevention)


## Viruses

**Viruses** are **acellular** microorganisms, which means they are not composed of cells. Essentially, a virus consists of proteins and genetic material—either DNA or RNA, but never both—that are inert outside of a host organism. However, by incorporating themselves into a host cell, viruses are able to co-opt the host's cellular mechanisms to multiply and infect other hosts.

Viruses can infect all types of cells, from human cells to the cells of other microorganisms. In humans, viruses are responsible for numerous diseases, from the common cold to deadly Ebola (**Figure 1.20**). However, many viruses do not cause disease.

11. C. Greenaway “Dracunculiasis (Guinea Worm Disease).” *Canadian Medical Association Journal* 170 no. 4 (2004):495–500.

12. World Health Organization. “Dracunculiasis (Guinea-Worm Disease).” *WHO*. 2015. <http://www.who.int/mediacentre/factsheets/fs359/en/>. Accessed October 2, 2015.



**Figure 1.20** (a) Members of the Coronavirus family can cause respiratory infections like the common cold, severe acute respiratory syndrome (SARS), and Middle East respiratory syndrome (MERS). Here they are viewed under a transmission electron microscope (TEM). (b) Ebolavirus, a member of the Filovirus family, as visualized using a TEM. (credit b: modification of work by Thomas W. Geisbert)



### Check Your Understanding

- Are helminths microorganisms? Explain why or why not.
- How are viruses different from other microorganisms?

## Microbiology as a Field of Study

**Microbiology** is a broad term that encompasses the study of all different types of microorganisms. But in practice, microbiologists tend to specialize in one of several subfields. For example, **bacteriology** is the study of bacteria; **mycology** is the study of fungi; **protozoology** is the study of protozoa; **parasitology** is the study of helminths and other parasites; and **virology** is the study of viruses (**Figure 1.21**). **Immunology**, the study of the immune system, is often included in the study of microbiology because host–pathogen interactions are central to our understanding of infectious disease processes. Microbiologists can also specialize in certain areas of microbiology, such as clinical microbiology, environmental microbiology, applied microbiology, or food microbiology.

In this textbook, we are primarily concerned with clinical applications of microbiology, but since the various subfields of microbiology are highly interrelated, we will often discuss applications that are not strictly clinical.





**Figure 1.21** A virologist samples eggs from this nest to be tested for the influenza A virus, which causes avian flu in birds. (credit: Don Becker)

### Eye on Ethics



#### Bioethics in Microbiology

In the 1940s, the U.S. government was looking for a solution to a medical problem: the prevalence of sexually transmitted diseases (STDs) among soldiers. Several now-infamous government-funded studies used human subjects to research common STDs and treatments. In one such study, American researchers intentionally exposed more than 1300 human subjects in Guatemala to syphilis, gonorrhea, and chancroid to determine the ability of penicillin and other antibiotics to combat these diseases. Subjects of the study included Guatemalan soldiers, prisoners, prostitutes, and psychiatric patients—none of whom were informed that they were taking part in the study. Researchers exposed subjects to STDs by various methods, from facilitating intercourse with infected prostitutes to inoculating subjects with the bacteria known to cause the diseases. This latter method involved making a small wound on the subject's genitals or elsewhere on the body, and then putting bacteria directly into the wound.<sup>[13]</sup> In 2011, a U.S. government commission tasked with investigating the experiment revealed that only some of the subjects were treated with penicillin, and 83 subjects died by 1953, likely as a result of the study.<sup>[14]</sup>

Unfortunately, this is one of many horrific examples of microbiology experiments that have violated basic ethical standards. Even if this study had led to a life-saving medical breakthrough (it did not), few would argue that its methods were ethically sound or morally justifiable. But not every case is so clear cut. Professionals working in clinical settings are frequently confronted with ethical dilemmas, such as working with patients who

decline a vaccine or life-saving blood transfusion. These are just two examples of life-and-death decisions that may intersect with the religious and philosophical beliefs of both the patient and the health-care professional.

No matter how noble the goal, microbiology studies and clinical practice must be guided by a certain set of ethical principles. Studies must be done with integrity. Patients and research subjects provide informed consent (not only agreeing to be treated or studied but demonstrating an understanding of the purpose of the study and any risks involved). Patients' rights must be respected. Procedures must be approved by an institutional review board. When working with patients, accurate record-keeping, honest communication, and confidentiality are paramount. Animals used for research must be treated humanely, and all protocols must be approved by an institutional animal care and use committee. These are just a few of the ethical principles explored in the *Eye on Ethics* boxes throughout this book.

## Clinical Focus

### Resolution

Cora's CSF samples show no signs of inflammation or infection, as would be expected with a viral infection. However, there is a high concentration of a particular protein, 14-3-3 protein, in her CSF. An electroencephalogram (EEG) of her brain function is also abnormal. The EEG resembles that of a patient with a neurodegenerative disease like Alzheimer's or Huntington's, but Cora's rapid cognitive decline is not consistent with either of these. Instead, her doctor concludes that Cora has Creutzfeldt-Jakob disease (CJD), a type of transmissible spongiform encephalopathy (TSE).

CJD is an extremely rare disease, with only about 300 cases in the United States each year. It is not caused by a bacterium, fungus, or virus, but rather by prions—which do not fit neatly into any particular category of microbe. Like viruses, prions are not found on the tree of life because they are acellular. Prions are extremely small, about one-tenth the size of a typical virus. They contain no genetic material and are composed solely of a type of abnormal protein.

CJD can have several different causes. It can be acquired through exposure to the brain or nervous-system tissue of an infected person or animal. Consuming meat from an infected animal is one way such exposure can occur. There have also been rare cases of exposure to CJD through contact with contaminated surgical equipment<sup>[15]</sup> and from cornea and growth-hormone donors who unknowingly had CJD.<sup>[16][17]</sup> In rare cases, the disease results from a specific genetic mutation that can sometimes be hereditary. However, in approximately 85% of patients with CJD, the cause of the disease is spontaneous (or sporadic) and has no identifiable cause.<sup>[18]</sup> Based on her symptoms and their rapid progression, Cora is diagnosed with sporadic CJD.

Unfortunately for Cora, CJD is a fatal disease for which there is no approved treatment. Approximately 90% of patients die within 1 year of diagnosis.<sup>[19]</sup> Her doctors focus on limiting her pain and cognitive symptoms as her disease progresses. Eight months later, Cora dies. Her CJD diagnosis is confirmed with a brain autopsy.

Go back to the *previous Clinical Focus box*.

13. Kara Rogers. "Guatemala Syphilis Experiment: American Medical Research Project". *Encyclopaedia Britannica*. <http://www.britannica.com/event/Guatemala-syphilis-experiment>. Accessed June 24, 2015.

14. Susan Donaldson James. "Syphilis Experiments Shock, But So Do Third-World Drug Trials." *ABC World News*. August 30, 2011. <http://abcnews.go.com/Health/guatemala-syphilis-experiments-shock-us-drug-trials-exploit/story?id=14414902>. Accessed June 24, 2015.

15. Greg Botelho. "Case of Creutzfeldt-Jakob Disease Confirmed in New Hampshire." *CNN*. 2013. <http://www.cnn.com/2013/09/20/health/creutzfeldt-jakob-brain-disease/>.

16. P. Rudge et al. "Iatrogenic CJD Due to Pituitary-Derived Growth Hormone With Genetically Determined Incubation Times of Up to 40 Years." *Brain* 138 no. 11 (2015): 3386–3399.

## Summary

---

### 1.1 What Our Ancestors Knew

- **Microorganisms** (or **microbes**) are living organisms that are generally too small to be seen without a microscope.
- Throughout history, humans have used microbes to make fermented foods such as beer, bread, cheese, and wine.
- Long before the invention of the microscope, some people theorized that infection and disease were spread by living things that were too small to be seen. They also correctly intuited certain principles regarding the spread of disease and immunity.
- Antonie van Leeuwenhoek, using a microscope, was the first to actually describe observations of bacteria, in 1675.
- During the Golden Age of Microbiology (1857–1914), microbiologists, including Louis Pasteur and Robert Koch, discovered many new connections between the fields of microbiology and medicine.

### 1.2 A Systematic Approach

- Carolus Linnaeus developed a taxonomic system for categorizing organisms into related groups.
- **Binomial nomenclature** assigns organisms Latinized scientific names with a genus and species designation.
- A **phylogenetic tree** is a way of showing how different organisms are thought to be related to one another from an evolutionary standpoint.
- The first phylogenetic tree contained kingdoms for plants and animals; Ernst Haeckel proposed adding kingdom for protists.
- Robert Whittaker's tree contained five kingdoms: Animalia, Plantae, Protista, Fungi, and Monera.
- Carl Woese used small subunit ribosomal RNA to create a phylogenetic tree that groups organisms into three domains based on their genetic similarity.
- Bergey's manuals of determinative and systemic bacteriology are the standard references for identifying and classifying bacteria, respectively.
- Bacteria can be identified through biochemical tests, DNA/RNA analysis, and serological testing methods.

### 1.3 Types of Microorganisms

- Microorganisms are very diverse and are found in all three domains of life: Archaea, Bacteria, and Eukarya.
- **Archaea** and **bacteria** are classified as prokaryotes because they lack a cellular nucleus. Archaea differ from bacteria in evolutionary history, genetics, metabolic pathways, and cell wall and membrane composition.
- Archaea inhabit nearly every environment on earth, but no archaea have been identified as human pathogens.
- **Eukaryotes** studied in microbiology include algae, protozoa, fungi, and helminths.
- **Algae** are plant-like organisms that can be either unicellular or multicellular, and derive energy via photosynthesis.
- **Protozoa** are unicellular organisms with complex cell structures; most are motile.
- Microscopic **fungi** include **molds** and **yeasts**.
- **Helminths** are multicellular parasitic worms. They are included in the field of microbiology because their eggs and larvae are often microscopic.
- **Viruses** are acellular microorganisms that require a host to reproduce.

---

17. J.G. Heckmann et al. "Transmission of Creutzfeldt-Jakob Disease via a Corneal Transplant." *Journal of Neurology, Neurosurgery & Psychiatry* 63 no. 3 (1997): 388–390.

18. National Institute of Neurological Disorders and Stroke. "Creutzfeldt-Jakob Disease Fact Sheet." *NIH*. 2015. [http://www.ninds.nih.gov/disorders/cjd/detail\\_cjd.htm#288133058](http://www.ninds.nih.gov/disorders/cjd/detail_cjd.htm#288133058).

19. National Institute of Neurological Disorders and Stroke. "Creutzfeldt-Jakob Disease Fact Sheet." *NIH*. 2015. [http://www.ninds.nih.gov/disorders/cjd/detail\\_cjd.htm#288133058](http://www.ninds.nih.gov/disorders/cjd/detail_cjd.htm#288133058). Accessed June 22, 2015.

- The field of microbiology is extremely broad. Microbiologists typically specialize in one of many subfields, but all health professionals need a solid foundation in clinical microbiology.

## Review Questions

### Multiple Choice

- Which of the following foods is NOT made by fermentation?
  - beer
  - bread
  - cheese
  - orange juice
- Who is considered the “father of Western medicine”?
  - Marcus Terentius Varro
  - Thucydides
  - Antonie van Leeuwenhoek
  - Hippocrates
- Who was the first to observe “animalcules” under the microscope?
  - Antonie van Leeuwenhoek
  - Ötzi the Iceman
  - Marcus Terentius Varro
  - Robert Koch
- Who proposed that swamps might harbor tiny, disease-causing animals too small to see?
  - Thucydides
  - Marcus Terentius Varro
  - Hippocrates
  - Louis Pasteur
- Which of the following was NOT a kingdom in Linnaeus’s taxonomy?
  - animal
  - mineral
  - protist
  - plant
- Which of the following is a correct usage of binomial nomenclature?
  - Homo Sapiens*
  - homo sapiens*
  - Homo sapiens*
  - Homo Sapiens*
- Which scientist proposed adding a kingdom for protists?
  - Carolus Linnaeus
  - Carl Woese
  - Robert Whittaker
  - Ernst Haeckel
- Which of the following is NOT a domain in Woese and Fox’s phylogenetic tree?
  - Plantae
  - Bacteria
  - Archaea
  - Eukarya
- Which of the following is the standard resource for identifying bacteria?
  - Systema Naturae*
  - Bergey’s *Manual of Determinative Bacteriology*
  - Woese and Fox’s phylogenetic tree
  - Haeckel’s *General Morphology of Organisms*
- Which of the following types of microorganisms is photosynthetic?
  - yeast
  - virus
  - helminth
  - alga
- Which of the following is a prokaryotic microorganism?
  - helminth
  - protozoan
  - cyanobacterium
  - mold
- Which of the following is acellular?
  - virus
  - bacterium
  - fungus
  - protozoan
- Which of the following is a type of fungal microorganism?
  - bacterium
  - protozoan
  - alga
  - yeast
- Which of the following is not a subfield of microbiology?
  - bacteriology
  - botany
  - clinical microbiology
  - virology

**Fill in the Blank**

15. Thucydides is known as the father of \_\_\_\_\_.
16. Researchers think that Ötzi the Iceman may have been infected with \_\_\_\_\_ disease.
17. The process by which microbes turn grape juice into wine is called \_\_\_\_\_.
18. In binomial nomenclature, an organism's scientific name includes its \_\_\_\_\_ and \_\_\_\_\_.
19. Whittaker proposed adding the kingdoms \_\_\_\_\_ and \_\_\_\_\_ to his phylogenetic tree.
20. \_\_\_\_\_ are organisms without membrane-bound nuclei.
21. \_\_\_\_\_ are microorganisms that are not included in phylogenetic trees because they are acellular.
22. A \_\_\_\_\_ is a disease-causing microorganism.
23. Multicellular parasitic worms studied by microbiologists are called \_\_\_\_\_.
24. The study of viruses is \_\_\_\_\_.
25. The cells of prokaryotic organisms lack a \_\_\_\_\_.


**Short Answer**

26. What did Thucydides learn by observing the Athenian plague?
27. Why was the invention of the microscope important for microbiology?
28. What are some ways people use microbes?
29. What is a phylogenetic tree?
30. Which of the five kingdoms in Whittaker's phylogenetic tree are prokaryotic, and which are eukaryotic?
31. What molecule did Woese and Fox use to construct their phylogenetic tree?
32. Name some techniques that can be used to identify and differentiate species of bacteria.
33. Describe the differences between bacteria and archaea.
34. Name three structures that various protozoa use for locomotion.
35. Describe the actual and relative sizes of a virus, a bacterium, and a plant or animal cell.

**Critical Thinking**


36. Explain how the discovery of fermented foods likely benefited our ancestors.
37. What evidence would you use to support this statement: Ancient people thought that disease was transmitted by things they could not see.
38. Why is using binomial nomenclature more useful than using common names?

39. Label the three Domains found on modern phylogenetic trees.



40. Contrast the behavior of a virus outside versus inside a cell.


41. Where would a virus, bacterium, animal cell, and a prion belong on this chart?





## Chapter 2

# How We See the Invisible World



**Figure 2.1** Different types of microscopy are used to visualize different structures. Brightfield microscopy (left) renders a darker image on a lighter background, producing a clear image of these *Bacillus anthracis* cells in cerebrospinal fluid (the rod-shaped bacterial cells are surrounded by larger white blood cells). Darkfield microscopy (right) increases contrast, rendering a brighter image on a darker background, as demonstrated by this image of the bacterium *Borrelia burgdorferi*, which causes Lyme disease. (credit right: modification of work by American Society for Microbiology)

### Chapter Outline

- 2.1 The Properties of Light
- 2.2 Peering Into the Invisible World
- 2.3 Instruments of Microscopy
- 2.4 Staining Microscopic Specimens

## Introduction

When we look at a rainbow, its colors span the full spectrum of light that the human eye can detect and differentiate. Each hue represents a different frequency of visible light, processed by our eyes and brains and rendered as red, orange, yellow, green, or one of the many other familiar colors that have always been a part of the human experience. But only recently have humans developed an understanding of the properties of light that allow us to see images in color.

Over the past several centuries, we have learned to manipulate light to peer into previously invisible worlds—those too small or too far away to be seen by the naked eye. Through a microscope, we can examine microbial cells and colonies, using various techniques to manipulate color, size, and contrast in ways that help us identify species and diagnose disease.


**Figure 2.1** illustrates how we can apply the properties of light to visualize and magnify images; but these stunning micrographs are just two examples of the numerous types of images we are now able to produce with different microscopic technologies. This chapter explores how various types of microscopes manipulate light in order to provide a window into the world of microorganisms. By understanding how various kinds of microscopes work, we can produce highly detailed images of microbes that can be useful for both research and clinical applications.

## 2.1 The Properties of Light

### Learning Objectives

- Identify and define the characteristics of electromagnetic radiation (EMR) used in microscopy
- Explain how lenses are used in microscopy to manipulate visible and ultraviolet (UV) light

Visible light consists of electromagnetic waves that behave like other waves. Hence, many of the properties of light that are relevant to microscopy can be understood in terms of light's behavior as a wave. An important property of light waves is the **wavelength**, or the distance between one peak of a wave and the next peak. The height of each peak (or depth of each trough) is called the **amplitude**. In contrast, the **frequency** of the wave is the rate of vibration of the wave, or the number of wavelengths within a specified time period (**Figure 2.2**).



**Figure 2.2** (a) The amplitude is the height of a wave, whereas the wavelength is the distance between one peak and the next. (b) These waves have different frequencies, or rates of vibration. The wave at the top has the lowest frequency, since it has the fewest peaks per unit time. The wave at the bottom has the highest frequency.

### Interactions of Light

Light waves interact with materials by being reflected, absorbed, or transmitted. **Reflection** occurs when a wave bounces off of a material. For example, a red piece of cloth may reflect red light to our eyes while absorbing other colors of light. **Absorbance** occurs when a material captures the energy of a light wave. In the case of glow-in-the-dark plastics, the energy from light can be absorbed and then later re-emitted as another form of phosphorescence. Transmission occurs when a wave travels through a material, like light through glass (the process of transmission is called **transmittance**). When a material allows a large proportion of light to be transmitted, it may do so because it

### Clinical Focus

#### Part 1

Cindy, a 17-year-old counselor at a summer sports camp, scraped her knee playing basketball 2 weeks ago. At the time, she thought it was only a minor abrasion that would heal, like many others before it. Instead, the wound began to look like an insect bite and has continued to become increasingly painful and swollen.


The camp nurse examines the lesion and observes a large amount of pus oozing from the surface. Concerned that Cindy may have developed a potentially aggressive infection, she swabs the wound to collect a sample from the infection site. Then she cleans out the pus and dresses the wound, instructing Cindy to keep the area clean and to come back the next day. When Cindy leaves, the nurse sends the sample to the closest medical lab to be analyzed under a microscope.

- What are some things we can learn about these bacteria by looking at them under a microscope?

Jump to the **next** Clinical Focus box.



is thinner, or more transparent (having more **transparency** and less **opacity**). **Figure 2.3** illustrates the difference between transparency and opacity.



**Figure 2.3** (a) A Petri dish is made of transparent plastic or glass, which allows transmission of a high proportion of light. This transparency allows us to see through the sides of the dish to view the contents. (b) This slice of an iron meteorite is opaque (i.e., it has opacity). Light is not transmitted through the material, making it impossible to see the part of the hand covered by the object. (credit a: modification of work by Umberto Salvagnin; credit b: modification of work by “Waifer X”/Flickr)

Light waves can also interact with each other by **interference**, creating complex patterns of motion. Dropping two pebbles into a puddle causes the waves on the puddle’s surface to interact, creating complex interference patterns. Light waves can interact in the same way.

In addition to interfering with each other, light waves can also interact with small objects or openings by bending or scattering. This is called **diffraction**. Diffraction is larger when the object is smaller relative to the wavelength of the light (the distance between two consecutive peaks of a light wave). Often, when waves diffract in different directions around an obstacle or opening, they will interfere with each other.




### Check Your Understanding

- If a light wave has a long wavelength, is it likely to have a low or high frequency?
- If an object is transparent, does it reflect, absorb, or transmit light?

## Lenses and Refraction

In the context of microscopy, **refraction** is perhaps the most important behavior exhibited by light waves. Refraction occurs when light waves change direction as they enter a new medium (**Figure 2.4**). Different transparent materials transmit light at different speeds; thus, light can change speed when passing from one material to another. This change in speed usually also causes a change in direction (refraction), with the degree of change dependent on the angle of the incoming light.




**Figure 2.4** (a) Refraction occurs when light passes from one medium, such as air, to another, such as glass, changing the direction of the light rays. (b) As shown in this diagram, light rays passing from one medium to another may be either refracted or reflected. (credit a: modification of work by "ajizai"/Wikimedia Commons)

The extent to which a material slows transmission speed relative to empty space is called the **refractive index** of that material. Large differences between the refractive indices of two materials will result in a large amount of refraction when light passes from one material to the other. For example, light moves much more slowly through water than through air, so light entering water from air can change direction greatly. We say that the water has a higher refractive index than air (**Figure 2.5**).



**Figure 2.5** This straight pole appears to bend at an angle as it enters the water. This optical illusion is due to the large difference between the refractive indices of air and water.

When light crosses a boundary into a material with a higher refractive index, its direction turns to be closer to perpendicular to the boundary (i.e., more toward a normal to that boundary; see **Figure 2.5**). This is the principle behind lenses. We can think of a lens as an object with a curved boundary (or a collection of prisms) that collects all of the light that strikes it and refracts it so that it all meets at a single point called the **image point (focus)**. A convex lens can be used to magnify because it can focus at closer range than the human eye, producing a larger image. Concave lenses and mirrors can also be used in microscopes to redirect the light path. **Figure 2.6** shows the **focal point** (the image point when light entering the lens is parallel) and the **focal length** (the distance to the focal point) for convex and concave lenses.



**Figure 2.6** (a) A lens is like a collection of prisms, such as the one shown here. (b) When light passes through a convex lens, it is refracted toward a focal point on the other side of the lens. The focal length is the distance to the focal point. (c) Light passing through a concave lens is refracted away from a focal point in front of the lens.

The human eye contains a lens that enables us to see images. This lens focuses the light reflecting off of objects in front of the eye onto the surface of the retina, which is like a screen in the back of the eye. Artificial lenses placed in front of the eye (contact lenses, glasses, or microscopic lenses) focus light before it is focused (again) by the lens of the eye, manipulating the image that ends up on the retina (e.g., by making it appear larger).

Images are commonly manipulated by controlling the distances between the object, the lens, and the screen, as well as the curvature of the lens. For example, for a given amount of curvature, when an object is closer to the lens, the focal points are farther from the lens. As a result, it is often necessary to manipulate these distances to create a focused image on a screen. Similarly, more curvature creates image points closer to the lens and a larger image when the image is in focus. This property is often described in terms of the focal distance, or distance to the focal point.




### Check Your Understanding

- Explain how a lens focuses light at the image point.
- Name some factors that affect the focal length of a lens.

## Electromagnetic Spectrum and Color

Visible light is just one form of electromagnetic radiation (EMR), a type of energy that is all around us. Other forms of EMR include microwaves, X-rays, and radio waves, among others. The different types of EMR fall on the electromagnetic spectrum, which is defined in terms of wavelength and frequency. The spectrum of visible light occupies a relatively small range of frequencies between infrared and ultraviolet light (**Figure 2.7**).




**Figure 2.7** The electromagnetic spectrum ranges from high-frequency gamma rays to low-frequency radio waves. Visible light is the relatively small range of electromagnetic frequencies that can be sensed by the human eye. On the electromagnetic spectrum, visible light falls between ultraviolet and infrared light. (credit: modification of work by Johannes Ahlmann)

Whereas wavelength represents the distance between adjacent peaks of a light wave, frequency, in a simplified definition, represents the rate of oscillation. Waves with higher frequencies have shorter wavelengths and, therefore, have more oscillations per unit time than lower-frequency waves. Higher-frequency waves also contain more energy than lower-frequency waves. This energy is delivered as elementary particles called photons. Higher-frequency waves deliver more energetic photons than lower-frequency waves.

Photons with different energies interact differently with the retina. In the spectrum of visible light, each color corresponds to a particular frequency and wavelength (**Figure 2.7**). The lowest frequency of visible light appears as the color red, whereas the highest appears as the color violet. When the retina receives visible light of many different frequencies, we perceive this as white light. However, white light can be separated into its component colors using refraction. If we pass white light through a prism, different colors will be refracted in different directions, creating a rainbow-like spectrum on a screen behind the prism. This separation of colors is called **dispersion**, and it occurs because, for a given material, the refractive index is different for different frequencies of light.

Certain materials can refract nonvisible forms of EMR and, in effect, transform them into visible light. Certain **fluorescent** dyes, for instance, absorb ultraviolet or blue light and then use the energy to emit photons of a different color, giving off light rather than simply vibrating. This occurs because the energy absorption causes electrons to jump to higher energy states, after which they then almost immediately fall back down to their ground states, emitting specific amounts of energy as photons. Not all of the energy is emitted in a given photon, so the emitted photons will be of lower energy and, thus, of lower frequency than the absorbed ones. Thus, a dye such as Texas red may be excited by blue light, but emit red light; or a dye such as fluorescein isothiocyanate (FITC) may absorb (invisible) high-energy ultraviolet light and emit green light (**Figure 2.8**). In some materials, the photons may be emitted following a delay after absorption; in this case, the process is called **phosphorescence**. Glow-in-the-dark plastic works by using phosphorescent material.



**Figure 2.8** The fluorescent dyes absorbed by these bovine pulmonary artery endothelial cells emit brilliant colors when excited by ultraviolet light under a fluorescence microscope. Various cell structures absorb different dyes. The nuclei are stained blue with 4',6-diamidino-2-phenylindole (DAPI); microtubules are marked green by an antibody bound to FITC; and actin filaments are labeled red with phalloidin bound to tetramethylrhodamine (TRITC).



### Check Your Understanding

- Which has a higher frequency: red light or green light?
- Explain why dispersion occurs when white light passes through a prism.
- Why do fluorescent dyes emit a different color of light than they absorb?

## Magnification, Resolution, and Contrast

Microscopes magnify images and use the properties of light to create useful images of small objects. **Magnification** is defined as the ability of a lens to enlarge the image of an object when compared to the real object. For example, a magnification of  $10\times$  means that the image appears 10 times the size of the object as viewed with the naked eye.

Greater magnification typically improves our ability to see details of small objects, but magnification alone is not sufficient to make the most useful images. It is often useful to enhance the **resolution** of objects: the ability to tell that two separate points or objects are separate. A low-resolution image appears fuzzy, whereas a high-resolution image appears sharp. Two factors affect resolution. The first is wavelength. Shorter wavelengths are able to resolve smaller objects; thus, an electron microscope has a much higher resolution than a light microscope, since it uses an electron beam with a very short wavelength, as opposed to the long-wavelength visible light used by a light microscope. The second factor that affects resolution is **numerical aperture**, which is a measure of a lens's ability to gather light. The higher the numerical aperture, the better the resolution.

### Link to Learning



Read this [article \(http://www.openstaxcollege.org//22aperture\)](http://www.openstaxcollege.org//22aperture) to learn more about factors that can increase or decrease the numerical aperture of a lens.

Even when a microscope has high resolution, it can be difficult to distinguish small structures in many specimens because microorganisms are relatively transparent. It is often necessary to increase **contrast** to detect different



structures in a specimen. Various types of microscopes use different features of light or electrons to increase contrast—visible differences between the parts of a specimen (see [Section 2.3](#)). Additionally, dyes that bind to some structures but not others can be used to improve the contrast between images of relatively transparent objects (see [Section 2.4](#)).



### Check Your Understanding

- Explain the difference between magnification and resolution.
- Explain the difference between resolution and contrast.
- Name two factors that affect resolution.

## 2.2 Peering Into the Invisible World

---


### Learning Objectives

- Describe historical developments and individual contributions that led to the invention and development of the microscope
- Compare and contrast the features of simple and compound microscopes

Some of the fundamental characteristics and functions of microscopes can be understood in the context of the history of their use. Italian scholar Girolamo Fracastoro is regarded as the first person to formally postulate that disease was spread by tiny invisible *seminaria*, or “seeds of the contagion.” In his book *De Contagione* (1546), he proposed that these seeds could attach themselves to certain objects (which he called *fomes* [cloth]) that supported their transfer from person to person. However, since the technology for seeing such tiny objects did not yet exist, the existence of the *seminaria* remained hypothetical for a little over a century—an invisible world waiting to be revealed.

### Early Microscopes


Antonie van Leeuwenhoek, sometimes hailed as “the Father of Microbiology,” is typically credited as the first person to have created microscopes powerful enough to view microbes ([Figure 2.9](#)). Born in the city of Delft in the Dutch Republic, van Leeuwenhoek began his career selling fabrics. However, he later became interested in lens making (perhaps to look at threads) and his innovative techniques produced microscopes that allowed him to observe microorganisms as no one had before. In 1674, he described his observations of single-celled organisms, whose existence was previously unknown, in a series of letters to the Royal Society of London. His report was initially met with skepticism, but his claims were soon verified and he became something of a celebrity in the scientific community.



**Figure 2.9** (a) Antonie van Leeuwenhoek (1632–1723) is credited as being the first person to observe microbes, including bacteria, which he called “animalcules” and “wee little beasties.” (b) Even though van Leeuwenhoek’s microscopes were simple microscopes (as seen in this replica), they were more powerful and provided better resolution than the compound microscopes of his day. (credit b: modification of work by “Wellcome Images”/Wikimedia Commons) (c) Though more famous for developing the telescope, Galileo Galilei (1564–1642) was also one of the pioneers of microscopy.

While van Leeuwenhoek is credited with the discovery of microorganisms, others before him had contributed to the development of the microscope. These included eyeglass makers in the Netherlands in the late 1500s, as well as the Italian astronomer Galileo Galilei, who used a **compound microscope** to examine insect parts (??). Whereas van Leeuwenhoek used a **simple microscope**, in which light is passed through just one lens, Galileo’s compound microscope was more sophisticated, passing light through two sets of lenses.

Van Leeuwenhoek’s contemporary, the Englishman Robert Hooke (1635–1703), also made important contributions to microscopy, publishing in his book *Micrographia* (1665) many observations using compound microscopes. Viewing a thin sample of cork through his microscope, he was the first to observe the structures that we now know as cells (**Figure 2.10**). Hooke described these structures as resembling “Honey-comb,” and as “small Boxes or Bladders of Air,” noting that each “Cavern, Bubble, or Cell” is distinct from the others (in Latin, “cell” literally means “small room”). They likely appeared to Hooke to be filled with air because the cork cells were dead, with only the rigid cell walls providing the structure.



**Figure 2.10** Robert Hooke used his (a) compound microscope to view (b) cork cells. Both of these engravings are from his seminal work *Micrographia*, published in 1665.



## Check Your Understanding

- Explain the difference between simple and compound microscopes.
- Compare and contrast the contributions of van Leeuwenhoek, Hooke, and Galileo to early microscopy.

### Micro Connections

#### Who Invented the Microscope?

While Antonie van Leeuwenhoek and Robert Hooke generally receive much of the credit for early advances in microscopy, neither can claim to be the inventor of the microscope. Some argue that this designation should belong to Hans and Zaccharias Janssen, Dutch spectacle-makers who may have invented the telescope, the simple microscope, and the compound microscope during the late 1500s or early 1600s (**Figure 2.11**). Unfortunately, little is known for sure about the Janssens, not even the exact dates of their births and deaths. The Janssens were secretive about their work and never published. It is also possible that the Janssens did not invent anything at all; their neighbor, Hans Lippershey, also developed microscopes and telescopes during the same time frame, and he is often credited with inventing the telescope. The historical records from the time are as fuzzy and imprecise as the images viewed through those early lenses, and any archived records have been lost over the centuries.

By contrast, van Leeuwenhoek and Hooke can thank ample documentation of their work for their respective legacies. Like Janssen, van Leeuwenhoek began his work in obscurity, leaving behind few records. However, his friend, the prominent physician Reinier de Graaf, wrote a letter to the editor of the *Philosophical Transactions of the Royal Society of London* calling attention to van Leeuwenhoek's powerful microscopes. From 1673 onward, van Leeuwenhoek began regularly submitting letters to the Royal Society detailing his observations. In 1674, his report describing single-celled organisms produced controversy in the scientific community, but his observations were soon confirmed when the society sent a delegation to investigate his findings. He subsequently enjoyed considerable celebrity, at one point even entertaining a visit by the czar of Russia.

Similarly, Robert Hooke had his observations using microscopes published by the Royal Society in a book called *Micrographia* in 1665. The book became a bestseller and greatly increased interest in microscopy throughout much of Europe.



**Figure 2.11** Zaccharias Janssen, along with his father Hans, may have invented the telescope, the simple microscope, and the compound microscope during the late 1500s or early 1600s. The historical evidence is inconclusive.

## 2.3 Instruments of Microscopy

### Learning Objectives

- Identify and describe the parts of a brightfield microscope
- Calculate total magnification for a compound microscope
- Describe the distinguishing features and typical uses for various types of light microscopes, electron microscopes, and scanning probe microscopes

The early pioneers of microscopy opened a window into the invisible world of microorganisms. But microscopy continued to advance in the centuries that followed. In 1830, Joseph Jackson Lister created an essentially modern light microscope. The 20th century saw the development of microscopes that leveraged nonvisible light, such as fluorescence microscopy, which uses an ultraviolet light source, and electron microscopy, which uses short-wavelength electron beams. These advances led to major improvements in magnification, resolution, and contrast. By comparison, the relatively rudimentary microscopes of van Leeuwenhoek and his contemporaries were far less powerful than even the most basic microscopes in use today. In this section, we will survey the broad range of modern microscopic technology and common applications for each type of microscope.

### Light Microscopy

Many types of microscopes fall under the category of light microscopes, which use light to visualize images. Examples of light microscopes include brightfield microscopes, darkfield microscopes, phase-contrast microscopes, differential interference contrast microscopes, fluorescence microscopes, confocal scanning laser microscopes, and two-photon microscopes. These various types of light microscopes can be used to complement each other in diagnostics and research.


## Brightfield Microscopes

The **brightfield microscope**, perhaps the most commonly used type of microscope, is a compound microscope with two or more lenses that produce a dark image on a bright background. Some brightfield microscopes are **monocular** (having a single eyepiece), though most newer brightfield microscopes are **binocular** (having two eyepieces), like the one shown in **Figure 2.12**; in either case, each eyepiece contains a lens called an **ocular lens**. The ocular lenses typically magnify images 10 times ( $10\times$ ). At the other end of the body tube are a set of **objective lenses** on a rotating nosepiece. The magnification of these objective lenses typically ranges from  $4\times$  to  $100\times$ , with the magnification for each lens designated on the metal casing of the lens. The ocular and objective lenses work together to create a magnified image. The **total magnification** is the product of the ocular magnification times the objective magnification:

$$\text{ocular magnification} \times \text{objective magnification}$$

For example, if a  $40\times$  objective lens is selected and the ocular lens is  $10\times$ , the total magnification would be

$$(40\times)(10\times) = 400\times$$



**Figure 2.12** Components of a typical brightfield microscope.

The item being viewed is called a specimen. The specimen is placed on a glass slide, which is then clipped into place on the **stage** (a platform) of the microscope. Once the slide is secured, the specimen on the slide is positioned over the light using the **x-y mechanical stage knobs**. These knobs move the slide on the surface of the stage, but do not raise or lower the stage. Once the specimen is centered over the light, the stage position can be raised or lowered to focus the image. The **coarse focusing knob** is used for large-scale movements with  $4\times$  and  $10\times$  objective lenses; the **fine focusing knob** is used for small-scale movements, especially with  $40\times$  or  $100\times$  objective lenses.


When images are magnified, they become dimmer because there is less light per unit area of image. Highly magnified images produced by microscopes, therefore, require intense lighting. In a brightfield microscope, this light is provided by an **illuminator**, which is typically a high-intensity bulb below the stage. Light from the illuminator passes up through **condenser lens** (located below the stage), which focuses all of the light rays on the specimen to maximize illumination. The position of the condenser can be optimized using the attached condenser focus knob; once the optimal distance is established, the condenser should not be moved to adjust the brightness. If less-than-maximal light



levels are needed, the amount of light striking the specimen can be easily adjusted by opening or closing a **diaphragm** between the condenser and the specimen. In some cases, brightness can also be adjusted using the **rheostat**, a dimmer switch that controls the intensity of the illuminator.

A brightfield microscope creates an image by directing light from the illuminator at the specimen; this light is differentially transmitted, absorbed, reflected, or refracted by different structures. Different colors can behave differently as they interact with **chromophores** (pigments that absorb and reflect particular wavelengths of light) in parts of the specimen. Often, chromophores are artificially added to the specimen using stains, which serve to increase contrast and resolution. In general, structures in the specimen will appear darker, to various extents, than the bright background, creating maximally sharp images at magnifications up to about  $1000\times$ . Further magnification would create a larger image, but without increased resolution. This allows us to see objects as small as bacteria, which are visible at about  $400\times$  or so, but not smaller objects such as viruses.

At very high magnifications, resolution may be compromised when light passes through the small amount of air between the specimen and the lens. This is due to the large difference between the refractive indices of air and glass; the air scatters the light rays before they can be focused by the lens. To solve this problem, a drop of oil can be used to fill the space between the specimen and an **oil immersion lens**, a special lens designed to be used with immersion oils. Since the oil has a refractive index very similar to that of glass, it increases the maximum angle at which light leaving the specimen can strike the lens. This increases the light collected and, thus, the resolution of the image (**Figure 2.13**). A variety of oils can be used for different types of light.



**Figure 2.13** (a) Oil immersion lenses like this one are used to improve resolution. (b) Because immersion oil and glass have very similar refractive indices, there is a minimal amount of refraction before the light reaches the lens. Without immersion oil, light scatters as it passes through the air above the slide, degrading the resolution of the image.

## Micro Connections

### Microscope Maintenance: Best Practices

Even a very powerful microscope cannot deliver high-resolution images if it is not properly cleaned and maintained. Since lenses are carefully designed and manufactured to refract light with a high degree of precision, even a slightly dirty or scratched lens will refract light in unintended ways, degrading the image of the specimen. In addition, microscopes are rather delicate instruments, and great care must be taken to avoid damaging parts and surfaces. Among other things, proper care of a microscope includes the following:

- cleaning the lenses with lens paper
- not allowing lenses to contact the slide (e.g., by rapidly changing the focus)
- protecting the bulb (if there is one) from breakage
- not pushing an objective into a slide
- not using the coarse focusing knob when using the 40× or greater objective lenses
- only using immersion oil with a specialized oil objective, usually the 100× objective
- cleaning oil from immersion lenses after using the microscope
- cleaning any oil accidentally transferred from other lenses
- covering the microscope or placing it in a cabinet when not in use

### Link to Learning




Visit the online resources linked below for simulations and demonstrations involving the use of microscopes. Keep in mind that execution of specific techniques and procedures can vary depending on the specific instrument you are using. Thus, it is important to learn and practice with an actual microscope in a laboratory setting under expert supervision.

- University of Delaware's **Virtual Microscope**  
(<http://www.openstaxcollege.org//22virtualsim>)
- St. John's University **Microscope Tutorials**  
(<http://www.openstaxcollege.org//22microtut>)

### Darkfield Microscopy

A **darkfield microscope** is a brightfield microscope that has a small but significant modification to the condenser. A small, opaque disk (about 1 cm in diameter) is placed between the illuminator and the condenser lens. This opaque light stop, as the disk is called, blocks most of the light from the illuminator as it passes through the condenser on its way to the objective lens, producing a hollow cone of light that is focused on the specimen. The only light that reaches the objective is light that has been refracted or reflected by structures in the specimen. The resulting image typically shows bright objects on a dark background (**Figure 2.14**).



**Figure 2.14** An opaque light stop inserted into a brightfield microscope is used to produce a darkfield image. The light stop blocks light traveling directly from the illuminator to the objective lens, allowing only light reflected or refracted off the specimen to reach the eye.

Darkfield microscopy can often create high-contrast, high-resolution images of specimens without the use of stains, which is particularly useful for viewing live specimens that might be killed or otherwise compromised by the stains. For example, thin spirochetes like *Treponema pallidum*, the causative agent of syphilis, can be best viewed using a darkfield microscope (**Figure 2.15**).



**Figure 2.15** Use of a darkfield microscope allows us to view living, unstained samples of the spirochete *Treponema pallidum*. Similar to a photographic negative, the spirochetes appear bright against a dark background. (credit: Centers for Disease Control and Prevention/C.W. Hubbard)



### Check Your Understanding

- Identify the key differences between brightfield and darkfield microscopy.

## Clinical Focus

### Part 2

Wound infections like Cindy's can be caused by many different types of bacteria, some of which can spread rapidly with serious complications. Identifying the specific cause is very important to select a medication that can kill or stop the growth of the bacteria.

After calling a local doctor about Cindy's case, the camp nurse sends the sample from the wound to the closest medical laboratory. Unfortunately, since the camp is in a remote area, the nearest lab is small and poorly equipped. A more modern lab would likely use other methods to culture, grow, and identify the bacteria, but in this case, the technician decides to make a wet mount from the specimen and view it under a brightfield microscope. In a wet mount, a small drop of water is added to the slide, and a cover slip is placed over the specimen to keep it in place before it is positioned under the objective lens.


Under the brightfield microscope, the technician can barely see the bacteria cells because they are nearly transparent against the bright background. To increase contrast, the technician inserts an opaque light stop above the illuminator. The resulting darkfield image clearly shows that the bacteria cells are spherical and grouped in clusters, like grapes.

- Why is it important to identify the shape and growth patterns of cells in a specimen?
- What other types of microscopy could be used effectively to view this specimen?

Jump to the **next** Clinical Focus box. Go back to the **previous** Clinical Focus box.


## Phase-Contrast Microscopes

**Phase-contrast microscopes** use refraction and interference caused by structures in a specimen to create high-contrast, high-resolution images without staining. It is the oldest and simplest type of microscope that creates an image by altering the wavelengths of light rays passing through the specimen. To create altered wavelength paths, an annular stop is used in the condenser. The annular stop produces a hollow cone of light that is focused on the specimen before reaching the objective lens. The objective contains a phase plate containing a phase ring. As a result, light traveling directly from the illuminator passes through the phase ring while light refracted or reflected by the specimen passes through the plate. This causes waves traveling through the ring to be about one-half of a wavelength out of phase with those passing through the plate. Because waves have peaks and troughs, they can add together (if in phase together) or cancel each other out (if out of phase). When the wavelengths are out of phase, wave troughs will cancel out wave peaks, which is called destructive interference. Structures that refract light then appear dark against a bright background of only unrefracted light. More generally, structures that differ in features such as refractive index will differ in levels of darkness (**Figure 2.16**).



**Figure 2.16** This diagram of a phase-contrast microscope illustrates phase differences between light passing through the object and background. These differences are produced by passing the rays through different parts of a phase plate. The light rays are superimposed in the image plane, producing contrast due to their interference.

Because it increases contrast without requiring stains, phase-contrast microscopy is often used to observe live specimens. Certain structures, such as organelles in eukaryotic cells and endospores in prokaryotic cells, are especially well visualized with phase-contrast microscopy (**Figure 2.17**).




**Figure 2.17** This figure compares a brightfield image (left) with a phase-contrast image (right) of the same unstained simple squamous epithelial cells. The cells are in the center and bottom right of each photograph (the irregular item above the cells is acellular debris). Notice that the unstained cells in the brightfield image are almost invisible against the background, whereas the cells in the phase-contrast image appear to glow against the background, revealing far more detail. (credit: "Clearly kefir"/Wikimedia Commons)



## Differential Interference Contrast Microscopes

**Differential interference contrast (DIC) microscopes** (also known as Nomarski optics) are similar to phase-contrast microscopes in that they use interference patterns to enhance contrast between different features of a specimen. In a DIC microscope, two beams of light are created in which the direction of wave movement (polarization) differs. Once the beams pass through either the specimen or specimen-free space, they are recombined and effects of the specimens cause differences in the interference patterns generated by the combining of the beams. This results in high-contrast images of living organisms with a three-dimensional appearance. These microscopes are especially useful in distinguishing structures within live, unstained specimens. (Figure 2.18)



**Figure 2.18** A DIC image of *Fonsecaea pedrosoi* grown on modified Leonian's agar. This fungus causes chromoblastomycosis, a chronic skin infection common in tropical and subtropical climates.



### Check Your Understanding

- What are some advantages of phase-contrast and DIC microscopy?

## Fluorescence Microscopes

A **fluorescence microscope** uses fluorescent chromophores called **fluorochromes**, which are capable of absorbing energy from a light source and then emitting this energy as visible light. Fluorochromes include naturally fluorescent substances (such as chlorophylls) as well as fluorescent stains that are added to the specimen to create contrast. Dyes such as Texas red and FITC are examples of fluorochromes. Other examples include the nucleic acid dyes 4',6'-diamidino-2-phenylindole (DAPI) and acridine orange.


The microscope transmits an excitation light, generally a form of EMR with a short wavelength, such as ultraviolet or blue light, toward the specimen; the chromophores absorb the excitation light and emit visible light with longer wavelengths. The excitation light is then filtered out (in part because ultraviolet light is harmful to the eyes) so that only visible light passes through the ocular lens. This produces an image of the specimen in bright colors against a dark background.

Fluorescence microscopes are especially useful in clinical microbiology. They can be used to identify pathogens, to find particular species within an environment, or to find the locations of particular molecules and structures within a cell. Approaches have also been developed to distinguish living from dead cells using fluorescence microscopy based upon whether they take up particular fluorochromes. Sometimes, multiple fluorochromes are used on the same specimen to show different structures or features.

One of the most important applications of fluorescence microscopy is a technique called **immunofluorescence**, which is used to identify certain disease-causing microbes by observing whether antibodies bind to them. (Antibodies are protein molecules produced by the immune system that attach to specific pathogens to kill or inhibit them.) There are

two approaches to this technique: direct immunofluorescence assay (DFA) and indirect immunofluorescence assay (IFA). In DFA, specific antibodies (e.g., those that target the rabies virus) are stained with a fluorochrome. If the specimen contains the targeted pathogen, one can observe the antibodies binding to the pathogen under the fluorescent microscope. This is called a primary antibody stain because the stained antibodies attach directly to the pathogen.

In IFA, secondary antibodies are stained with a fluorochrome rather than primary antibodies. Secondary antibodies do not attach directly to the pathogen, but they do bind to primary antibodies. When the unstained primary antibodies bind to the pathogen, the fluorescent secondary antibodies can be observed binding to the primary antibodies. Thus, the secondary antibodies are attached indirectly to the pathogen. Since multiple secondary antibodies can often attach to a primary antibody, IFA increases the number of fluorescent antibodies attached to the specimen, making it easier to visualize features in the specimen (Figure 2.19).



**Figure 2.19** (a) A direct immunofluorescent stain is used to visualize *Neisseria gonorrhoeae*, the bacterium that causes gonorrhea. (b) An indirect immunofluorescent stain is used to visualize larvae of *Schistosoma mansoni*, a parasitic worm that causes schistosomiasis, an intestinal disease common in the tropics. (c) In direct immunofluorescence, the stain is absorbed by a primary antibody, which binds to the antigen. In indirect immunofluorescence, the stain is absorbed by a secondary antibody, which binds to a primary antibody, which, in turn, binds to the antigen. (credit a: modification of work by Centers for Disease Control and Prevention; credit b: modification of work by Centers for Disease Control and Prevention/Dr. Sulzer)




### Check Your Understanding

- Why must fluorochromes be used to examine a specimen under a fluorescence microscope?

### Confocal Microscopes

Whereas other forms of light microscopy create an image that is maximally focused at a single distance from the observer (the depth, or z-plane), a **confocal microscope** uses a laser to scan multiple z-planes successively. This produces numerous two-dimensional, high-resolution images at various depths, which can be constructed into a three-dimensional image by a computer. As with fluorescence microscopes, fluorescent stains are generally used to increase

contrast and resolution. Image clarity is further enhanced by a narrow aperture that eliminates any light that is not from the z-plane. Confocal microscopes are thus very useful for examining thick specimens such as biofilms, which can be examined alive and unfixed (**Figure 2.20**).



**Figure 2.20** Confocal microscopy can be used to visualize structures such as this roof-dwelling cyanobacterium biofilm. (credit: American Society for Microbiology)

### Link to Learning



Explore a rotating three-dimensional **view** (<http://www.openstaxcollege.org//22biofilm3d>) of a biofilm as observed under a confocal microscope. After navigating to the webpage, click the “play” button to launch the video.

## Two-Photon Microscopes

While the original fluorescent and confocal microscopes allowed better visualization of unique features in specimens, there were still problems that prevented optimum visualization. The effective sensitivity of fluorescence microscopy when viewing thick specimens was generally limited by out-of-focus flare, which resulted in poor resolution. This limitation was greatly reduced in the confocal microscope through the use of a confocal pinhole to reject out-of-focus background fluorescence with thin ( $<1 \mu\text{m}$ ), unblurred optical sections. However, even the confocal microscopes lacked the resolution needed for viewing thick tissue samples. These problems were resolved with the development of the **two-photon microscope**, which uses a scanning technique, fluorochromes, and long-wavelength light (such as infrared) to visualize specimens. The low energy associated with the long-wavelength light means that two photons must strike a location at the same time to excite the fluorochrome. The low energy of the excitation light is less damaging to cells, and the long wavelength of the excitation light more easily penetrates deep into thick specimens. This makes the two-photon microscope useful for examining living cells within intact tissues—brain slices, embryos, whole organs, and even entire animals.

Currently, use of two-photon microscopes is limited to advanced clinical and research laboratories because of the high costs of the instruments. A single two-photon microscope typically costs between \$300,000 and \$500,000, and the lasers used to excite the dyes used on specimens are also very expensive. However, as technology improves, two-photon microscopes may become more readily available in clinical settings.



## Check Your Understanding

- What types of specimens are best examined using confocal or two-photon microscopy?

## Electron Microscopy

The maximum theoretical resolution of images created by light microscopes is ultimately limited by the wavelengths of visible light. Most light microscopes can only magnify  $1000\times$ , and a few can magnify up to  $1500\times$ , but this does not begin to approach the magnifying power of an **electron microscope (EM)**, which uses short-wavelength electron beams rather than light to increase magnification and resolution.

Electrons, like electromagnetic radiation, can behave as waves, but with wavelengths of  $0.005\text{ nm}$ , they can produce much better resolution than visible light. An EM can produce a sharp image that is magnified up to  $100,000\times$ . Thus, EMs can resolve subcellular structures as well as some molecular structures (e.g., single strands of DNA); however, electron microscopy cannot be used on living material because of the methods needed to prepare the specimens.

There are two basic types of EM: the **transmission electron microscope (TEM)** and the **scanning electron microscope (SEM)** (Figure 2.21). The TEM is somewhat analogous to the brightfield light microscope in terms of the way it functions. However, it uses an electron beam from above the specimen that is focused using a magnetic lens (rather than a glass lens) and projected through the specimen onto a detector. Electrons pass through the specimen, and then the detector captures the image (Figure 2.22).




(a)



(b)

**Figure 2.21** (a) A transmission electron microscope (TEM). (b) A scanning electron microscope (SEM). (credit a: modification of work by "Deshi"/Wikimedia Commons; credit b: modification of work by "ZEISS Microscopy"/Flickr)




**Figure 2.22** Electron microscopes use magnets to focus electron beams similarly to the way that light microscopes use lenses to focus light.

For electrons to pass through the specimen in a TEM, the specimen must be extremely thin (20–100 nm thick). The image is produced because of varying opacity in various parts of the specimen. This opacity can be enhanced by staining the specimen with materials such as heavy metals, which are electron dense. TEM requires that the beam and specimen be in a vacuum and that the specimen be very thin and dehydrated. The specific steps needed to prepare a specimen for observation under an EM are discussed in detail in the next section.


SEMs form images of surfaces of specimens, usually from electrons that are knocked off of specimens by a beam of electrons. This can create highly detailed images with a three-dimensional appearance that are displayed on a monitor (**Figure 2.23**). Typically, specimens are dried and prepared with fixatives that reduce artifacts, such as shriveling, that can be produced by drying, before being sputter-coated with a thin layer of metal such as gold. Whereas transmission electron microscopy requires very thin sections and allows one to see internal structures such as organelles and the interior of membranes, scanning electron microscopy can be used to view the surfaces of larger objects (such as a pollen grain) as well as the surfaces of very small samples (**Figure 2.24**). Some EMs can magnify an image up to  $2,000,000\times$ .<sup>[1]</sup>

1. “JEM-ARM200F Transmission Electron Microscope,” *JEOL USA Inc*, <http://www.jeolusa.com/PRODUCTS/TransmissionElectronMicroscopes%28TEM%29/200kV/JEM-ARM200F/tabid/663/Default.aspx#195028-specifications>. Accessed 8/28/2015.





**Figure 2.23** These schematic illustrations compare the components of transmission electron microscopes and scanning electron microscopes.



**Figure 2.24** (a) This TEM image of cells in a biofilm shows well-defined internal structures of the cells because of varying levels of opacity in the specimen. (b) This color-enhanced SEM image of the bacterium *Staphylococcus aureus* illustrates the ability of scanning electron microscopy to render three-dimensional images of the surface structure of cells. (credit a: modification of work by American Society for Microbiology; credit b: modification of work by Centers for Disease Control and Prevention)



### Check Your Understanding

- What are some advantages and disadvantages of electron microscopy, as opposed to light microscopy, for examining microbiological specimens?
- What kinds of specimens are best examined using TEM? SEM?

## Micro Connections

### Using Microscopy to Study Biofilms

A biofilm is a complex community of one or more microorganism species, typically forming as a slimy coating attached to a surface because of the production of an extrapolymeric substance (EPS) that attaches to a surface or at the interface between surfaces (e.g., between air and water). In nature, biofilms are abundant and frequently occupy complex niches within ecosystems (Figure 2.25). In medicine, biofilms can coat medical devices and exist within the body. Because they possess unique characteristics, such as increased resistance against the immune system and to antimicrobial drugs, biofilms are of particular interest to microbiologists and clinicians alike.

Because biofilms are thick, they cannot be observed very well using light microscopy; slicing a biofilm to create a thinner specimen might kill or disturb the microbial community. Confocal microscopy provides clearer images of biofilms because it can focus on one z-plane at a time and produce a three-dimensional image of a thick specimen. Fluorescent dyes can be helpful in identifying cells within the matrix. Additionally, techniques such as immunofluorescence and fluorescence in situ hybridization (FISH), in which fluorescent probes are used to bind to DNA, can be used.

Electron microscopy can be used to observe biofilms, but only after dehydrating the specimen, which produces undesirable artifacts and distorts the specimen. In addition to these approaches, it is possible to follow water currents through the shapes (such as cones and mushrooms) of biofilms, using video of the movement of fluorescently coated beads (Figure 2.26).





Diagram showing five stages of biofilm development of *Pseudomonas aeruginosa*. All photomicrographs are shown to same scale.

**Figure 2.25** A biofilm forms when planktonic (free-floating) bacteria of one or more species adhere to a surface, produce slime, and form a colony. (credit: Public Library of Science)




**Figure 2.26** In this image, multiple species of bacteria grow in a biofilm on stainless steel (stained with DAPI for epifluorescence microscopy). (credit: Ricardo Murga, Rodney Donlan)

## Scanning Probe Microscopy

A **scanning probe microscope** does not use light or electrons, but rather very sharp probes that are passed over the surface of the specimen and interact with it directly. This produces information that can be assembled into images with magnifications up to  $100,000,000\times$ . Such large magnifications can be used to observe individual atoms on surfaces. To date, these techniques have been used primarily for research rather than for diagnostics.

There are two types of scanning probe microscope: the **scanning tunneling microscope (STM)** and the **atomic force microscope (AFM)**. An STM uses a probe that is passed just above the specimen as a constant voltage bias creates the potential for an electric current between the probe and the specimen. This current occurs via quantum tunneling of electrons between the probe and the specimen, and the intensity of the current is dependent upon the distance between the probe and the specimen. The probe is moved horizontally above the surface and the intensity of the current is measured. Scanning tunneling microscopy can effectively map the structure of surfaces at a resolution at which individual atoms can be detected.

Similar to an STM, AFMs have a thin probe that is passed just above the specimen. However, rather than measuring variations in the current at a constant height above the specimen, an AFM establishes a constant current and measures variations in the height of the probe tip as it passes over the specimen. As the probe tip is passed over the specimen, forces between the atoms (van der Waals forces, capillary forces, chemical bonding, electrostatic forces, and others) cause it to move up and down. Deflection of the probe tip is determined and measured using Hooke's law of elasticity, and this information is used to construct images of the surface of the specimen with resolution at the atomic level (**Figure 2.27**).



**Figure 2.27** STMs and AFMs allow us to view images at the atomic level. (a) This STM image of a pure gold surface shows individual atoms of gold arranged in columns. (b) This AFM image shows long, strand-like molecules of nanocellulose, a laboratory-created substance derived from plant fibers. (credit a: modification of work by "Erwinrossen"/Wikimedia Commons)










### Check Your Understanding

- Which has higher magnification, a light microscope or a scanning probe microscope?
- Name one advantage and one limitation of scanning probe microscopy.



**Figure 2.28**, **Figure 2.29**, and **Figure 2.30** summarize the microscopy techniques for light microscopes, electron microscopes, and scanning probe microscopes, respectively.





<b>LIGHT MICROSCOPES</b> <b>Magnification: up to about 1000×</b> Use visible or ultraviolet light to produce an image.		
Microscope Type	Key Uses	Sample Images
<b>Brightfield</b>	Commonly used in a wide variety of laboratory applications as the standard microscope; produces an image on a bright background.  <b>Example:</b> <i>Bacillus</i> sp. showing endospores.	
<b>Darkfield</b>	Increases contrast without staining by producing a bright image on a darker background; especially useful for viewing live specimens.  <b>Example:</b> <i>Borrelia burgdorferi</i>	
<b>Phase contrast</b>	Uses refraction and interference caused by structures in the specimen to create high-contrast, high-resolution images without staining, making it useful for viewing live specimens, and structures such as endospores and organelles.  <b>Example:</b> <i>Pseudomonas</i> sp.	
<b>Differential interference contrast (DIC)</b>	Uses interference patterns to enhance contrast between different features of a specimen to produce high-contrast images of living organisms with a three-dimensional appearance, making it especially useful in distinguishing structures within live, unstained specimens; images viewed reveal detailed structures within cells.  <b>Example:</b> <i>Escherichia coli</i> O157:H7	
<b>Fluorescence</b>	Uses fluorescent stains to produce an image; can be used to identify pathogens, to find particular species, to distinguish living from dead cells, or to find locations of particular molecules within a cell; also used for immunofluorescence.  <b>Example:</b> <i>P. putida</i> stained with fluorescent dyes to visualize the capsule.	
<b>Confocal</b>	Uses a laser to scan multiple z-planes successively, producing numerous two-dimensional, high-resolution images at various depths that can be constructed into a three-dimensional image by a computer, making this useful for examining thick specimens such as biofilms.  <b>Example:</b> <i>Escherichia coli</i> stained with acridine orange dye to show the nucleoid regions of the cells.	
<b>Two-photon</b>	Uses a scanning technique, fluorochromes, and long-wavelength light (such as infrared) to penetrate deep into thick specimens such as biofilms.  <b>Example:</b> Mouse intestine cells stained with fluorescent dye.	

**Figure 2.28** (credit “Brightfield”: modification of work by American Society for Microbiology; credit “Darkfield”: modification of work by American Society for Microbiology; credit “Phase contrast”: modification of work by American Society for Microbiology; credit “DIC”: modification of work by American Society for Microbiology; credit “Fluorescence”: modification of work by American Society for Microbiology; credit “Confocal”: modification of work by American Society for Microbiology; credit “Two-photon”: modification of work by Alberto Diaspro, Paolo Bianchini, Giuseppe Vicidomini, Mario Faretta, Paola Ramoino, Cesare Usai)



ELECTRON MICROSCOPES      Magnification: 20–100,000× or more		
Use electron beams focused with magnets to produce an image.		
Microscope Type	Key Uses	Sample Images
<b>Transmission (TEM)</b>	<p>Uses electron beams that pass through a specimen to visualize small images; useful to observe small, thin specimens such as tissue sections and subcellular structures.</p> <p><b>Example:</b> <i>Ebola virus</i></p>	
<b>Scanning (SEM)</b>	<p>Uses electron beams to visualize surfaces; useful to observe the three-dimensional surface details of specimens.</p> <p><b>Example:</b> <i>Campylobacter jejuni</i></p>	

**Figure 2.29** (credit “TEM”: modification of work by American Society for Microbiology; credit “SEM”: modification of work by American Society for Microbiology)

SCANNING PROBE MICROSCOPES      Magnification: 100–100,000,000× or more		
Use very short probes that are passed over the surface of the specimen and interact with it directly.		
Microscope Type	Key Uses	Sample Images
<b>Scanning tunneling (STM)</b>	<p>Uses a probe passed horizontally at a constant distance just above the specimen while the intensity of the current is measured; can map the structure of surfaces at the atomic level; works best on conducting materials but can also be used to examine organic materials such as DNA, if fixed on a surface.</p> <p><b>Example:</b> Image of surface reconstruction on a clean gold [Au(100)] surface, as visualized using scanning tunneling microscopy.</p>	
<b>Atomic force (AFM)</b>	<p>Can be used in several ways, including using a laser focused on a cantilever to measure the bending of the tip or a probe passed above the specimen while the height needed to maintain a constant current is measured; useful to observe specimens at the atomic level and can be more easily used with nonconducting samples.</p> <p><b>Example:</b> AFM height image of carboxymethylated nanocellulose adsorbed on a silica surface.</p>	

**Figure 2.30** (credit “STM”: modification of work by “Erwinrossen”/Wikimedia Commons)

## 2.4 Staining Microscopic Specimens

---

### Learning Objectives

- Differentiate between simple and differential stains
- Describe the unique features of commonly used stains
- Explain the procedures and name clinical applications for Gram, endospore, acid-fast, negative capsule, and flagella staining

In their natural state, most of the cells and microorganisms that we observe under the microscope lack color and contrast. This makes it difficult, if not impossible, to detect important cellular structures and their distinguishing characteristics without artificially treating specimens. We have already alluded to certain techniques involving stains and fluorescent dyes, and in this section we will discuss specific techniques for sample preparation in greater detail. Indeed, numerous methods have been developed to identify specific microbes, cellular structures, DNA sequences, or indicators of infection in tissue samples, under the microscope. Here, we will focus on the most clinically relevant techniques.


### Preparing Specimens for Light Microscopy

In clinical settings, light microscopes are the most commonly used microscopes. There are two basic types of preparation used to view specimens with a light microscope: wet mounts and fixed specimens.

The simplest type of preparation is the **wet mount**, in which the specimen is placed on the slide in a drop of liquid. Some specimens, such as a drop of urine, are already in a liquid form and can be deposited on the slide using a dropper. Solid specimens, such as a skin scraping, can be placed on the slide before adding a drop of liquid to prepare the wet mount. Sometimes the liquid used is simply water, but often stains are added to enhance contrast. Once the liquid has been added to the slide, a coverslip is placed on top and the specimen is ready for examination under the microscope.

The second method of preparing specimens for light microscopy is **fixation**. The “fixing” of a sample refers to the process of attaching cells to a slide. Fixation is often achieved either by heating (heat fixing) or chemically treating the specimen. In addition to attaching the specimen to the slide, fixation also kills microorganisms in the specimen, stopping their movement and metabolism while preserving the integrity of their cellular components for observation.


To heat-fix a sample, a thin layer of the specimen is spread on the slide (called a **smear**), and the slide is then briefly heated over a heat source (**Figure 2.31**). Chemical fixatives are often preferable to heat for tissue specimens. Chemical agents such as acetic acid, ethanol, methanol, formaldehyde (formalin), and glutaraldehyde can denature proteins, stop biochemical reactions, and stabilize cell structures in tissue samples (**Figure 2.31**).



**Figure 2.31** (a) A specimen can be heat-fixed by using a slide warmer like this one. (b) Another method for heat-fixing a specimen is to hold a slide with a smear over a microincinerator. (c) This tissue sample is being fixed in a solution of formalin (also known as formaldehyde). Chemical fixation kills microorganisms in the specimen, stopping degradation of the tissues and preserving their structure so that they can be examined later under the microscope. (credit a: modification of work by Nina Parker; credit b: modification of work by Nina Parker; credit c: modification of work by “University of Bristol”/YouTube)

In addition to fixation, **staining** is almost always applied to color certain features of a specimen before examining it under a light microscope. Stains, or dyes, contain salts made up of a positive ion and a negative ion. Depending on the type of dye, the positive or the negative ion may be the chromophore (the colored ion); the other, uncolored ion is called the counterion. If the chromophore is the positively charged ion, the stain is classified as a **basic dye**; if the negative ion is the chromophore, the stain is considered an **acidic dye**.

Dyes are selected for staining based on the chemical properties of the dye and the specimen being observed, which determine how the dye will interact with the specimen. In most cases, it is preferable to use a **positive stain**, a dye that will be absorbed by the cells or organisms being observed, adding color to objects of interest to make them stand out against the background. However, there are scenarios in which it is advantageous to use a **negative stain**, which is absorbed by the background but not by the cells or organisms in the specimen. Negative staining produces an outline or silhouette of the organisms against a colorful background (**Figure 2.32**).



**Figure 2.32** (a) These *Bacillus anthracis* cells have absorbed crystal violet, a basic positive stain. (b) This specimen of *Spinoloricus*, a microscopic marine organism, has been stained with rose bengal, a positive acidic stain. (c) These *B. megaterium* appear to be white because they have not absorbed the negative red stain applied to the slide. (credit a: modification of work by Centers for Disease Control and Prevention; credit b: modification of work by Roberto Danovaro, Antonio Pusceddu, Cristina Gambi, Iben Heiner, Reinhardt Mobjerg Kristensen; credit c: modification of work by Anh-Hue Tu)

Because cells typically have negatively charged cell walls, the positive chromophores in basic dyes tend to stick to the cell walls, making them positive stains. Thus, commonly used basic dyes such as basic fuchsin, crystal violet,

malachite green, methylene blue, and safranin typically serve as positive stains. On the other hand, the negatively charged chromophores in acidic dyes are repelled by negatively charged cell walls, making them negative stains. Commonly used acidic dyes include acid fuchsin, eosin, and rose bengal. **Figure 2.40** provides more detail.

Some staining techniques involve the application of only one dye to the sample; others require more than one dye. In **simple staining**, a single dye is used to emphasize particular structures in the specimen. A simple stain will generally make all of the organisms in a sample appear to be the same color, even if the sample contains more than one type of organism. In contrast, **differential staining** distinguishes organisms based on their interactions with multiple stains. In other words, two organisms in a differentially stained sample may appear to be different colors. Differential staining techniques commonly used in clinical settings include Gram staining, acid-fast staining, endospore staining, flagella staining, and capsule staining. **Figure 2.41** provides more detail on these differential staining techniques.











### Check Your Understanding

- Explain why it is important to fix a specimen before viewing it under a light microscope.
- What types of specimens should be chemically fixed as opposed to heat-fixed?
- Why might an acidic dye react differently with a given specimen than a basic dye?
- Explain the difference between a positive stain and a negative stain.
- Explain the difference between simple and differential staining.

The **Gram stain procedure** is a differential staining procedure that involves multiple steps. It was developed by Danish microbiologist Hans Christian Gram in 1884 as an effective method to distinguish between bacteria with different types of cell walls, and even today it remains one of the most frequently used staining techniques. The steps of the Gram stain procedure are listed below and illustrated in **Figure 2.33**.

1. First, crystal violet, a **primary stain**, is applied to a heat-fixed smear, giving all of the cells a purple color.
2. Next, Gram's iodine, a **mordant**, is added. A mordant is a substance used to set or stabilize stains or dyes; in this case, Gram's iodine acts like a trapping agent that complexes with the crystal violet, making the crystal violet-iodine complex clump and stay contained in thick layers of peptidoglycan in the cell walls.
3. Next, a **decolorizing agent** is added, usually ethanol or an acetone/ethanol solution. Cells that have thick peptidoglycan layers in their cell walls are much less affected by the decolorizing agent; they generally retain the crystal violet dye and remain purple. However, the decolorizing agent more easily washes the dye out of cells with thinner peptidoglycan layers, making them again colorless.
4. Finally, a secondary **counterstain**, usually safranin, is added. This stains the decolorized cells pink and is less noticeable in the cells that still contain the crystal violet dye.




Gram stain process			
Gram staining steps	Cell effects	Gram-positive	Gram-negative
<b>Step 1</b> <b>Crystal violet</b> <i>primary stain added to specimen smear.</i>	Stains cells purple or blue.		
<b>Step 2</b> <b>Iodine</b> <i>mordant makes dye less soluble so it adheres to cell walls.</i>	Cells remain purple or blue.		
<b>Step 3</b> <b>Alcohol</b> <i>decolorizer washes away stain from gram-negative cell walls.</i>	Gram-positive cells remain purple or blue. Gram-negative cells are colorless.		
<b>Step 4</b> <b>Safranin</b> <i>counterstain allows dye adherence to gram-negative cells.</i>	Gram-positive cells remain purple or blue. Gram-negative cells appear pink or red.		

**Figure 2.33** Gram-staining is a differential staining technique that uses a primary stain and a secondary counterstain to distinguish between gram-positive and gram-negative bacteria.

The purple, crystal-violet stained cells are referred to as gram-positive cells, while the red, safranin-dyed cells are gram-negative (**Figure 2.34**). However, there are several important considerations in interpreting the results of a Gram stain. First, older bacterial cells may have damage to their cell walls that causes them to appear gram-negative even if the species is gram-positive. Thus, it is best to use fresh bacterial cultures for Gram staining. Second, errors such as leaving on decolorizer too long can affect the results. In some cases, most cells will appear gram-positive while a few appear gram-negative (as in **Figure 2.34**). This suggests damage to the individual cells or that decolorizer was left on for too long; the cells should still be classified as gram-positive if they are all the same species rather than a mixed culture.

Besides their differing interactions with dyes and decolorizing agents, the chemical differences between gram-positive and gram-negative cells have other implications with clinical relevance. For example, Gram staining can help clinicians classify bacterial pathogens in a sample into categories associated with specific properties. Gram-negative bacteria tend to be more resistant to certain antibiotics than gram-positive bacteria. We will discuss this and other applications of Gram staining in more detail in later chapters.





**Figure 2.34** In this specimen, the gram-positive bacterium *Staphylococcus aureus* retains crystal violet dye even after the decolorizing agent is added. Gram-negative *Escherichia coli*, the most common Gram stain quality-control bacterium, is decolorized, and is only visible after the addition of the pink counterstain safranin. (credit: American Society for Microbiology)



### Check Your Understanding

- Explain the role of Gram's iodine in the Gram stain procedure.
- Explain the role of alcohol in the Gram stain procedure.
- What color are gram-positive and gram-negative cells, respectively, after the Gram stain procedure?


### Clinical Focus

#### Part 3

Viewing Cindy's specimen under the darkfield microscope has provided the technician with some important clues about the identity of the microbe causing her infection. However, more information is needed to make a conclusive diagnosis. The technician decides to make a Gram stain of the specimen. This technique is commonly used as an early step in identifying pathogenic bacteria. After completing the Gram stain procedure, the technician views the slide under the brightfield microscope and sees purple, grape-like clusters of spherical cells (**Figure 2.35**).

- Are these bacteria gram-positive or gram-negative?
- What does this reveal about their cell walls?

Jump to the **next** Clinical Focus box. Go back to the **previous** Clinical Focus box.



**Figure 2.35** (credit: American Society for Microbiology)

## Acid-Fast Stains

Acid-fast staining is another commonly used, differential staining technique that can be an important diagnostic tool. An **acid-fast stain** is able to differentiate two types of gram-positive cells: those that have waxy mycolic acids in their cell walls, and those that do not. Two different methods for acid-fast staining are the **Ziehl-Neelsen technique** and the **Kinyoun technique**. Both use carbolfuchsin as the primary stain. The waxy, acid-fast cells retain the carbolfuchsin even after a decolorizing agent (an acid-alcohol solution) is applied. A secondary counterstain, methylene blue, is then applied, which renders non-acid-fast cells blue.

The fundamental difference between the two carbolfuchsin-based methods is whether heat is used during the primary staining process. The Ziehl-Neelsen method uses heat to infuse the carbolfuchsin into the acid-fast cells, whereas the Kinyoun method does not use heat. Both techniques are important diagnostic tools because a number of specific diseases are caused by acid-fast bacteria (AFB). If AFB are present in a tissue sample, their red or pink color can be seen clearly against the blue background of the surrounding tissue cells (**Figure 2.36**).



### Check Your Understanding


- Why are acid-fast stains useful?

## Micro Connections

### Using Microscopy to Diagnose Tuberculosis

*Mycobacterium tuberculosis*, the bacterium that causes tuberculosis, can be detected in specimens based on the presence of acid-fast bacilli. Often, a smear is prepared from a sample of the patient's sputum and then stained using the Ziehl-Neelsen technique (**Figure 2.36**). If acid-fast bacteria are confirmed, they are generally cultured to make a positive identification. Variations of this approach can be used as a first step in determining whether *M. tuberculosis* or other acid-fast bacteria are present, though samples from elsewhere in the body (such as urine) may contain other *Mycobacterium* species.

An alternative approach for determining the presence of *M. tuberculosis* is immunofluorescence. In this technique, fluorochrome-labeled antibodies bind to *M. tuberculosis*, if present. Antibody-specific fluorescent dyes can be used to view the mycobacteria with a fluorescence microscope.




**Figure 2.36** Ziehl-Neelsen staining has rendered these *Mycobacterium tuberculosis* cells red and the surrounding growth indicator medium blue. (credit: American Society for Microbiology)


## Capsule Staining

Certain bacteria and yeasts have a protective outer structure called a capsule. Since the presence of a capsule is directly related to a microbe's virulence (its ability to cause disease), the ability to determine whether cells in a sample have capsules is an important diagnostic tool. Capsules do not absorb most basic dyes; therefore, a negative staining technique (staining around the cells) is typically used for **capsule staining**. The dye stains the background but does not penetrate the capsules, which appear like halos around the borders of the cell. The specimen does not need to be heat-fixed prior to negative staining.

One common negative staining technique for identifying encapsulated yeast and bacteria is to add a few drops of India ink or nigrosin to a specimen. Other capsular stains can also be used to negatively stain encapsulated cells (**Figure 2.37**). Alternatively, positive and negative staining techniques can be combined to visualize capsules: The positive stain colors the body of the cell, and the negative stain colors the background but not the capsule, leaving halo around each cell.



(a)



(b)

**Figure 2.37** (a) India-ink was used to stain the background around these cells of the yeast *Cryptococcus neoformans*. The halos surrounding the cells are the polysaccharide capsules. (b) Crystal violet and copper sulfate dyes cannot penetrate the encapsulated *Bacillus* cells in this negatively stained sample. Encapsulated cells appear to have a light-blue halo. (credit a: modification of work by American Society for Microbiology; credit b: modification of work by American Society for Microbiology)




## Check Your Understanding

- How does negative staining help us visualize capsules?

## Endospore Staining

Endospores are structures produced within certain bacterial cells that allow them to survive harsh conditions. Gram staining alone cannot be used to visualize endospores, which appear clear when Gram-stained cells are viewed. **Endospore staining** uses two stains to differentiate endospores from the rest of the cell. The Schaeffer-Fulton method (the most commonly used endospore-staining technique) uses heat to push the primary stain (malachite green) into the endospore. Washing with water decolorizes the cell, but the endospore retains the green stain. The cell is then counterstained pink with safranin. The resulting image reveals the shape and location of endospores, if they are present. The green endospores will appear either within the pink vegetative cells or as separate from the pink cells altogether. If no endospores are present, then only the pink vegetative cells will be visible (**Figure 2.38**).



**Figure 2.38** A stained preparation of *Bacillus subtilis* showing endospores as green and the vegetative cells as pink. (credit: American Society for Microbiology)

Endospore-staining techniques are important for identifying *Bacillus* and *Clostridium*, two genera of endospore-producing bacteria that contain clinically significant species. Among others, *B. anthracis* (which causes anthrax) has been of particular interest because of concern that its spores could be used as a bioterrorism agent. *C. difficile* is a particularly important species responsible for the typically hospital-acquired infection known as “C. diff.”




## Check Your Understanding

- Is endospore staining an example of positive, negative, or differential staining?




## Flagella Staining

Flagella (singular: flagellum) are tail-like cellular structures used for locomotion by some bacteria, archaea, and eukaryotes. Because they are so thin, flagella typically cannot be seen under a light microscope without a specialized **flagella staining** technique. Flagella staining thickens the flagella by first applying mordant (generally tannic acid, but sometimes potassium alum), which coats the flagella; then the specimen is stained with pararosaniline (most commonly) or basic fuchsin (**Figure 2.39**).








**Figure 2.39** A flagella stain of *Bacillus cereus*, a common cause of foodborne illness, reveals that the cells have numerous flagella, used for locomotion. (credit: Centers for Disease Control and Prevention)

Though flagella staining is uncommon in clinical settings, the technique is commonly used by microbiologists, since the location and number of flagella can be useful in classifying and identifying bacteria in a sample. When using this technique, it is important to handle the specimen with great care; flagella are delicate structures that can easily be damaged or pulled off, compromising attempts to accurately locate and count the number of flagella.

SIMPLE STAINS				
Stain Type	Specific Dyes	Purpose	Outcome	Sample Images
<b>Basic stains</b>	Methylene blue, crystal violet, malachite green, basic fuchsin, carbolfuchsin, safranin	Stain negatively charged molecules and structures, such as nucleic acids and proteins	Positive stain	
<b>Acidic stains</b>	Eosin, acid fuchsin, rose bengal, Congo red	Stain positively charged molecules and structures, such as proteins	Can be either a positive or negative stain, depending on the cell's chemistry.	
<b>Negative stains</b>	India ink, nigrosin	Stains background, not specimen	Dark background with light specimen	

**Figure 2.40** (credit “basic stains”: modification of work by Centers for Disease Control and Prevention; credit “Acidic stains”: modification of work by Roberto Danovaro, Antonio Pusceddu, Cristina Gambi, Iben Heiner, Reinhardt Moberg Kristensen; credit “Negative stains”: modification of work by Anh-Hue Tu)



DIFFERENTIAL STAINS				
Stain Type	Specific Dyes	Purpose	Outcome	Sample Images
<b>Gram stain</b>	Uses crystal violet, Gram's iodine, ethanol (decolorizer), and safranin	Used to distinguish cells by cell-wall type (gram-positive, gram-negative)	Gram-positive cells stain purple/violet. Gram-negative cells stain pink.	
<b>Acid-fast stain</b>	After staining with basic fuchsin, acid-fast bacteria resist decolorization by acid-alcohol. Non acid-fast bacteria are counterstained with methylene blue.	Used to distinguish acid-fast bacteria such as <i>M. tuberculosis</i> , from non-acid-fast cells	Acid-fast bacteria are red; non-acid-fast cells are blue.	
<b>Endospore stain</b>	Uses heat to stain endospores with malachite green (Schaeffer-Fulton procedure), then cell is washed and counterstained with safranin.	Used to distinguish organisms with endospores from those without; used to study the endospore.	Endospores appear bluish-green; other structures appear pink to red.	
<b>Flagella stain</b>	Flagella are coated with a tannic acid or potassium alum mordant, then stained using either pararosaniline or basic fuchsin.	Used to view and study flagella in bacteria that have them.	Flagella are visible if present.	
<b>Capsule stain</b>	Negative staining with India ink or nigrosin is used to stain the background, leaving a clear area of the cell and the capsule. Counterstaining can be used to stain the cell while leaving the capsule clear.	Used to distinguish cells with capsules from those without.	Capsules appear clear or as halos if present.	

**Figure 2.41** (credit “Gram stain”: modification of work by American Society for Microbiology; credit “Acid-fast stain”: modification of work by American Society for Microbiology; credit “Endospore stain”: modification of work by American Society for Microbiology; credit “Capsule stain” : modification of work by American Society for Microbiology; credit “Flagella stain”: modification of work by Centers for Disease Control and Prevention)

## Preparing Specimens for Electron Microscopy

Samples to be analyzed using a TEM must have very thin sections. But cells are too soft to cut thinly, even with diamond knives. To cut cells without damage, the cells must be embedded in plastic resin and then dehydrated through a series of soaks in ethanol solutions (50%, 60%, 70%, and so on). The ethanol replaces the water in the cells, and the resin dissolves in ethanol and enters the cell, where it solidifies. Next, **thin sections** are cut using a specialized device called an **ultramicrotome** (Figure 2.42). Finally, samples are fixed to fine copper wire or carbon-fiber grids and stained—not with colored dyes, but with substances like uranyl acetate or osmium tetroxide, which contain electron-dense heavy metal atoms.



**Figure 2.42** (a) An ultramicrotome used to prepare specimens for a TEM. (b) A technician uses an ultramicrotome to slice a specimen into thin sections. (credit a: modification of work by “Frost Museum”/Flickr; credit b: modification of work by “U.S. Fish and Wildlife Service Northeast Region”/Flickr)

When samples are prepared for viewing using an SEM, they must also be dehydrated using an ethanol series. However, they must be even drier than is necessary for a TEM. Critical point drying with inert liquid carbon dioxide under pressure is used to displace the water from the specimen. After drying, the specimens are sputter-coated with metal by knocking atoms off of a palladium target, with energetic particles. Sputter-coating prevents specimens from becoming charged by the SEM’s electron beam.




### Check Your Understanding

- Why is it important to dehydrate cells before examining them under an electron microscope?
- Name the device that is used to create thin sections of specimens for electron microscopy.

### Micro Connections


#### Using Microscopy to Diagnose Syphilis

The causative agent of syphilis is *Treponema pallidum*, a flexible, spiral cell (spirochete) that can be very thin ( $<0.15\ \mu\text{m}$ ) and match the refractive index of the medium, making it difficult to view using brightfield microscopy. Additionally, this species has not been successfully cultured in the laboratory on an artificial medium; therefore, diagnosis depends upon successful identification using microscopic techniques and serology (analysis of body fluids, often looking for antibodies to a pathogen). Since fixation and staining would kill the cells, darkfield microscopy is typically used for observing live specimens and viewing their movements. However, other approaches can also be used. For example, the cells can be thickened with silver particles (in tissue sections) and observed using a light microscope. It is also possible to use fluorescence or electron microscopy to view *Treponema* (Figure 2.43).



**Figure 2.43** (a) Living, unstained *Treponema pallidum* spirochetes can be viewed under a darkfield microscope. (b) In this brightfield image, a modified Steiner silver stain is used to visualize *T. pallidum* spirochetes. Though the stain kills the cells, it increases the contrast to make them more visible. (c) While not used for standard diagnostic testing, *T. pallidum* can also be examined using scanning electron microscopy. (credit a: modification of work by Centers for Disease Control and Prevention; credit b: modification of work by Centers for Disease Control and Prevention; credit c: modification of work by Centers for Disease Control and Prevention)

In clinical settings, indirect immunofluorescence is often used to identify *Treponema*. A primary, unstained antibody attaches directly to the pathogen surface, and secondary antibodies “tagged” with a fluorescent stain attach to the primary antibody. Multiple secondary antibodies can attach to each primary antibody, amplifying the amount of stain attached to each *Treponema* cell, making them easier to spot (**Figure 2.44**).



**Figure 2.44** Indirect immunofluorescence can be used to identify *T. pallidum*, the causative agent of syphilis, in a specimen.

## Preparation and Staining for Other Microscopes

Samples for fluorescence and confocal microscopy are prepared similarly to samples for light microscopy, except that the dyes are fluorochromes. Stains are often diluted in liquid before applying to the slide. Some dyes attach to an antibody to stain specific proteins on specific types of cells (immunofluorescence); others may attach to DNA molecules in a process called fluorescence in situ hybridization (FISH), causing cells to be stained based on whether they have a specific DNA sequence.

Sample preparation for two-photon microscopy is similar to fluorescence microscopy, except for the use of infrared dyes. Specimens for STM need to be on a very clean and atomically smooth surface. They are often mica coated with Au(111). Toluene vapor is a common fixative.



## Check Your Understanding

- What is the main difference between preparing a sample for fluorescence microscopy versus light microscopy?

## Link to Learning



Cornell University's **Case Studies in Microscopy** (<http://www.openstaxcollege.org/l/22cornellstud>) offers a series of clinical problems based on real-life events. Each case study walks you through a clinical problem using appropriate techniques in microscopy at each step.

## Clinical Focus

### Resolution

From the results of the Gram stain, the technician now knows that Cindy's infection is caused by spherical, gram-positive bacteria that form grape-like clusters, which is typical of staphylococcal bacteria. After some additional testing, the technician determines that these bacteria are the medically important species known as *Staphylococcus aureus*, a common culprit in wound infections. Because some strains of *S. aureus* are resistant to many antibiotics, skin infections may spread to other areas of the body and become serious, sometimes even resulting in amputations or death if the correct antibiotics are not used.

After testing several antibiotics, the lab is able to identify one that is effective against this particular strain of *S. aureus*. Cindy's doctor quickly prescribes the medication and emphasizes the importance of taking the entire course of antibiotics, even if the infection appears to clear up before the last scheduled dose. This reduces the risk that any especially resistant bacteria could survive, causing a second infection or spreading to another person.

Go back to the **previous** Clinical Focus box.

## Eye on Ethics



### Microscopy and Antibiotic Resistance

As the use of antibiotics has proliferated in medicine, as well as agriculture, microbes have evolved to become more resistant. Strains of bacteria such as methicillin-resistant *S. aureus* (MRSA), which has developed a high level of resistance to many antibiotics, are an increasingly worrying problem, so much so that research is underway to develop new and more diversified antibiotics.

Fluorescence microscopy can be useful in testing the effectiveness of new antibiotics against resistant strains like MRSA. In a test of one new antibiotic derived from a marine bacterium, MC21-A (bromophene), researchers used the fluorescent dye SYTOX Green to stain samples of MRSA. SYTOX Green is often used



to distinguish dead cells from living cells, with fluorescence microscopy. Live cells will not absorb the dye, but cells killed by an antibiotic will absorb the dye, since the antibiotic has damaged the bacterial cell membrane. In this particular case, MRSA bacteria that had been exposed to MC21-A did, indeed, appear green under the fluorescence microscope, leading researchers to conclude that it is an effective antibiotic against MRSA.

Of course, some argue that developing new antibiotics will only lead to even more antibiotic-resistant microbes, so-called superbugs that could spawn epidemics before new treatments can be developed. For this reason, many health professionals are beginning to exercise more discretion in prescribing antibiotics. Whereas antibiotics were once routinely prescribed for common illnesses without a definite diagnosis, doctors and hospitals are much more likely to conduct additional testing to determine whether an antibiotic is necessary and appropriate before prescribing.

A sick patient might reasonably object to this stingy approach to prescribing antibiotics. To the patient who simply wants to feel better as quickly as possible, the potential benefits of taking an antibiotic may seem to outweigh any immediate health risks that might occur if the antibiotic is ineffective. But at what point do the risks of widespread antibiotic use supersede the desire to use them in individual cases?

## Summary

---

### 2.1 The Properties of Light

- Light waves interacting with materials may be **reflected**, **absorbed**, or **transmitted**, depending on the properties of the material.
- Light waves can interact with each other (**interference**) or be distorted by interactions with small objects or openings (**diffraction**).
- **Refraction** occurs when light waves change speed and direction as they pass from one medium to another. Differences in the **refraction indices** of two materials determine the magnitude of directional changes when light passes from one to the other.
- A **lens** is a medium with a curved surface that refracts and focuses light to produce an image.
- Visible light is part of the **electromagnetic spectrum**; light waves of different frequencies and wavelengths are distinguished as colors by the human eye.
- A prism can separate the colors of white light (**dispersion**) because different frequencies of light have different refractive indices for a given material.
- **Fluorescent dyes** and **phosphorescent** materials can effectively transform nonvisible electromagnetic radiation into visible light.
- The power of a microscope can be described in terms of its **magnification** and **resolution**.
- Resolution can be increased by shortening wavelength, increasing the **numerical aperture** of the lens, or using stains that enhance contrast.

### 2.2 Peering Into the Invisible World

- **Antonie van Leeuwenhoek** is credited with the first observation of microbes, including protists and bacteria, with simple microscopes that he made.
- **Robert Hooke** was the first to describe what we now call cells.
- **Simple microscopes** have a single lens, while **compound microscopes** have multiple lenses.

### 2.3 Instruments of Microscopy

- Numerous types of microscopes use various technologies to generate micrographs. Most are useful for a particular type of specimen or application.



- **Light microscopy** uses lenses to focus light on a specimen to produce an image. Commonly used light microscopes include **brightfield, darkfield, phase-contrast, differential interference contrast, fluorescence, confocal,** and **two-photon** microscopes.
- **Electron microscopy** focuses electrons on the specimen using magnets, producing much greater magnification than light microscopy. The **transmission electron microscope (TEM)** and **scanning electron microscope (SEM)** are two common forms.
- **Scanning probe microscopy** produces images of even greater magnification by measuring feedback from sharp probes that interact with the specimen. Probe microscopes include the **scanning tunneling microscope (STM)** and the **atomic force microscope (AFM)**.

## 2.4 Staining Microscopic Specimens

- Samples must be properly prepared for microscopy. This may involve **staining, fixation,** and/or cutting **thin sections**.
- A variety of staining techniques can be used with light microscopy, including **Gram staining, acid-fast staining, capsule staining, endospore staining,** and **flagella staining**.
- Samples for TEM require very thin sections, whereas samples for SEM require sputter-coating.
- Preparation for fluorescence microscopy is similar to that for light microscopy, except that fluorochromes are used.

## Review Questions

### Multiple Choice

- Which of the following has the highest energy?
  - light with a long wavelength
  - light with an intermediate wavelength
  - light with a short wavelength
  - It is impossible to tell from the information given.
- You place a specimen under the microscope and notice that parts of the specimen begin to emit light immediately. These materials can be described as \_\_\_\_\_.
  - fluorescent
  - phosphorescent
  - transparent
  - opaque
- Who was the first to describe “cells” in dead cork tissue?
  - Hans Janssen
  - Zaccharias Janssen
  - Antonie van Leeuwenhoek
  - Robert Hooke
- Who is the probable inventor of the compound microscope?
  - Girolamo Fracastoro
  - Zaccharias Janssen
  - Antonie van Leeuwenhoek
  - Robert Hooke
- Which would be the best choice for viewing internal structures of a living protist such as a *Paramecium*?
  - a brightfield microscope with a stain
  - a brightfield microscope without a stain
  - a darkfield microscope
  - a transmission electron microscope
- Which type of microscope is especially useful for viewing thick structures such as biofilms?
  - a transmission electron microscope
  - a scanning electron microscope
  - a phase-contrast microscope
  - a confocal scanning laser microscope
  - an atomic force microscope
- Which type of microscope would be the best choice for viewing very small surface structures of a cell?
  - a transmission electron microscope
  - a scanning electron microscope
  - a brightfield microscope
  - a darkfield microscope
  - a phase-contrast microscope
- What type of microscope uses an annular stop?
  - a transmission electron microscope
  - a scanning electron microscope
  - a brightfield microscope
  - a darkfield microscope
  - a phase-contrast microscope

9. What type of microscope uses a cone of light so that light only hits the specimen indirectly, producing a darker image on a brighter background?

- a. a transmission electron microscope
- b. a scanning electron microscope
- c. a brightfield microscope
- d. a darkfield microscope
- e. a phase-contrast microscope

10. What mordant is used in Gram staining?

- a. crystal violet
- b. safranin
- c. acid-alcohol
- d. iodine

11. What is one difference between specimen preparation for a transmission electron microscope (TEM) and preparation for a scanning electron microscope (SEM)?

- a. Only the TEM specimen requires sputter coating.
- b. Only the SEM specimen requires sputter-coating.
- c. Only the TEM specimen must be dehydrated.
- d. Only the SEM specimen must be dehydrated.

#### Fill in the Blank

12. When you see light bend as it moves from air into water, you are observing \_\_\_\_\_.

13. A microscope that uses multiple lenses is called a \_\_\_\_\_ microscope.

14. Chromophores that absorb and then emit light are called \_\_\_\_\_.

15. In a(n) \_\_\_\_\_ microscope, a probe located just above the specimen moves up and down in response to forces between the atoms and the tip of the probe.

16. What is the total magnification of a specimen that is being viewed with a standard ocular lens and a 40× objective lens?

17. Ziehl-Neelsen staining, a type of \_\_\_\_\_ staining, is diagnostic for *Mycobacterium tuberculosis*.

18. The \_\_\_\_\_ is used to differentiate bacterial cells based on the components of their cell walls.

#### Short Answer

19. Explain how a prism separates white light into different colors.


20. Why is Antonie van Leeuwenhoek's work much better known than that of Zaccharias Janssen?

21. Why did the cork cells observed by Robert Hooke appear to be empty, as opposed to being full of other structures?

22. What is the function of the condenser in a brightfield microscope?

#### Art Connection

23. Label each component of the brightfield microscope.



24. How could you identify whether a particular bacterial sample contained specimens with mycolic acid-rich cell walls?

### Critical Thinking

25. In **Figure 2.7**, which of the following has the lowest energy?

- visible light
- X-rays
- ultraviolet rays
- infrared rays

26. When focusing a light microscope, why is it best to adjust the focus using the coarse focusing knob before using the fine focusing knob?


27. You need to identify structures within a cell using a microscope. However, the image appears very blurry even though you have a high magnification. What are some things that you could try to improve the resolution of the image? Describe the most basic factors that affect resolution when you first put the slide onto the stage; then consider more specific factors that could affect resolution for 40 $\times$  and 100 $\times$  lenses.

28. You use the Gram staining procedure to stain an L-form bacterium (a bacterium that lacks a cell wall). What color will the bacterium be after the staining procedure is finished?



## Chapter 3

# The Cell



**Figure 3.1** Microorganisms vary visually in their size and shape, as can be observed microscopically; but they also vary in invisible ways, such as in their metabolic capabilities. (credit “microbacterium tuberculosis bacteria”: modification of work by NIAID)

### Chapter Outline

- 3.1 Spontaneous Generation
- 3.2 Foundations of Modern Cell Theory
- 3.3 Unique Characteristics of Prokaryotic Cells
- 3.4 Unique Characteristics of Eukaryotic Cells

## Introduction

Life takes many forms, from giant redwood trees towering hundreds of feet in the air to the tiniest known microbes, which measure only a few billionths of a meter. Humans have long pondered life’s origins and debated the defining characteristics of life, but our understanding of these concepts has changed radically since the invention of the microscope. In the 17th century, observations of microscopic life led to the development of the cell theory: the idea that the fundamental unit of life is the cell, that all organisms contain at least one cell, and that cells only come from other cells.

Despite sharing certain characteristics, cells may vary significantly. The two main types of cells are prokaryotic cells (lacking a nucleus) and eukaryotic cells (containing a well-organized, membrane-bound nucleus). Each type of cell exhibits remarkable variety in structure, function, and metabolic activity (**Figure 3.1**). This chapter will focus on the historical discoveries that have shaped our current understanding of microbes, including their origins and their role in human disease. We will then explore the distinguishing structures found in prokaryotic and eukaryotic cells.



## 3.1 Spontaneous Generation

### Learning Objectives

- Explain the theory of spontaneous generation and why people once accepted it as an explanation for the existence of certain types of organisms
- Explain how certain individuals (van Helmont, Redi, Needham, Spallanzani, and Pasteur) tried to prove or disprove spontaneous generation

Humans have been asking for millennia: Where does new life come from? Religion, philosophy, and science have all wrestled with this question. One of the oldest explanations was the theory of spontaneous generation, which can be traced back to the ancient Greeks and was widely accepted through the Middle Ages.

### The Theory of Spontaneous Generation

The Greek philosopher Aristotle (384–322 BC) was one of the earliest recorded scholars to articulate the theory of **spontaneous generation**, the notion that life can arise from nonliving matter. Aristotle proposed that life arose from nonliving material if the material contained *pneuma* (“vital heat”). As evidence, he noted several instances of the appearance of animals from environments previously devoid of such animals, such as the seemingly sudden appearance of fish in a new puddle of water.<sup>[1]</sup>

This theory persisted into the 17th century, when scientists undertook additional experimentation to support or disprove it. By this time, the proponents of the theory cited how frogs simply seem to appear along the muddy banks of the Nile River in Egypt during the annual flooding. Others observed that mice simply appeared among grain stored in barns with thatched roofs. When the roof leaked and the grain molded, mice appeared. Jan Baptista van Helmont, a 17th century Flemish scientist, proposed that mice could arise from rags and wheat kernels left in an open container for 3 weeks. In reality, such habitats provided ideal food sources and shelter for mouse populations to flourish.

However, one of van Helmont’s contemporaries, Italian physician Francesco Redi (1626–1697), performed an experiment in 1668 that was one of the first to refute the idea that maggots (the larvae of flies) spontaneously generate on meat left out in the open air. He predicted that preventing flies from having direct contact with the meat would also prevent the appearance of maggots. Redi left meat in each of six containers (**Figure 3.2**): Two were open to the air, two were covered with gauze, and two were tightly sealed. His hypothesis was supported when maggots developed in the uncovered jars, but no maggots appeared in either the gauze-covered or the tightly sealed jars. He concluded that maggots could only form when flies were allowed to lay eggs in the meat, and that the maggots were the offspring of flies, not the product of spontaneous generation.

### Clinical Focus


#### Part 1

Barbara is a 19-year-old college student living in the dormitory. In January, she came down with a sore throat, headache, mild fever, chills, and a violent but unproductive (i.e., no mucus) cough. To treat these symptoms, Barbara began taking an over-the-counter cold medication, which did not seem to work. In fact, over the next few days, while some of Barbara’s symptoms began to resolve, her cough and fever persisted, and she felt very tired and weak.

- What types of respiratory disease may be responsible?

Jump to the **next** Clinical Focus box

1. K. Zwier. “Aristotle on Spontaneous Generation.” <http://www.sju.edu/int/academics/cas/resources/gppc/pdf/Karen%20R.%20Zwier.pdf>



**Figure 3.2** Francesco Redi's experimental setup consisted of an open container, a container sealed with a cork top, and a container covered in mesh that let in air but not flies. Maggots only appeared on the meat in the open container. However, maggots were also found on the gauze of the gauze-covered container.

In 1745, John Needham (1713–1781) published a report of his own experiments, in which he briefly boiled broth infused with plant or animal matter, hoping to kill all preexisting microbes.<sup>[2]</sup> He then sealed the flasks. After a few days, Needham observed that the broth had become cloudy and a single drop contained numerous microscopic creatures. He argued that the new microbes must have arisen spontaneously. In reality, however, he likely did not boil the broth enough to kill all preexisting microbes.

Lazzaro Spallanzani (1729–1799) did not agree with Needham's conclusions, however, and performed hundreds of carefully executed experiments using heated broth.<sup>[3]</sup> As in Needham's experiment, broth in sealed jars and unsealed jars was infused with plant and animal matter. Spallanzani's results contradicted the findings of Needham: Heated but sealed flasks remained clear, without any signs of spontaneous growth, unless the flasks were subsequently opened to the air. This suggested that microbes were introduced into these flasks from the air. In response to Spallanzani's findings, Needham argued that life originates from a "life force" that was destroyed during Spallanzani's extended boiling. Any subsequent sealing of the flasks then prevented new life force from entering and causing spontaneous generation (**Figure 3.3**).

2. E. Capanna. "Lazzaro Spallanzani: At the Roots of Modern Biology." *Journal of Experimental Zoology* 285 no. 3 (1999):178–196.

3. R. Mancini, M. Nigro, G. Ippolito. "Lazzaro Spallanzani and His Refutation of the Theory of Spontaneous Generation." *Le Infezioni in Medicina* 15 no. 3 (2007):199–206.



**Figure 3.3** (a) Francesco Redi, who demonstrated that maggots were the offspring of flies, not products of spontaneous generation. (b) John Needham, who argued that microbes arose spontaneously in broth from a “life force.” (c) Lazzaro Spallanzani, whose experiments with broth aimed to disprove those of Needham.



### Check Your Understanding

- Describe the theory of spontaneous generation and the observations used as evidence to support it.
- Explain how the experiments of Redi and Spallanzani challenged the theory of spontaneous generation.


## Disproving Spontaneous Generation

The debate over spontaneous generation continued well into the 19th century, with scientists serving as proponents of both sides. To settle the debate, the Paris Academy of Sciences offered a prize for resolution of the problem. Louis Pasteur, a prominent French chemist who had been studying microbial fermentation and the causes of wine spoilage, accepted the challenge and would ultimately win both the debate and the Academy’s prize (**Figure 3.4**). In 1858, Pasteur filtered air through a gun-cotton filter and, upon microscopic examination of the cotton, found it full of microorganisms, suggesting that the exposure of a broth to air was not introducing a “life force” to the broth but rather airborne microorganisms.

Later, Pasteur made a series of flasks with long, twisted necks (“swan-neck” flasks), in which he boiled broth to sterilize it (**Figure 3.4**). His design allowed air inside the flasks to be exchanged with air from the outside, but prevented the introduction of any airborne microorganisms, which would get caught in the twists and bends of the flasks’ necks. If a life force besides the airborne microorganisms were responsible for microbial growth within the sterilized flasks, it would have access to the broth, whereas the microorganisms would not. He correctly predicted that sterilized broth in his swan-neck flasks would remain sterile as long as the swan necks remained intact. However, should the necks be broken, microorganisms would be introduced, contaminating the flasks and allowing microbial growth within the broth.

Pasteur’s set of experiments irrefutably disproved the theory of spontaneous generation and earned him the prestigious Alhumbert Prize from the Paris Academy of Sciences in 1862. In a subsequent lecture in 1864, Pasteur articulated “*Omne vivum ex vivo*” (“Life only comes from life”). In this lecture, Pasteur recounted his famous swan-neck flask experiment, stating that “...life is a germ and a germ is life. Never will the doctrine of spontaneous generation recover from the mortal blow of this simple experiment.”<sup>[4]</sup> To Pasteur’s credit, it never has.

4. R. Vallery-Radot. *The Life of Pasteur*, trans. R.L. Devonshire. New York: McClure, Phillips and Co, 1902, 1:142.



**Figure 3.4** (a) French scientist Louis Pasteur, who definitively refuted the long-disputed theory of spontaneous generation. (b) The unique swan-neck feature of the flasks used in Pasteur's experiment allowed air to enter the flask but prevented the entry of bacterial and fungal spores. (c) Pasteur's experiment consisted of two parts. In the first part, the broth in the flask was boiled to sterilize it. When this broth was cooled, it remained free of contamination. In the second part of the experiment, the flask was boiled and then the neck was broken off. The broth in this flask became contaminated.



### Check Your Understanding

- How did Pasteur's experimental design allow air to enter, but not microbes, and why was this important?
- What was the control group in Pasteur's experiment and what did it show?

## 3.2 Foundations of Modern Cell Theory

### Learning Objectives

- Explain the key points of cell theory and the individual contributions of Hooke, Schleiden, Schwann, Remak, and Virchow.
- Explain the key points of endosymbiotic theory and cite the evidence that supports this concept.
- Explain the contributions of Semmelweis, Snow, Pasteur, Lister, and Koch to the development of germ theory.


While some scientists were arguing over the theory of spontaneous generation, other scientists were making discoveries leading to a better understanding of what we now call the cell theory. Modern cell theory has two basic tenets:

- All cells only come from other cells (the principle of biogenesis).
- Cells are the fundamental units of organisms.

Today, these tenets are fundamental to our understanding of life on earth. However, modern cell theory grew out of the collective work of many scientists.

### The Origins of Cell Theory

The English scientist Robert Hooke first used the term “cells” in 1665 to describe the small chambers within cork that he observed under a microscope of his own design. To Hooke, thin sections of cork resembled “Honey-comb,” or “small Boxes or Bladders of Air.” He noted that each “Cavern, Bubble, or Cell” was distinct from the others (**Figure 3.5**). At the time, Hooke was not aware that the cork cells were long dead and, therefore, lacked the internal structures found within living cells.



**Figure 3.5** Robert Hooke (1635–1703) was the first to describe cells based upon his microscopic observations of cork. This illustration was published in his work *Micrographia*.

Despite Hooke’s early description of cells, their significance as the fundamental unit of life was not yet recognized. Nearly 200 years later, in 1838, Matthias Schleiden (1804–1881), a German botanist who made extensive microscopic observations of plant tissues, described them as being composed of cells. Visualizing plant cells was relatively easy because plant cells are clearly separated by their thick cell walls. Schleiden believed that cells formed through crystallization, rather than cell division.



Theodor Schwann (1810–1882), a noted German physiologist, made similar microscopic observations of animal tissue. In 1839, after a conversation with Schleiden, Schwann realized that similarities existed between plant and animal tissues. This laid the foundation for the idea that cells are the fundamental components of plants and animals.

In the 1850s, two Polish scientists living in Germany pushed this idea further, culminating in what we recognize today as the modern cell theory. In 1852, Robert Remak (1815–1865), a prominent neurologist and embryologist, published convincing evidence that cells are derived from other cells as a result of cell division. However, this idea was questioned by many in the scientific community. Three years later, Rudolf Virchow (1821–1902), a well-respected pathologist, published an editorial essay entitled “Cellular Pathology,” which popularized the concept of cell theory using the Latin phrase *omnis cellula a cellula* (“all cells arise from cells”), which is essentially the second tenet of modern cell theory.<sup>[5]</sup> Given the similarity of Virchow’s work to Remak’s, there is some controversy as to which scientist should receive credit for articulating cell theory. See the following Eye on Ethics feature for more about this controversy.

### Eye on Ethics



#### Science and Plagiarism

Rudolf Virchow, a prominent, Polish-born, German scientist, is often remembered as the “Father of Pathology.” Well known for innovative approaches, he was one of the first to determine the causes of various diseases by examining their effects on tissues and organs. He was also among the first to use animals in his research and, as a result of his work, he was the first to name numerous diseases and created many other medical terms. Over the course of his career, he published more than 2,000 papers and headed various important medical facilities, including the Charité – Universitätsmedizin Berlin, a prominent Berlin hospital and medical school. But he is, perhaps, best remembered for his 1855 editorial essay titled “Cellular Pathology,” published in *Archiv für Pathologische Anatomie und Physiologie*, a journal that Virchow himself cofounded and still exists today.

Despite his significant scientific legacy, there is some controversy regarding this essay, in which Virchow proposed the central tenet of modern cell theory—that all cells arise from other cells. Robert Remak, a former colleague who worked in the same laboratory as Virchow at the University of Berlin, had published the same idea 3 years before. Though it appears Virchow was familiar with Remak’s work, he neglected to credit Remak’s ideas in his essay. When Remak wrote a letter to Virchow pointing out similarities between Virchow’s ideas and his own, Virchow was dismissive. In 1858, in the preface to one of his books, Virchow wrote that his 1855 publication was just an editorial piece, not a scientific paper, and thus there was no need to cite Remak’s work.


By today’s standards, Virchow’s editorial piece would certainly be considered an act of plagiarism, since he presented Remak’s ideas as his own. However, in the 19th century, standards for academic integrity were much less clear. Virchow’s strong reputation, coupled with the fact that Remak was a Jew in a somewhat anti-Semitic political climate, shielded him from any significant repercussions. Today, the process of peer review and the ease of access to the scientific literature help discourage plagiarism. Although scientists are still motivated to publish original ideas that advance scientific knowledge, those who would consider plagiarizing are well aware of the serious consequences.

In academia, plagiarism represents the theft of both individual thought and research—an offense that can destroy reputations and end careers.<sup>[6] [7] [8] [9]</sup>

5. M. Schultz. “Rudolph Virchow.” *Emerging Infectious Diseases* 14 no. 9 (2008):1480–1481.

6. B. Kisch. “Forgotten Leaders in Modern Medicine, Valentin, Gouby, Remak, Auerbach.” *Transactions of the American Philosophical Society* 44 (1954):139–317.

7. H. Harris. *The Birth of the Cell*. New Haven, CT: Yale University Press, 2000:133.



**Figure 3.6** (a) Rudolf Virchow (1821–1902) popularized the cell theory in an 1855 essay entitled “Cellular Pathology.” (b) The idea that all cells originate from other cells was first published in 1852 by his contemporary and former colleague Robert Remak (1815–1865).



### Check Your Understanding

- What are the key points of the cell theory?
- What contributions did Rudolf Virchow and Robert Remak make to the development of the cell theory?

## Endosymbiotic Theory

As scientists were making progress toward understanding the role of cells in plant and animal tissues, others were examining the structures within the cells themselves. In 1831, Scottish botanist Robert Brown (1773–1858) was the first to describe observations of nuclei, which he observed in plant cells. Then, in the early 1880s, German botanist Andreas Schimper (1856–1901) was the first to describe the chloroplasts of plant cells, identifying their role in starch formation during photosynthesis and noting that they divided independent of the nucleus.

Based upon the chloroplasts’ ability to reproduce independently, Russian botanist Konstantin Mereschkowski (1855–1921) suggested in 1905 that chloroplasts may have originated from ancestral photosynthetic bacteria living symbiotically inside a eukaryotic cell. He proposed a similar origin for the nucleus of plant cells. This was the first articulation of the endosymbiotic hypothesis, and would explain how eukaryotic cells evolved from ancestral bacteria.

Mereschkowski’s endosymbiotic hypothesis was furthered by American anatomist Ivan Wallin (1883–1969), who began to experimentally examine the similarities between mitochondria, chloroplasts, and bacteria—in other words, to put the endosymbiotic hypothesis to the test using objective investigation. Wallin published a series of papers in the 1920s supporting the endosymbiotic hypothesis, including a 1926 publication co-authored with Mereschkowski. Wallin claimed he could culture mitochondria outside of their eukaryotic host cells. Many scientists dismissed his cultures of mitochondria as resulting from bacterial contamination. Modern genome sequencing work supports the

8. C. Webster (ed.). *Biology, Medicine and Society 1840-1940*. Cambridge, UK; Cambridge University Press, 1981:118–119.

9. C. Zuchora-Walske. *Key Discoveries in Life Science*. Minneapolis, MN: Lerner Publishing, 2015:12–13.

dissenting scientists by showing that much of the genome of mitochondria had been transferred to the host cell's nucleus, preventing the mitochondria from being able to live on their own.<sup>[10] [11]</sup>

Wallin's ideas regarding the endosymbiotic hypothesis were largely ignored for the next 50 years because scientists were unaware that these organelles contained their own DNA. However, with the discovery of mitochondrial and chloroplast DNA in the 1960s, the endosymbiotic hypothesis was resurrected. Lynn Margulis (1938–2011), an American geneticist, published her ideas regarding the endosymbiotic hypothesis of the origins of mitochondria and chloroplasts in 1967.<sup>[12]</sup> In the decade leading up to her publication, advances in microscopy had allowed scientists to differentiate prokaryotic cells from eukaryotic cells. In her publication, Margulis reviewed the literature and argued that the eukaryotic organelles such as mitochondria and chloroplasts are of prokaryotic origin. She presented a growing body of microscopic, genetic, molecular biology, fossil, and geological data to support her claims.

Again, this hypothesis was not initially popular, but mounting genetic evidence due to the advent of DNA sequencing supported the **endosymbiotic theory**, which is now defined as the theory that mitochondria and chloroplasts arose as a result of prokaryotic cells establishing a symbiotic relationship within a eukaryotic host. With Margulis' initial endosymbiotic theory gaining wide acceptance (**Figure 3.7**), she expanded on the theory in her 1981 book *Symbiosis in Cell Evolution*. In it, she explains how endosymbiosis is a major driving factor in the evolution of organisms. More recent genetic sequencing and phylogenetic analysis show that mitochondrial DNA and chloroplast DNA are highly related to their bacterial counterparts, both in DNA sequence and chromosome structure. However, mitochondrial DNA and chloroplast DNA are reduced compared with nuclear DNA because many of the genes have moved from the organelles into the host cell's nucleus. Additionally, mitochondrial and chloroplast ribosomes are structurally similar to bacterial ribosomes, rather than to the eukaryotic ribosomes of their hosts. Last, the binary fission of these organelles strongly resembles the binary fission of bacteria, as compared with mitosis performed by eukaryotic cells. Since Margulis' original proposal, scientists have observed several examples of bacterial endosymbionts in modern-day eukaryotic cells. Examples include the endosymbiotic bacteria found within the guts of certain insects, such as cockroaches,<sup>[13]</sup> and photosynthetic bacteria-like organelles found in protists.<sup>[14]</sup>

---

10. T. Embley, W. Martin. "Eukaryotic Evolution, Changes, and Challenges." *Nature* Vol. 440 (2006):623–630.


11. O.G. Berg, C.G. Kurland. "Why Mitochondrial Genes Are Most Often Found in Nuclei." *Molecular Biology and Evolution* 17 no. 6 (2000):951–961.

12. L. Sagan. "On the Origin of Mitosing Cells." *Journal of Theoretical Biology* 14 no. 3 (1967):225–274.

13. A.E. Douglas. "The Microbial Dimension in Insect Nutritional Ecology." *Functional Ecology* 23 (2009):38–47.

14. J.M. Jaynes, L.P. Vernon. "The Cyanelle of *Cyanophora paradoxa*: Almost a Cyanobacterial Chloroplast." *Trends in Biochemical Sciences* 7 no. 1 (1982):22–24.

### The Endosymbiotic Theory



**Figure 3.7** According to the endosymbiotic theory, mitochondria and chloroplasts are each derived from the uptake of bacteria. These bacteria established a symbiotic relationship with their host cell that eventually led to the bacteria evolving into mitochondria and chloroplasts.



### Check Your Understanding

- What does the modern endosymbiotic theory state?
- What evidence supports the endosymbiotic theory?

## The Germ Theory of Disease


Prior to the discovery of microbes during the 17th century, other theories circulated about the origins of disease. For example, the ancient Greeks proposed the miasma theory, which held that disease originated from particles emanating from decomposing matter, such as that in sewage or cesspits. Such particles infected humans in close proximity to the rotting material. Diseases including the Black Death, which ravaged Europe's population during the Middle Ages, were thought to have originated in this way.

In 1546, Italian physician Girolamo Fracastoro proposed, in his essay *De Contagione et Contagiosis Morbis*, that seed-like spores may be transferred between individuals through direct contact, exposure to contaminated clothing, or through the air. We now recognize Fracastoro as an early proponent of the **germ theory of disease**, which states that diseases may result from microbial infection. However, in the 16th century, Fracastoro's ideas were not widely accepted and would be largely forgotten until the 19th century.

In 1847, Hungarian obstetrician Ignaz Semmelweis observed that mothers who gave birth in hospital wards staffed by physicians and medical students were more likely to suffer and die from puerperal fever after childbirth (10%–20% mortality rate) than were mothers in wards staffed by midwives (1% mortality rate). Semmelweis observed medical students performing autopsies and then subsequently carrying out vaginal examinations on living patients without washing their hands in between. He suspected that the students carried disease from the autopsies to the patients they

examined. His suspicions were supported by the untimely death of a friend, a physician who contracted a fatal wound infection after a postmortem examination of a woman who had died of a puerperal infection. The dead physician's wound had been caused by a scalpel used during the examination, and his subsequent illness and death closely paralleled that of the dead patient.

Although Semmelweis did not know the true cause of puerperal fever, he proposed that physicians were somehow transferring the causative agent to their patients. He suggested that the number of puerperal fever cases could be reduced if physicians and medical students simply washed their hands with chlorinated lime water before and after examining every patient. When this practice was implemented, the maternal mortality rate in mothers cared for by physicians dropped to the same 1% mortality rate observed among mothers cared for by midwives. This demonstrated that handwashing was a very effective method for preventing disease transmission (**Figure 3.8**). Despite this great success, many discounted Semmelweis's work at the time, and physicians were slow to adopt the simple procedure of handwashing to prevent infections in their patients because it contradicted established norms for that time period.



**Figure 3.8** Ignaz Semmelweis (1818–1865) was a proponent of the importance of handwashing to prevent transfer of disease between patients by physicians.

Around the same time Semmelweis was promoting handwashing, in 1848, British physician John Snow conducted studies to track the source of cholera outbreaks in London. By tracing the outbreaks to two specific water sources, both of which were contaminated by sewage, Snow ultimately demonstrated that cholera bacteria were transmitted via drinking water. Snow's work is influential in that it represents the first known epidemiological study, and it resulted in the first known public health response to an epidemic. The work of both Semmelweis and Snow clearly refuted the prevailing miasma theory of the day, showing that disease is not only transmitted through the air but also through contaminated items.

Although the work of Semmelweis and Snow successfully showed the role of sanitation in preventing infectious disease, the cause of disease was not fully understood. The subsequent work of Louis Pasteur, Robert Koch, and Joseph Lister would further substantiate the germ theory of disease (**Figure 3.9**).

While studying the causes of beer and wine spoilage in 1856, Louis Pasteur discovered properties of fermentation by microorganisms. He had demonstrated with his swan-neck flask experiments (**Figure 3.4**) that airborne microbes, not spontaneous generation, were the cause of food spoilage, and he suggested that if microbes were responsible for food spoilage and fermentation, they could also be responsible for causing infection. This was the foundation for the germ theory of disease.

Meanwhile, British surgeon Joseph Lister was trying to determine the causes of postsurgical infections. Many physicians did not give credence to the idea that microbes on their hands, on their clothes, or in the air could infect patients' surgical wounds, despite the fact that 50% of surgical patients, on average, were dying of postsurgical infections.<sup>[15]</sup> Lister, however, was familiar with the work of Semmelweis and Pasteur; therefore, he insisted on

---

15. Alexander, J. Wesley. "The Contributions of Infection Control to a Century of Progress" *Annals of Surgery* 201:423-428, 1985.




handwashing and extreme cleanliness during surgery. In 1867, to further decrease the incidence of postsurgical wound infections, Lister began using carbolic acid (phenol) spray disinfectant/antiseptic during surgery. His extremely successful efforts to reduce postsurgical infection caused his techniques to become a standard medical practice.

A few years later, Robert Koch proposed a series of postulates (Koch's postulates) based on the idea that the cause of a specific disease could be attributed to a specific microbe. Using these postulates, Koch and his colleagues were able to definitively identify the causative pathogens of specific diseases, including anthrax, tuberculosis, and cholera. Koch's "one microbe, one disease" concept was the culmination of the 19th century's paradigm shift away from miasma theory and toward the germ theory of disease. Koch's postulates are discussed more thoroughly in **How Pathogens Cause Disease** (<http://cnx.org/content/m58867/latest/>).



(a)



(b)


**Figure 3.9** (a) Joseph Lister developed procedures for the proper care of surgical wounds and the sterilization of surgical equipment. (b) Robert Koch established a protocol to determine the cause of infectious disease. Both scientists contributed significantly to the acceptance of the germ theory of disease.

## Clinical Focus

### Part 2

After suffering a fever, congestion, cough, and increasing aches and pains for several days, Barbara suspects that she has a case of the flu. She decides to visit the health center at her university. The PA tells Barbara that her symptoms could be due to a range of diseases, such as influenza, bronchitis, pneumonia, or tuberculosis.


During her physical examination, the PA notes that Barbara's heart rate is slightly elevated. Using a pulse oximeter, a small device that clips on her finger, he finds that Barbara has hypoxemia—a lower-than-normal level of oxygen in the blood. Using a stethoscope, the PA listens for abnormal sounds made by Barbara's heart, lungs, and digestive system. As Barbara breathes, the PA hears a crackling sound and notes a slight shortness of breath. He collects a sputum sample, noting the greenish color of the mucus, and orders a chest radiograph, which shows a "shadow" in the left lung. All of these signs are suggestive of pneumonia, a condition in which the lungs fill with mucus (**Figure 3.10**).



**Figure 3.10** This is a chest radiograph typical of pneumonia. Because X-ray images are negative images, a “shadow” is seen as a white area within the lung that should otherwise be black. In this case, the left lung shows a shadow as a result of pockets in the lung that have become filled with fluid.

- What kinds of infectious agents are known to cause pneumonia?

Jump to the **next** Clinical Focus box. Go back to the **previous** Clinical Focus box.



**Figure 3.11**



## Check Your Understanding

- Compare and contrast the miasma theory of disease with the germ theory of disease.
- How did Joseph Lister's work contribute to the debate between the miasma theory and germ theory and how did this increase the success of medical procedures?

## 3.3 Unique Characteristics of Prokaryotic Cells

### Learning Objectives

- Explain the distinguishing characteristics of prokaryotic cells
- Describe common cell morphologies and cellular arrangements typical of prokaryotic cells and explain how cells maintain their morphology
- Describe internal and external structures of prokaryotic cells in terms of their physical structure, chemical structure, and function
- Compare the distinguishing characteristics of bacterial and archaeal cells

Cell theory states that the cell is the fundamental unit of life. However, cells vary significantly in size, shape, structure, and function. At the simplest level of construction, all cells possess a few fundamental components. These include **cytoplasm** (a gel-like substance composed of water and dissolved chemicals needed for growth), which is contained within a plasma membrane (also called a cell membrane or cytoplasmic membrane); one or more chromosomes, which contain the genetic blueprints of the cell; and **ribosomes**, organelles used for the production of proteins.


Beyond these basic components, cells can vary greatly between organisms, and even within the same multicellular organism. The two largest categories of cells—**prokaryotic cells** and **eukaryotic cells**—are defined by major differences in several cell structures. Prokaryotic cells lack a nucleus surrounded by a complex nuclear membrane and generally have a single, circular chromosome located in a nucleoid. Eukaryotic cells have a nucleus surrounded by a complex nuclear membrane that contains multiple, rod-shaped chromosomes.<sup>[16]</sup>

All plant cells and animal cells are eukaryotic. Some microorganisms are composed of prokaryotic cells, whereas others are composed of eukaryotic cells. Prokaryotic microorganisms are classified within the domains Archaea and Bacteria, whereas eukaryotic organisms are classified within the domain Eukarya.

The structures inside a cell are analogous to the organs inside a human body, with unique structures suited to specific functions. Some of the structures found in prokaryotic cells are similar to those found in some eukaryotic cells; others are unique to prokaryotes. Although there are some exceptions, eukaryotic cells tend to be larger than prokaryotic cells. The comparatively larger size of eukaryotic cells dictates the need to compartmentalize various chemical processes within different areas of the cell, using complex membrane-bound organelles. In contrast, prokaryotic cells generally lack membrane-bound organelles; however, they often contain inclusions that compartmentalize their cytoplasm. **Figure 3.12** illustrates structures typically associated with prokaryotic cells. These structures are described in more detail in the next section.

---

16. Y.-H.M. Chan, W.F. Marshall. "Scaling Properties of Cell and Organelle Size." *Organogenesis* 6 no. 2 (2010):88–96.



**Figure 3.12** A typical prokaryotic cell contains a cell membrane, chromosomal DNA that is concentrated in a nucleoid, ribosomes, and a cell wall. Some prokaryotic cells may also possess flagella, pili, fimbriae, and capsules.

### Common Cell Morphologies and Arrangements

Individual cells of a particular prokaryotic organism are typically similar in shape, or **cell morphology**. Although thousands of prokaryotic organisms have been identified, only a handful of cell morphologies are commonly seen microscopically. **Figure 3.13** names and illustrates cell morphologies commonly found in prokaryotic cells. In addition to cellular shape, prokaryotic cells of the same species may group together in certain distinctive arrangements depending on the plane of cell division. Some common arrangements are shown in **Figure 3.14**.




















Common Prokaryotic Cell Shapes			
Name	Description	Illustration	Image
Coccus (pl. cocci)	Round		
Bacillus (pl. bacilli)	Rod		
Vibrio (pl. vibrios)	Curved rod		
Coccobacillus (pl. coccobacilli)	Short rod		
Spirillum (pl. spirilla)	Spiral		
Spirochete (pl. spirochetes)	Long, loose, helical spiral		

Figure 3.13



Common Prokaryotic Cell Arrangements		
Name	Description	Illustration
Coccus (pl. cocci)	Single coccus	
Diplococcus (pl. diplococci)	Pair of two cocci	
Tetrad (pl. tetrads)	Grouping of four cells arranged in a square	
Streptococcus (pl. streptococci)	Chain of cocci	
Staphylococcus (pl. staphylococci)	Cluster of cocci	
Bacillus (pl. bacilli)	Single rod	
Streptobacillus (pl. streptobacilli)	Chain of rods	


**Figure 3.14**

In most prokaryotic cells, morphology is maintained by the **cell wall** in combination with cytoskeletal elements. The cell wall is a structure found in most prokaryotes and some eukaryotes; it envelops the cell membrane, protecting the cell from changes in **osmotic pressure** (Figure 3.15). Osmotic pressure occurs because of differences in the concentration of solutes on opposing sides of a semipermeable membrane. Water is able to pass through a semipermeable membrane, but solutes (dissolved molecules like salts, sugars, and other compounds) cannot. When the concentration of solutes is greater on one side of the membrane, water diffuses across the membrane from the side with the lower concentration (more water) to the side with the higher concentration (less water) until the concentrations on both sides become equal. This diffusion of water is called **osmosis**, and it can cause extreme osmotic pressure on a cell when its external environment changes.


The external environment of a cell can be described as an isotonic, hypertonic, or hypotonic medium. In an **isotonic medium**, the solute concentrations inside and outside the cell are approximately equal, so there is no net movement of water across the cell membrane. In a **hypertonic medium**, the solute concentration outside the cell exceeds that inside the cell, so water diffuses out of the cell and into the external medium. In a **hypotonic medium**, the solute concentration inside the cell exceeds that outside of the cell, so water will move by osmosis into the cell. This causes the cell to swell and potentially lyse, or burst.

The degree to which a particular cell is able to withstand changes in osmotic pressure is called tonicity. Cells that have a cell wall are better able to withstand subtle changes in osmotic pressure and maintain their shape. In hypertonic environments, cells that lack a cell wall can become dehydrated, causing **crenation**, or shriveling of the cell; the plasma membrane contracts and appears scalloped or notched (Figure 3.15). By contrast, cells that possess a cell wall undergo **plasmolysis** rather than crenation. In plasmolysis, the plasma membrane contracts and detaches from the cell wall, and there is a decrease in interior volume, but the cell wall remains intact, thus allowing the cell to maintain some shape and integrity for a period of time (Figure 3.16). Likewise, cells that lack a cell wall are more

prone to lysis in hypotonic environments. The presence of a cell wall allows the cell to maintain its shape and integrity for a longer time before lysing (Figure 3.16).



**Figure 3.15** In cells that lack a cell wall, changes in osmotic pressure can lead to crenation in hypertonic environments or cell lysis in hypotonic environments.



**Figure 3.16** In prokaryotic cells, the cell wall provides some protection against changes in osmotic pressure, allowing it to maintain its shape longer. The cell membrane is typically attached to the cell wall in an isotonic medium (left). In a hypertonic medium, the cell membrane detaches from the cell wall and contracts (plasmolysis) as water leaves the cell. In a hypotonic medium (right), the cell wall prevents the cell membrane from expanding to the point of bursting, although lysis will eventually occur if too much water is absorbed.




## Check Your Understanding

- Explain the difference between cell morphology and arrangement.
- What advantages do cell walls provide prokaryotic cells?

## The Nucleoid

All cellular life has a DNA genome organized into one or more chromosomes. Prokaryotic chromosomes are typically circular, haploid (unpaired), and not bound by a complex nuclear membrane. Prokaryotic DNA and DNA-associated proteins are concentrated within the **nucleoid** region of the cell (**Figure 3.17**). In general, prokaryotic DNA interacts with **nucleoid-associated proteins (NAPs)** that assist in the organization and packaging of the chromosome. In bacteria, NAPs function similar to histones, which are the DNA-organizing proteins found in eukaryotic cells. In archaea, the nucleoid is organized by either NAPs or histone-like DNA organizing proteins.




**Figure 3.17** The nucleoid region (the area enclosed by the green dashed line) is a condensed area of DNA found within prokaryotic cells. Because of the density of the area, it does not readily stain and appears lighter in color when viewed with a transmission electron microscope.

## Plasmids

Prokaryotic cells may also contain extrachromosomal DNA, or DNA that is not part of the chromosome. This extrachromosomal DNA is found in **plasmids**, which are small, circular, double-stranded DNA molecules. Cells that have plasmids often have hundreds of them within a single cell. Plasmids are more commonly found in bacteria; however, plasmids have been found in archaea and eukaryotic organisms. Plasmids often carry genes that confer advantageous traits such as antibiotic resistance; thus, they are important to the survival of the organism. We will discuss plasmids in more detail in **Mechanisms of Microbial Genetics** (<http://cnx.org/content/m58839/latest/>).

## Ribosomes

All cellular life synthesizes proteins, and organisms in all three domains of life possess ribosomes, structures responsible protein synthesis. However, ribosomes in each of the three domains are structurally different. Ribosomes, themselves, are constructed from proteins, along with ribosomal RNA (rRNA). Prokaryotic ribosomes are found in the cytoplasm. They are called **70S ribosomes** because they have a size of 70S (**Figure 3.18**), whereas eukaryotic cytoplasmic ribosomes have a size of 80S. (The S stands for Svedberg unit, a measure of sedimentation in an ultracentrifuge, which is based on size, shape, and surface qualities of the structure being analyzed). Although they are the same size, bacterial and archaeal ribosomes have different proteins and rRNA molecules, and the archaeal versions are more similar to their eukaryotic counterparts than to those found in bacteria.




**Figure 3.18** Prokaryotic ribosomes (70S) are composed of two subunits: the 30S (small subunit) and the 50S (large subunit), each of which are composed of protein and rRNA components.

## Inclusions

As single-celled organisms living in unstable environments, some prokaryotic cells have the ability to store excess nutrients within cytoplasmic structures called **inclusions**. Storing nutrients in a polymerized form is advantageous because it reduces the buildup of osmotic pressure that occurs as a cell accumulates solutes. Various types of inclusions store glycogen and starches, which contain carbon that cells can access for energy. **Volutin** granules, also called **metachromatic granules** because of their staining characteristics, are inclusions that store polymerized inorganic phosphate that can be used in metabolism and assist in the formation of biofilms. Microbes known to contain volutin granules include the archaea *Methanosarcina*, the bacterium *Corynebacterium diphtheriae*, and the unicellular eukaryotic alga *Chlamydomonas*. Sulfur granules, another type of inclusion, are found in sulfur bacteria of the genus *Thiobacillus*; these granules store elemental sulfur, which the bacteria use for metabolism.

Occasionally, certain types of inclusions are surrounded by a phospholipid monolayer embedded with protein. **Polyhydroxybutyrate (PHB)**, which can be produced by species of *Bacillus* and *Pseudomonas*, is an example of an inclusion that displays this type of monolayer structure. Industrially, PHB has also been used as a source of biodegradable polymers for bioplastics. Several different types of inclusions are shown in **Figure 3.19**.



**Figure 3.19** Prokaryotic cells may have various types of inclusions. (a) A transmission electron micrograph of polyhydroxybutyrate lipid droplets. (b) A light micrograph of volutin granules. (c) A phase-contrast micrograph of sulfur granules. (d) A transmission electron micrograph of magnetosomes. (e) A transmission electron micrograph of gas vacuoles.

Some prokaryotic cells have other types of inclusions that serve purposes other than nutrient storage. For example, some prokaryotic cells produce gas vacuoles, accumulations of small, protein-lined vesicles of gas. These gas vacuoles allow the prokaryotic cells that synthesize them to alter their buoyancy so that they can adjust their location in the water column. Magnetotactic bacteria, such as *Magnetospirillum magnetotacticum*, contain **magnetosomes**, which are inclusions of magnetic iron oxide or iron sulfide surrounded by a lipid layer. These allow cells to align along a magnetic field, aiding their movement (**Figure 3.19**). Cyanobacteria such as *Anabaena cylindrica* and bacteria such as *Halothiobacillus neapolitanus* produce **carboxysome** inclusions. Carboxysomes are composed of outer shells of thousands of protein subunits. Their interior is filled with ribulose-1,5-bisphosphate carboxylase/oxygenase (RuBisCO) and carbonic anhydrase. Both of these compounds are used for carbon metabolism. Some prokaryotic cells also possess carboxysomes that sequester functionally related enzymes in one location. These structures are considered proto-organelles because they compartmentalize important compounds or chemical reactions, much like many eukaryotic organelles.

## Endospores

Bacterial cells are generally observed as **vegetative cells**, but some genera of bacteria have the ability to form **endospores**, structures that essentially protect the bacterial genome in a dormant state when environmental conditions are unfavorable. Endospores (not to be confused with the reproductive spores formed by fungi) allow some bacterial cells to survive long periods without food or water, as well as exposure to chemicals, extreme temperatures, and even radiation. **Table 3.1** compares the characteristics of vegetative cells and endospores.





### Characteristics of Vegetative Cells versus Endospores


Vegetative Cells	Endospores
Sensitive to extreme temperatures and radiation	Resistant to extreme temperatures and radiation
Gram-positive	Do not absorb Gram stain, only special endospore stains (see <b>Staining Microscopic Specimens</b> )
Normal water content and enzymatic activity	Dehydrated; no metabolic activity
Capable of active growth and metabolism	Dormant; no growth or metabolic activity

**Table 3.1**

The process by which vegetative cells transform into endospores is called **sporulation**, and it generally begins when nutrients become depleted or environmental conditions become otherwise unfavorable (**Figure 3.20**). The process begins with the formation of a septum in the vegetative bacterial cell. The septum divides the cell asymmetrically, separating a DNA forespore from the mother cell. The forespore, which will form the core of the endospore, is essentially a copy of the cell's chromosomes, and is separated from the mother cell by a second membrane. A cortex gradually forms around the forespore by laying down layers of calcium and dipicolinic acid between membranes. A protein spore coat then forms around the cortex while the DNA of the mother cell disintegrates. Further maturation of the endospore occurs with the formation of an outermost exosporium. The endospore is released upon disintegration of the mother cell, completing sporulation.



(b)



(c)

**Figure 3.20** (a) Sporulation begins following asymmetric cell division. The forespore becomes surrounded by a double layer of membrane, a cortex, and a protein spore coat, before being released as a mature endospore upon disintegration of the mother cell. (b) An electron micrograph of a *Carboxydotherrnus hydrogenoformans* endospore. (c) These *Bacillus spp.* cells are undergoing sporulation. The endospores have been visualized using Malachite Green spore stain. (credit b: Jonathan Eisen)

Endospores of certain species have been shown to persist in a dormant state for extended periods of time, up to thousands of years.<sup>[17]</sup> However, when living conditions improve, endospores undergo **germination**, reentering a vegetative state. After germination, the cell becomes metabolically active again and is able to carry out all of its normal functions, including growth and cell division.

Not all bacteria have the ability to form endospores; however, there are a number of clinically significant endospore-forming gram-positive bacteria of the genera *Bacillus* and *Clostridium*. These include *B. anthracis*, the causative agent of anthrax, which produces endospores capable of survive for many decades<sup>[18]</sup>; *C. tetani* (causes tetanus); *C. difficile* (causes pseudomembranous colitis); *C. perfringens* (causes gas gangrene); and *C. botulinum* (causes botulism). Pathogens such as these are particularly difficult to combat because their endospores are so hard to kill. Special sterilization methods for endospore-forming bacteria are discussed in **Control of Microbial Growth** (<http://cnx.org/content/m58852/latest/>).



### Check Your Understanding

- What is an inclusion?
- What is the function of an endospore?


17. F. Rothfuss, M Bender, R Conrad. "Survival and Activity of Bacteria in a Deep, Aged Lake Sediment (Lake Constance)." *Microbial Ecology* 33 no. 1 (1997):69–77.

18. R. Sinclair et al. "Persistence of Category A Select Agents in the Environment." *Applied and Environmental Microbiology* 74 no. 3 (2008):555–563.

## Plasma Membrane

Structures that enclose the cytoplasm and internal structures of the cell are known collectively as the **cell envelope**. In prokaryotic cells, the structures of the cell envelope vary depending on the type of cell and organism. Most (but not all) prokaryotic cells have a cell wall, but the makeup of this cell wall varies. All cells (prokaryotic and eukaryotic) have a **plasma membrane** (also called **cytoplasmic membrane** or **cell membrane**) that exhibits selective permeability, allowing some molecules to enter or leave the cell while restricting the passage of others (**Figure 3.21**).

The structure of the plasma membrane is often described in terms of the **fluid mosaic model**, which refers to the ability of membrane components to move fluidly within the plane of the membrane, as well as the mosaic-like composition of the components, which include a diverse array of lipid and protein components. The plasma membrane structure of most bacterial and eukaryotic cell types is a bilayer composed mainly of phospholipids formed with ester linkages and proteins. These phospholipids and proteins have the ability to move laterally within the plane of the membranes as well as between the two phospholipid layers.



**Figure 3.21** The bacterial plasma membrane is a phospholipid bilayer with a variety of embedded proteins that perform various functions for the cell. Note the presence of glycoproteins and glycolipids, whose carbohydrate components extend out from the surface of the cell. The abundance and arrangement of these proteins and lipids can vary greatly between species.

Archaeal membranes are fundamentally different from bacterial and eukaryotic membranes in a few significant ways. First, archaeal membrane phospholipids are formed with ether linkages, in contrast to the ester linkages found in bacterial or eukaryotic cell membranes. Second, archaeal phospholipids have branched chains, whereas those of bacterial and eukaryotic cells are straight chained. Finally, although some archaeal membranes can be formed of bilayers like those found in bacteria and eukaryotes, other archaeal plasma membranes are lipid monolayers.

Proteins on the cell's surface are important for a variety of functions, including cell-to-cell communication, and sensing environmental conditions and pathogenic virulence factors. Membrane proteins and phospholipids may have carbohydrates (sugars) associated with them and are called glycoproteins or glycolipids, respectively. These glycoprotein and glycolipid complexes extend out from the surface of the cell, allowing the cell to interact with the external environment (**Figure 3.21**). Glycoproteins and glycolipids in the plasma membrane can vary considerably


in chemical composition among archaea, bacteria, and eukaryotes, allowing scientists to use them to characterize unique species.

Plasma membranes from different cells types also contain unique phospholipids, which contain fatty acids. As described in [Using Biochemistry to Identify Microorganisms \(http://cnx.org/content/m58817/latest/\)](http://cnx.org/content/m58817/latest/), phospholipid fatty acid analysis (PLFA) profiles can be used to identify unique types of cells based on differences in fatty acids. Archaea, bacteria, and eukaryotes each have a unique PFLA profile.

## Membrane Transport Mechanisms


One of the most important functions of the plasma membrane is to control the transport of molecules into and out of the cell. Internal conditions must be maintained within a certain range despite any changes in the external environment. The transport of substances across the plasma membrane allows cells to do so.

Cells use various modes of transport across the plasma membrane. For example, molecules moving from a higher concentration to a lower concentration with the concentration gradient are transported by simple diffusion, also known as passive transport (**Figure 3.22**). Some small molecules, like carbon dioxide, may cross the membrane bilayer directly by simple diffusion. However, charged molecules, as well as large molecules, need the help of carriers or channels in the membrane. These structures ferry molecules across the membrane, a process known as facilitated diffusion (**Figure 3.23**).




**Figure 3.22** Simple diffusion down a concentration gradient directly across the phospholipid bilayer.





**Figure 3.23** Facilitated diffusion down a concentration gradient through a membrane protein.

Active transport occurs when cells move molecules across their membrane *against* concentration gradients (**Figure 3.24**). A major difference between passive and active transport is that active transport requires adenosine triphosphate (ATP) or other forms of energy to move molecules “uphill.” Therefore, active transport structures are often called “pumps.”



**Figure 3.24** Active transport against a concentration gradient via a membrane pump that requires energy

Group translocation also transports substances into bacterial cells. In this case, as a molecule moves into a cell against its concentration gradient, it is chemically modified so that it does not require transport against an unfavorable concentration gradient. A common example of this is the bacterial phosphotransferase system, a series of carriers that phosphorylates (i.e., adds phosphate ions to) glucose or other sugars upon entry into cells. Since the phosphorylation of sugars is required during the early stages of sugar metabolism, the phosphotransferase system is considered to be an energy neutral system.

### Photosynthetic Membrane Structures

Some prokaryotic cells, namely cyanobacteria and photosynthetic bacteria, have membrane structures that enable them to perform photosynthesis. These structures consist of an infolding of the plasma membrane that encloses photosynthetic pigments such as green **chlorophylls** and bacteriochlorophylls. In cyanobacteria, these membrane structures are called thylakoids; in photosynthetic bacteria, they are called chromatophores, lamellae, or chlorosomes.




## Cell Wall

The primary function of the cell wall is to protect the cell from harsh conditions in the outside environment. When present, there are notable similarities and differences among the cell walls of archaea, bacteria, and eukaryotes.

The major component of bacterial cell walls is called **peptidoglycan** (or murein); it is only found in bacteria. Structurally, peptidoglycan resembles a layer of meshwork or fabric (**Figure 3.25**). Each layer is composed of long chains of alternating molecules of N-acetylglucosamine (NAG) and N-acetylmuramic acid (NAM). The structure of the long chains has significant two-dimensional tensile strength due to the formation of peptide bridges that connect NAG and NAM within each peptidoglycan layer. In gram-negative bacteria, tetrapeptide chains extending from each NAM unit are directly cross-linked, whereas in gram-positive bacteria, these tetrapeptide chains are linked by pentaglycine cross-bridges. Peptidoglycan subunits are made inside of the bacterial cell and then exported and assembled in layers, giving the cell its shape.

Since peptidoglycan is unique to bacteria, many antibiotic drugs are designed to interfere with peptidoglycan synthesis, weakening the cell wall and making bacterial cells more susceptible to the effects of osmotic pressure (see **Antibacterial Drugs** (<http://cnx.org/content/m58859/latest/>)). In addition, certain cells of the human immune system are able “recognize” bacterial pathogens by detecting peptidoglycan on the surface of a bacterial cell; these cells then engulf and destroy the bacterial cell, using enzymes such as lysozyme, which breaks down and digests the peptidoglycan in their cell walls (see **Pathogen Recognition and Phagocytosis** (<http://cnx.org/content/m58880/latest/>)).




**Figure 3.25** Peptidoglycan is composed of polymers of alternating NAM and NAG subunits, which are cross-linked by peptide bridges linking NAM subunits from various glycan chains. This provides the cell wall with tensile strength in two dimensions.


The Gram staining protocol (see **Staining Microscopic Specimens**) is used to differentiate two common types of cell wall structures (**Figure 3.26**). Gram-positive cells have a cell wall consisting of many layers of peptidoglycan totaling 30–100 nm in thickness. These peptidoglycan layers are commonly embedded with teichoic acids (TAs), carbohydrate chains that extend through and beyond the peptidoglycan layer.<sup>[19]</sup> TA is thought to stabilize peptidoglycan by increasing its rigidity. TA also plays a role in the ability of pathogenic gram-positive bacteria such as *Streptococcus* to bind to certain proteins on the surface of host cells, enhancing their ability to cause infection. In addition to peptidoglycan and TAs, a certain family of bacteria, the Mycobacteriaceae, have an additional external layer of waxy **mycolic acids** in their cell wall; as described in **Staining Microscopic Specimens**, these bacteria

19. T.J. Silhavy, D. Kahne, S. Walker. “The Bacterial Cell Envelope.” *Cold Spring Harbor Perspectives in Biology* 2 no. 5 (2010):a000414.

are referred to as acid-fast, since acid-fast stains must be used to penetrate the mycolic acid layer for purposes of microscopy (**Figure 3.27**).



**Figure 3.26** Bacteria contain two common cell wall structural types. Gram-positive cell walls are structurally simple, containing a thick layer of peptidoglycan with embedded teichoic acid external to the plasma membrane.<sup>[20]</sup> Gram-negative cell walls are structurally more complex, containing three layers: the inner membrane, a thin layer of peptidoglycan, and an outer membrane containing lipopolysaccharide.




**Figure 3.27** (a) Some gram-positive bacteria, including members of the Mycobacteriaceae, produce waxy mycolic acids found exterior to their structurally-distinct peptidoglycan. (b) The acid-fast staining protocol detects the presence of cell walls that are rich in mycolic acid. Acid-fast cells are stained red by carbolfuchsin.

Gram-negative cells have a much thinner layer of peptidoglycan (no more than about 4 nm thick<sup>[21]</sup>) than gram-positive cells, and the overall structure of their cell envelope is more complex. In gram-negative cells, a gel-like matrix occupies the **periplasmic space** between the cell wall and the plasma membrane, and there is a second lipid bilayer called the **outer membrane**, which is external to the peptidoglycan layer (**Figure 3.26**). This outer membrane is attached to the peptidoglycan by murein lipoprotein. The outer leaflet of the outer membrane contains the molecule **lipopolysaccharide (LPS)**, which functions as an endotoxin in infections involving gram-negative bacteria, contributing to symptoms such as fever, hemorrhaging, and septic shock. Each LPS molecule is composed

20. B. Zuber et al. "Granular Layer in the Periplasmic Space of Gram-Positive Bacteria and Fine Structures of *Enterococcus gallinarum* and *Streptococcus gordonii* Septa Revealed by Cryo-Electron Microscopy of Vitreous Sections." *Journal of Bacteriology* 188 no. 18 (2006):6652–6660

21. L. Gana, S. Chena, G.J. Jensen. "Molecular Organization of Gram-Negative Peptidoglycan." *Proceedings of the National Academy of Sciences of the United States of America* 105 no. 48 (2008):18953–18957.

of Lipid A, a core polysaccharide, and an O side chain that is composed of sugar-like molecules that comprise the external face of the LPS (**Figure 3.28**). The composition of the O side chain varies between different species and strains of bacteria. Parts of the O side chain called antigens can be detected using serological or immunological tests to identify specific pathogenic strains like *Escherichia coli* O157:H7, a deadly strain of bacteria that causes bloody diarrhea and kidney failure.



**Figure 3.28** The outer membrane of a gram-negative bacterial cell contains lipopolysaccharide (LPS), a toxin composed of Lipid A embedded in the outer membrane, a core polysaccharide, and the O side chain.

Archaeal cell wall structure differs from that of bacteria in several significant ways. First, archaeal cell walls do not contain peptidoglycan; instead, they contain a similar polymer called pseudopeptidoglycan (pseudomurein) in which NAM is replaced with a different subunit. Other archaea may have a layer of glycoproteins or polysaccharides that serves as the cell wall instead of pseudopeptidoglycan. Last, as is the case with some bacterial species, there are a few archaea that appear to lack cell walls entirely.


## Glycocalyx and S-Layers

Although most prokaryotic cells have cell walls, some may have additional cell envelope structures exterior to the cell wall, such as glycocalyxes and S-layers. A **glycocalyx** is a sugar coat, of which there are two important types: capsules and slime layers. A **capsule** is an organized layer located outside of the cell wall and usually composed of polysaccharides or proteins (**Figure 3.29**). A **slime layer** is a less tightly organized layer that is only loosely attached to the cell wall and can be more easily washed off. Slime layers may be composed of polysaccharides, glycoproteins, or glycolipids.

Glycocalyxes allows cells to adhere to surfaces, aiding in the formation of biofilms (colonies of microbes that form in layers on surfaces). In nature, most microbes live in mixed communities within biofilms, partly because the biofilm affords them some level of protection. Biofilms generally hold water like a sponge, preventing desiccation. They also protect cells from predation and hinder the action of antibiotics and disinfectants. All of these properties are



advantageous to the microbes living in a biofilm, but they present challenges in a clinical setting, where the goal is often to eliminate microbes.



**Figure 3.29** (a) Capsules are a type of glycocalyx composed of an organized layer of polysaccharides. (b) A capsule stain of *Pseudomonas aeruginosa*, a bacterial pathogen capable of causing many different types of infections in humans.

The ability to produce a capsule can contribute to a microbe's pathogenicity (ability to cause disease) because the capsule can make it more difficult for phagocytic cells (such as white blood cells) to engulf and kill the microorganism. *Streptococcus pneumoniae*, for example, produces a capsule that is well known to aid in this bacterium's pathogenicity. As explained in **Staining Microscopic specimens**, capsules are difficult to stain for microscopy, and are visualized by negative staining.

An **S-layer** is another type of cell envelope structure; it is composed of a mixture of structural proteins and glycoproteins. In bacteria, S-layers are found outside the cell wall, but in some archaea, the S-layer serves as the cell wall. The exact function of S-layers is not entirely understood, and they are difficult to study; but available evidence suggests that they may play a variety of functions in different prokaryotic cells, such as helping the cell withstand osmotic pressure and, for certain pathogens, interacting with the host immune system.

## Clinical Focus

### Part 3

After diagnosing Barbara with pneumonia, the PA writes her a prescription for amoxicillin, a commonly-prescribed type of penicillin derivative. More than a week later, despite taking the full course as directed, Barbara still feels weak and is not fully recovered, although she is still able to get through her daily activities. She returns to the health center for a follow-up visit.

Many types of bacteria, fungi, and viruses can cause pneumonia. Amoxicillin targets the peptidoglycan of bacterial cell walls. Since the amoxicillin has not resolved Barbara's symptoms, the PA concludes that the causative agent probably lacks peptidoglycan, meaning that the pathogen could be a virus, a fungus, or a bacterium that lacks peptidoglycan. Another possibility is that the pathogen is a bacterium containing peptidoglycan but has developed resistance to amoxicillin.

- How can the PA definitively identify the cause of Barbara's pneumonia?
- What form of treatment should the PA prescribe, given that the amoxicillin was ineffective?

Jump to the **next** Clinical Focus box. Go back to the **previous** Clinical Focus box.


## Filamentous Appendages

Many bacterial cells have protein appendages embedded within their cell envelopes that extend outward, allowing interaction with the environment. These appendages can attach to other surfaces, transfer DNA, or provide movement. Filamentous appendages include fimbriae, pili, and flagella.

### Fimbriae and Pili

Fimbriae and pili are structurally similar and, because differentiation between the two is problematic, these terms are often used interchangeably.<sup>[22] [23]</sup> The term **fimbriae** commonly refers to short bristle-like proteins projecting from the cell surface by the hundreds (**Figure 3.30**). Fimbriae enable a cell to attach to surfaces and to other cells. For pathogenic bacteria, adherence to host cells is important for colonization, infectivity, and virulence. Adherence to surfaces is also important in biofilm formation.

The term **pili** (singular: pilus) commonly refers to longer, less numerous protein appendages that aid in attachment to surfaces (**Figure 3.30**). A specific type of pilus, called the **F pilus** or **sex pilus**, is important in the transfer of DNA between bacterial cells, which occurs between members of the same generation when two cells physically transfer or exchange parts of their respective genomes (see **How Asexual Prokaryotes Achieve Genetic Diversity** (<http://cnx.org/content/m58845/latest/>)).



**Figure 3.30** Bacteria may produce two different types of protein appendages that aid in surface attachment. Fimbriae typically are more numerous and shorter, whereas pili (shown here) are longer and less numerous per cell.

### Micro Connections

#### Group A Strep

Before the structure and function of the various components of the bacterial cell envelope were well understood, scientists were already using cell envelope characteristics to classify bacteria. In 1933, Rebecca Lancefield proposed a method for serotyping various  $\beta$ -hemolytic strains of *Streptococcus* species using an agglutination assay, a technique using the clumping of bacteria to detect specific cell-surface antigens (**Figure 3.31**). In doing so, Lancefield discovered that one group of *S. pyogenes*, found in Group A, was associated with a variety of human diseases. She determined that various strains of Group A strep could be distinguished from each other based on variations in specific cell surface proteins that she named M proteins.

Today, more than 80 different strains of Group A strep have been identified based on M proteins. Various strains of Group A strep are associated with a wide variety of human infections, including streptococcal

22. J.A. Garnetta et al. "Structural Insights Into the Biogenesis and Biofilm Formation by the *Escherichia coli* Common Pilus." *Proceedings of the National Academy of Sciences of the United States of America* 109 no. 10 (2012):3950–3955.

23. T. Proft, E.N. Baker. "Pili in Gram-Negative and Gram-Positive Bacteria—Structure, Assembly and Their Role in Disease." *Cellular and Molecular Life Sciences* 66 (2009):613.




pharyngitis (strep throat), impetigo, toxic shock syndrome, scarlet fever, rheumatic fever, and necrotizing fasciitis. The M protein is an important virulence factor for Group A strep, helping these strains evade the immune system. Changes in M proteins appear to alter the infectivity of a particular strain of Group A strep.



**Figure 3.31** Rebecca Lancefield (1895–1981) was the first to develop a method for distinguishing various strains of  $\beta$ -hemolytic streptococci from each other based on their surface proteins.


## Flagella

**Flagella** are structures used by cells to move in aqueous environments. Bacterial flagella act like propellers. They are stiff spiral filaments composed of flagellin protein subunits that extend outward from the cell and spin in solution. The **basal body** is the motor for the flagellum and is embedded in the plasma membrane (**Figure 3.32**). A hook region connects the basal body to the filament. Gram-positive and gram-negative bacteria have different basal body configurations due to differences in cell wall structure.



**Figure 3.32** The basic structure of a bacterial flagellum consists of a basal body, hook, and filament. The basal body composition and arrangement differ between gram-positive and gram-negative bacteria.


Different types of motile bacteria exhibit different arrangements of flagella (**Figure 3.33**). A bacterium with a singular flagellum, typically located at one end of the cell (polar), is said to have a **monotrichous** flagellum. An example of a monotrichously flagellated bacterial pathogen is *Vibrio cholerae*, the gram-negative bacterium that causes cholera. Cells with **amphitrichous** flagella have a flagellum or tufts of flagella at each end. An example is *Spirillum minus*, the cause of spirillary (Asian) rat-bite fever or sodoku. Cells with **lophotrichous** flagella have a tuft at one end of the cell. The gram-negative bacillus *Pseudomonas aeruginosa*, an opportunistic pathogen known for causing many infections, including “swimmer’s ear” and burn wound infections, has lophotrichous flagella. Flagella that cover the entire surface of a bacterial cell are called **peritrichous** flagella. The gram-negative bacterium *E. coli* shows a peritrichous arrangement of flagella.




**Figure 3.33** Flagellated bacteria may exhibit multiple arrangements of their flagella. Common arrangements include monotrichous, amphitrichous, lophotrichous, or peritrichous.

Directional movement depends on the configuration of the flagella. Bacteria can move in response to a variety of environmental signals, including light (**phototaxis**), magnetic fields (**magnetotaxis**) using magnetosomes, and, most commonly, chemical gradients (**chemotaxis**). Purposeful movement toward a chemical attractant, like a food source, or away from a repellent, like a poisonous chemical, is achieved by increasing the length of **runs** and decreasing the length of **tumbles**. When running, flagella rotate in a counterclockwise direction, allowing the bacterial cell to move forward. In a peritrichous bacterium, the flagella are all bundled together in a very streamlined way (**Figure 3.34**), allowing for efficient movement. When tumbling, flagella are splayed out while rotating in a clockwise direction, creating a looping motion and preventing meaningful forward movement but reorienting the cell toward the direction of the attractant. When an attractant exists, runs and tumbles still occur; however, the length of runs is longer, while

the length of the tumbles is reduced, allowing overall movement toward the higher concentration of the attractant. When no chemical gradient exists, the lengths of runs and tumbles are more equal, and overall movement is more random (**Figure 3.35**).



**Figure 3.34** Bacteria achieve directional movement by changing the rotation of their flagella. In a cell with peritrichous flagella, the flagella bundle when they rotate in a counterclockwise direction, resulting in a run. However, when the flagella rotate in a clockwise direction, the flagella are no longer bundled, resulting in tumbles.



**Figure 3.35** Without a chemical gradient, flagellar rotation cycles between counterclockwise (run) and clockwise (tumble) with no overall directional movement. However, when a chemical gradient of an attractant exists, the length of runs is extended, while the length of tumbles is decreased. This leads to chemotaxis: an overall directional movement toward the higher concentration of the attractant.



### Check Your Understanding

- What is the peptidoglycan layer and how does it differ between gram-positive and gram-negative bacteria?
- Compare and contrast monotrichous, amphitrichous, lophotrichous, or peritrichous flagella.

## 3.4 Unique Characteristics of Eukaryotic Cells


---

### Learning Objectives

- Explain the distinguishing characteristics of eukaryotic cells
- Describe internal and external structures of prokaryotic cells in terms of their physical structure, chemical structure, and function
- Identify and describe structures and organelles unique to eukaryotic cells
- Compare and contrast similar structures found in prokaryotic and eukaryotic cells

Eukaryotic organisms include protozoans, algae, fungi, plants, and animals. Some eukaryotic cells are independent, single-celled microorganisms, whereas others are part of multicellular organisms. The cells of eukaryotic organisms have several distinguishing characteristics. Above all, eukaryotic cells are defined by the presence of a nucleus surrounded by a complex nuclear membrane. Also, eukaryotic cells are characterized by the presence of membrane-bound organelles in the cytoplasm. Organelles such as mitochondria, the endoplasmic reticulum (ER), Golgi apparatus, lysosomes, and peroxisomes are held in place by the **cytoskeleton**, an internal network that supports transport of intracellular components and helps maintain cell shape (**Figure 3.36**). The genome of eukaryotic cells is packaged in multiple, rod-shaped chromosomes as opposed to the single, circular-shaped chromosome that characterizes most prokaryotic cells. **Table 3.2** compares the characteristics of eukaryotic cell structures with those of bacteria and archaea.





**Figure 3.36** An illustration of a generalized, single-celled eukaryotic organism. Note that cells of eukaryotic organisms vary greatly in terms of structure and function, and a particular cell may not have all of the structures shown here.

### Summary of Cell Structures

Cell Structure	Prokaryotes		Eukaryotes
	Bacteria	Archaea	
Size	~0.5–1 $\mu\text{m}$	~0.5–1 $\mu\text{m}$	~5–20 $\mu\text{m}$
Surface area-to-volume ratio	High	High	Low
Nucleus	No	No	Yes

**Table 3.2**



### Summary of Cell Structures


Cell Structure	Prokaryotes		Eukaryotes
	Bacteria	Archaea	
Genome characteristics	<ul style="list-style-type: none"> <li>• Single chromosome</li> <li>• Circular</li> <li>• Haploid</li> <li>• Lacks histones</li> </ul>	<ul style="list-style-type: none"> <li>• Single chromosome</li> <li>• Circular</li> <li>• Haploid</li> <li>• Contains histones</li> </ul>	<ul style="list-style-type: none"> <li>• Multiple chromosomes</li> <li>• Linear</li> <li>• Haploid or diploid</li> <li>• Contains histones</li> </ul>
Cell division	Binary fission	Binary fission	Mitosis, meiosis
Membrane lipid composition	<ul style="list-style-type: none"> <li>• Ester-linked</li> <li>• Straight-chain fatty acids</li> <li>• Bilayer</li> </ul>	<ul style="list-style-type: none"> <li>• Ether-linked</li> <li>• Branched isoprenoids</li> <li>• Bilayer or monolayer</li> </ul>	<ul style="list-style-type: none"> <li>• Ester-linked</li> <li>• Straight-chain fatty acids</li> <li>• Sterols</li> <li>• Bilayer</li> </ul>
Cell wall composition	<ul style="list-style-type: none"> <li>• Peptidoglycan, or</li> <li>• None</li> </ul>	<ul style="list-style-type: none"> <li>• Pseudopeptidoglycan, or</li> <li>• Glycopeptide, or</li> <li>• Polysaccharide, or</li> <li>• Protein (S-layer), or</li> <li>• None</li> </ul>	<ul style="list-style-type: none"> <li>• Cellulose (plants, some algae)</li> <li>• Chitin (molluscs, insects, crustaceans, and fungi)</li> <li>• Silica (some algae)</li> <li>• Most others lack cell walls</li> </ul>
Motility structures	Rigid spiral flagella composed of flagellin	Rigid spiral flagella composed of archaeal flagellins	Flexible flagella and cilia composed of microtubules
Membrane-bound organelles	No	No	Yes
Endomembrane system	No	No	Yes (ER, Golgi, lysosomes)
Ribosomes	70S	70S	<ul style="list-style-type: none"> <li>• 80S in cytoplasm and rough ER</li> <li>• 70S in mitochondria, chloroplasts</li> </ul>

**Table 3.2**

## Cell Morphologies

Eukaryotic cells display a wide variety of different cell morphologies. Possible shapes include spheroid, ovoid, cuboidal, cylindrical, flat, lenticular, fusiform, discoidal, crescent, ring stellate, and polygonal (**Figure 3.37**). Some eukaryotic cells are irregular in shape, and some are capable of changing shape. The shape of a particular type of eukaryotic cell may be influenced by factors such as its primary function, the organization of its cytoskeleton, the

viscosity of its cytoplasm, the rigidity of its cell membrane or cell wall (if it has one), and the physical pressure exerted on it by the surrounding environment and/or adjoining cells.



**Figure 3.37** Eukaryotic cells come in a variety of cell shapes. (a) Spheroid *Chromulina* alga. (b) Fusiform shaped *Trypanosoma*. (c) Bell-shaped *Vorticella*. (d) Ovoid *Paramecium*. (e) Ring-shaped *Plasmodium ovale*.




### Check Your Understanding

- Identify two differences between eukaryotic and prokaryotic cells.

## Nucleus


Unlike prokaryotic cells, in which DNA is loosely contained in the nucleoid region, eukaryotic cells possess a **nucleus**, which is surrounded by a complex nuclear membrane that houses the DNA genome (**Figure 3.38**). By containing the cell's DNA, the nucleus ultimately controls all activities of the cell and also serves an essential role in reproduction and heredity. Eukaryotic cells typically have their DNA organized into multiple linear chromosomes. The DNA within the nucleus is highly organized and condensed to fit inside the nucleus, which is accomplished by wrapping the DNA around proteins called histones.



**Figure 3.38** Eukaryotic cells have a well-defined nucleus. The nucleus of this mammalian lung cell is the large, dark, oval-shaped structure in the lower right.

Although most eukaryotic cells have only one nucleus, exceptions exist. For example, protozoans of the genus *Paramecium* typically have two complete nuclei: a small nucleus that is used for reproduction (micronucleus) and a large nucleus that directs cellular metabolism (macronucleus). Additionally, some fungi transiently form cells with two nuclei, called heterokaryotic cells, during sexual reproduction. Cells whose nuclei divide, but whose cytoplasm does not, are called **coenocytes**.


The nucleus is bound by a complex **nuclear membrane**, often called the **nuclear envelope**, that consists of two distinct lipid bilayers that are contiguous with each other (**Figure 3.39**). Despite these connections between the inner and outer membranes, each membrane contains unique lipids and proteins on its inner and outer surfaces. The nuclear envelope contains nuclear pores, which are large, rosette-shaped protein complexes that control the movement of materials into and out of the nucleus. The overall shape of the nucleus is determined by the **nuclear lamina**, a meshwork of intermediate filaments found just inside the nuclear envelope membranes. Outside the nucleus, additional intermediate filaments form a looser mesh and serve to anchor the nucleus in position within the cell.



**Figure 3.39** In this fluorescent microscope image, all the intermediate filaments have been stained with a bright green fluorescent stain. The nuclear lamina is the intense bright green ring around the faint red nuclei.

### Nucleolus

The **nucleolus** is a dense region within the nucleus where ribosomal RNA (rRNA) biosynthesis occurs. In addition, the nucleolus is also the site where assembly of ribosomes begins. Preribosomal complexes are assembled from rRNA and proteins in the nucleolus; they are then transported out to the cytoplasm, where ribosome assembly is completed (**Figure 3.40**).



**Figure 3.40** (a) The nucleolus is the dark, dense area within the nucleus. It is the site of rRNA synthesis and preribosomal assembly. (b) Electron micrograph showing the nucleolus.

## Ribosomes

Ribosomes found in eukaryotic organelles such as mitochondria or chloroplasts have 70S ribosomes—the same size as prokaryotic ribosomes. However, nonorganelle-associated ribosomes in eukaryotic cells are **80S ribosomes**, composed of a 40S small subunit and a 60S large subunit. In terms of size and composition, this makes them distinct from the ribosomes of prokaryotic cells.

The two types of nonorganelle-associated eukaryotic ribosomes are defined by their location in the cell: **free ribosomes** and **membrane-bound ribosomes**. Free ribosomes are found in the cytoplasm and serve to synthesize water-soluble proteins; membrane-bound ribosomes are found attached to the rough endoplasmic reticulum and make proteins for insertion into the cell membrane or proteins destined for export from the cell.


The differences between eukaryotic and prokaryotic ribosomes are clinically relevant because certain antibiotic drugs are designed to target one or the other. For example, cycloheximide targets eukaryotic action, whereas chloramphenicol targets prokaryotic ribosomes.<sup>[24]</sup> Since human cells are eukaryotic, they generally are not harmed by antibiotics that destroy the prokaryotic ribosomes in bacteria. However, sometimes negative side effects may occur because mitochondria in human cells contain prokaryotic ribosomes.

## The Endomembrane System

The **endomembrane system**, unique to eukaryotic cells, is a series of membranous tubules, sacs, and flattened disks that synthesize many cell components and move materials around within the cell (**Figure 3.41**). Because of their larger cell size, eukaryotic cells require this system to transport materials that cannot be dispersed by diffusion alone. The endomembrane system comprises several organelles and connections between them, including the endoplasmic reticulum, Golgi apparatus, lysosomes, and vesicles.

24. A.E. Barnhill, M.T. Brewer, S.A. Carlson. “Adverse Effects of Antimicrobials via Predictable or Idiosyncratic Inhibition of Host Mitochondrial Components.” *Antimicrobial Agents and Chemotherapy* 56 no. 8 (2012):4046–4051.






**Figure 3.41** The endomembrane system is composed of a series of membranous intracellular structures that facilitate movement of materials throughout the cell and to the cell membrane.

### Endoplasmic Reticulum

The **endoplasmic reticulum (ER)** is an interconnected array of tubules and **cisternae** (flattened sacs) with a single lipid bilayer (**Figure 3.42**). The spaces inside of the cisternae are called **lumen** of the ER.

There are two types of ER, **rough endoplasmic reticulum (RER)** and **smooth endoplasmic reticulum (SER)**. These two different types of ER are sites for the synthesis of distinctly different types of molecules. RER is studded with ribosomes bound on the cytoplasmic side of the membrane. These ribosomes make proteins destined for the plasma membrane (**Figure 3.39**, **Figure 3.40**, and **Figure 3.42**). Following synthesis, these proteins are inserted into the membrane of the RER. Small sacs of the RER containing these newly synthesized proteins then bud off as **transport vesicles** and move either to the Golgi apparatus for further processing, directly to the plasma membrane, to the membrane of another organelle, or out of the cell (**Figure 3.41**). Transport vesicles are single-lipid, bilayer, membranous spheres with hollow interiors that carry molecules. SER does not have ribosomes and, therefore, appears “smooth.” It is involved in biosynthesis of lipids, carbohydrate metabolism, and detoxification of toxic compounds within the cell.



**Figure 3.42** The rough endoplasmic reticulum is studded with ribosomes for the synthesis of membrane proteins (which give it its rough appearance).




## Golgi Apparatus

The Golgi apparatus was discovered within the endomembrane system in 1898 by Italian scientist Camillo Golgi (1843–1926), who developed a novel staining technique that showed stacked membrane structures within the cells of *Plasmodium*, the causative agent of malaria. The **Golgi apparatus** is composed of a series of membranous disks called dictyosomes, each having a single lipid bilayer, that are stacked together (**Figure 3.43**).

Enzymes in the Golgi apparatus modify lipids and proteins transported from the ER to the Golgi, often adding carbohydrate components to them, producing glycolipids, glycoproteins, or proteoglycans. Glycolipids and glycoproteins are often inserted into the plasma membrane and are important for signal recognition by other cells or infectious particles. Different types of cells can be distinguished from one another by the structure and arrangement of the glycolipids and glycoproteins contained in their plasma membranes. These glycolipids and glycoproteins commonly also serve as cell surface receptors (**Figure 3.54**).

Transport vesicles leaving the ER fuse with a Golgi apparatus on its receiving, or *cis*, face. The proteins are processed within the Golgi apparatus, and then additional transport vesicles containing the modified proteins and lipids pinch off from the Golgi apparatus on its outgoing, or *trans*, face. These outgoing vesicles move to and fuse with the plasma membrane or the membrane of other organelles.

Exocytosis is the process by which **secretory vesicles** (spherical membranous sacs) release their contents to the cell's exterior (**Figure 3.41**). All cells have constitutive secretory pathways in which secretory vesicles transport soluble proteins that are released from the cell continually (constitutively). Certain specialized cells also have regulated secretory pathways, which are used to store soluble proteins in secretory vesicles. Regulated secretion involves substances that are only released in response to certain events or signals. For example, certain cells of the human immune system (e.g., mast cells) secrete histamine in response to the presence of foreign objects or pathogens in the body. Histamine is a compound that triggers various mechanisms used by the immune system to eliminate pathogens.



**Figure 3.43** A transmission electron micrograph (left) of a Golgi apparatus in a white blood cell. The illustration (right) shows the cup-shaped, stacked disks and several transport vesicles. The Golgi apparatus modifies lipids and proteins, producing glycolipids and glycoproteins, respectively, which are commonly inserted into the plasma membrane.

## Lysosomes

In the 1960s, Belgian scientist Christian de Duve (1917–2013) discovered **lysosomes**, membrane-bound organelles of the endomembrane system that contain digestive enzymes ([Figure 03\\_04\\_Lysosome](#)). Certain types of eukaryotic cells use lysosomes to break down various particles, such as food, damaged organelles or cellular debris,

microorganisms, or immune complexes. Compartmentalization of the digestive enzymes within the lysosome allows the cell to efficiently digest matter without harming the cytoplasmic components of the cell.



## Check Your Understanding


- Name the components of the endomembrane system and describe the function of each component.

## Peroxisomes

Christian de Duve is also credited with the discovery of **peroxisomes**, membrane-bound organelles that are not part of the endomembrane system (**Figure 3.44**). Peroxisomes form independently in the cytoplasm from the synthesis of peroxin proteins by free ribosomes and the incorporation of these peroxin proteins into existing peroxisomes. Growing peroxisomes then divide by a process similar to binary fission.

Peroxisomes were first named for their ability to produce hydrogen peroxide, a highly reactive molecule that helps to break down molecules such as uric acid, amino acids, and fatty acids. Peroxisomes also possess the enzyme catalase, which can degrade hydrogen peroxide. Along with the SER, peroxisomes also play a role in lipid biosynthesis. Like lysosomes, the compartmentalization of these degradative molecules within an organelle helps protect the cytoplasmic contents from unwanted damage.


The peroxisomes of certain organisms are specialized to meet their particular functional needs. For example, glyoxysomes are modified peroxisomes of yeasts and plant cells that perform several metabolic functions, including the production of sugar molecules. Similarly, glycosomes are modified peroxisomes made by certain trypanosomes, the pathogenic protozoans that cause Chagas disease and African sleeping sickness.



**Figure 3.44** A transmission electron micrograph (left) of a cell containing a peroxisome. The illustration (right) shows the location of peroxisomes in a cell. These eukaryotic structures play a role in lipid biosynthesis and breaking down various molecules. They may also have other specialized functions depending on the cell type.

## Cytoskeleton

Eukaryotic cells have an internal cytoskeleton made of **microfilaments**, **intermediate filaments**, and **microtubules**. This matrix of fibers and tubes provides structural support as well as a network over which materials can be transported within the cell and on which organelles can be anchored (**Figure 3.45**). For example, the process of exocytosis involves the movement of a vesicle via the cytoskeletal network to the plasma membrane, where it can release its contents.




**Figure 3.45** The cytoskeleton is a network of microfilaments, intermediate filaments, and microtubules found throughout the cytoplasm of a eukaryotic cell. In these fluorescently labeled animal cells, the microtubules are green, the actin microfilaments are red, the nucleus is blue, and keratin (a type of intermediate filament) is yellow.

Microfilaments are composed of two intertwined strands of actin, each composed of **actin** monomers forming filamentous cables 6 nm in diameter<sup>[25]</sup> (**Figure 3.46**). The actin filaments work together with motor proteins, like myosin, to effect muscle contraction in animals or the amoeboid movement of some eukaryotic microbes. In amoeboid organisms, actin can be found in two forms: a stiffer, polymerized, gel form and a more fluid, unpolymerized soluble form. Actin in the gel form creates stability in the ectoplasm, the gel-like area of cytoplasm just inside the plasma membrane of amoeboid protozoans.

Temporary extensions of the cytoplasmic membrane called **pseudopodia** (meaning “false feet”) are produced through the forward flow of soluble actin filaments into the pseudopodia, followed by the gel-sol cycling of the actin filaments, resulting in cell motility. Once the cytoplasm extends outward, forming a pseudopodium, the remaining cytoplasm flows up to join the leading edge, thereby creating forward locomotion. Beyond amoeboid movement, microfilaments are also involved in a variety of other processes in eukaryotic cells, including cytoplasmic streaming (the movement or circulation of cytoplasm within the cell), cleavage furrow formation during cell division, and muscle movement in animals (**Figure 3.46**). These functions are the result of the dynamic nature of microfilaments, which can polymerize and depolymerize relatively easily in response to cellular signals, and their interactions with molecular motors in different types of eukaryotic cells.

25. Fuchs E, Cleveland DW. “A Structural Scaffolding of Intermediate Filaments in Health and Disease.” *Science* 279 no. 5350 (1998):514–519.






**Figure 3.46** (a) A microfilament is composed of a pair of actin filaments. (b) Each actin filament is a string of polymerized actin monomers. (c) The dynamic nature of actin, due to its polymerization and depolymerization and its association with myosin, allows microfilaments to be involved in a variety of cellular processes, including amoeboid movement, cytoplasmic streaming, contractile ring formation during cell division, and muscle contraction in animals.


Intermediate filaments are a diverse group of cytoskeletal filaments that act as cables within the cell. They are termed “intermediate” because their 10-nm diameter is thicker than that of actin but thinner than that of microtubules.<sup>[26]</sup> They are composed of several strands of polymerized subunits that, in turn, are made up of a wide variety of monomers. Intermediate filaments tend to be more permanent in the cell and maintain the position of the nucleus. They also form the nuclear lamina (lining or layer) just inside the nuclear envelope. Additionally, intermediate filaments play a role in anchoring cells together in animal tissues. The intermediate filament protein desmin is found in desmosomes, the protein structures that join muscle cells together and help them resist external physical forces. The intermediate filament protein keratin is a structural protein found in hair, skin, and nails (**Figure 3.47**).

26. E. Fuchs, D.W. Cleveland. “A Structural Scaffolding of Intermediate Filaments in Health and Disease.” *Science* 279 no. 5350 (1998):514–519.



**Figure 3.47** (a) Intermediate filaments are composed of multiple strands of polymerized subunits. They are more permanent than other cytoskeletal structures and serve a variety of functions. (b) Intermediate filaments form much of the nuclear lamina. (c) Intermediate filaments form the desmosomes between cells in some animal tissues.

Microtubules are a third type of cytoskeletal fiber composed of tubulin dimers ( $\alpha$  tubulin and  $\beta$  tubulin). These form hollow tubes 23 nm in diameter that are used as girders within the cytoskeleton (**Figure 3.48**).<sup>[27]</sup> Like microfilaments, microtubules are dynamic and have the ability to rapidly assemble and disassemble. Microtubules also work with motor proteins (such as dynein and kinesin) to move organelles and vesicles around within the cytoplasm. Additionally, microtubules are the main components of eukaryotic flagella and cilia, composing both the filament and the basal body components (**Figure 3.55**).




**Figure 3.48** (a) Microtubules are hollow structures composed of polymerized tubulin dimers. (b) They are involved in several cellular processes, including the movement of organelles throughout the cytoplasm. Motor proteins carry organelles along microtubule tracks that crisscross the entire cell.

In addition, microtubules are involved in cell division, forming the mitotic spindle that serves to separate chromosomes during mitosis and meiosis. The mitotic spindle is produced by two **centrosomes**, which are essentially microtubule-organizing centers, at opposite ends of the cell. Each centrosome is composed of a pair of **centrioles** positioned at right angles to each other, and each centriole is an array of nine parallel microtubules arranged in triplets (**Figure 3.49**).

27. E. Fuchs, D.W. Cleveland. "A Structural Scaffolding of Intermediate Filaments in Health and Disease." *Science* 279 no. 5350 (1998):514–519.





**Figure 3.49** (a) A centrosome is composed of two centrioles positioned at right angles to each other. Each centriole is composed of nine triplets of microtubules held together by accessory proteins. (b) In animal cells, the centrosomes (arrows) serve as microtubule-organizing centers of the mitotic spindle during mitosis.




### Check Your Understanding

- Compare and contrast the three types of cytoskeletal structures described in this section.

## Mitochondria

The large, complex organelles in which aerobic cellular respiration occurs in eukaryotic cells are called **mitochondria** (**Figure 3.50**). The term “mitochondrion” was first coined by German microbiologist Carl Benda in 1898 and was later connected with the process of respiration by Otto Warburg in 1913. Scientists during the 1960s discovered that mitochondria have their own genome and 70S ribosomes. The mitochondrial genome was found to be bacterial, when it was sequenced in 1976. These findings ultimately supported the endosymbiotic theory proposed by Lynn Margulis, which states that mitochondria originally arose through an endosymbiotic event in which a bacterium capable of aerobic cellular respiration was taken up by phagocytosis into a host cell and remained as a viable intracellular component.


Each mitochondrion has two lipid membranes. The outer membrane is a remnant of the original host cell’s membrane structures. The inner membrane was derived from the bacterial plasma membrane. The electron transport chain for aerobic respiration uses integral proteins embedded in the inner membrane. The **mitochondrial matrix**, corresponding to the location of the original bacterium’s cytoplasm, is the current location of many metabolic enzymes. It also contains mitochondrial DNA and 70S ribosomes. Invaginations of the inner membrane, called cristae, evolved to increase surface area for the location of biochemical reactions. The folding patterns of the cristae differ among various types of eukaryotic cells and are used to distinguish different eukaryotic organisms from each other.



**Figure 3.50** Each mitochondrion is surrounded by two membranes, the inner of which is extensively folded into cristae and is the site of the intermembrane space. The mitochondrial matrix contains the mitochondrial DNA, ribosomes, and metabolic enzymes. The transmission electron micrograph of a mitochondrion, on the right, shows both membranes, including cristae and the mitochondrial matrix.

## Chloroplasts

Plant cells and algal cells contain **chloroplasts**, the organelles in which photosynthesis occurs (**Figure 3.51**). All chloroplasts have at least three membrane systems: the outer membrane, the inner membrane, and the thylakoid membrane system. Inside the outer and inner membranes is the chloroplast **stroma**, a gel-like fluid that makes up much of a chloroplast's volume, and in which the **thylakoid** system floats. The thylakoid system is a highly dynamic collection of folded membrane sacs. It is where the green photosynthetic pigment chlorophyll is found and the light reactions of photosynthesis occur. In most plant chloroplasts, the thylakoids are arranged in stacks called grana (singular: granum), whereas in some algal chloroplasts, the thylakoids are free floating.



**Figure 3.51** Photosynthesis takes place in chloroplasts, which have an outer membrane and an inner membrane. Stacks of thylakoids called grana form a third membrane layer.

Other organelles similar to mitochondria have arisen in other types of eukaryotes, but their roles differ. Hydrogenosomes are found in some anaerobic eukaryotes and serve as the location of anaerobic hydrogen production. Hydrogenosomes typically lack their own DNA and ribosomes. Kinetoplasts are a variation of the mitochondria found in some eukaryotic pathogens. In these organisms, each cell has a single, long, branched mitochondrion in which kinetoplast DNA, organized as multiple circular pieces of DNA, is found concentrated at one pole of the cell.

## Micro Connections

### Mitochondria-Related Organelles in Protozoan Parasites

Many protozoans, including several protozoan parasites that cause infections in humans, can be identified by their unusual appearance. Distinguishing features may include complex cell morphologies, the presence of unique organelles, or the absence of common organelles. The protozoan parasites *Giardia lamblia* and *Trichomonas vaginalis* are two examples.


*G. lamblia*, a frequent cause of diarrhea in humans and many other animals, is an anaerobic parasite that possesses two nuclei and several flagella. Its Golgi apparatus and endoplasmic reticulum are greatly reduced, and it lacks mitochondria completely. However, it does have organelles known as mitosomes, double-membrane-bound organelles that appear to be severely reduced mitochondria. This has led scientists to believe that *G. lamblia*'s ancestors once possessed mitochondria that evolved to become mitosomes. *T. vaginalis*, which causes the sexually transmitted infection vaginitis, is another protozoan parasite that lacks conventional mitochondria. Instead, it possesses hydrogenosomes, mitochondrial-related, double-membrane-bound organelles that produce molecular hydrogen used in cellular metabolism. Scientists believe that hydrogenosomes, like mitosomes, also evolved from mitochondria.<sup>[28]</sup>

## Plasma Membrane

The plasma membrane of eukaryotic cells is similar in structure to the prokaryotic plasma membrane in that it is composed mainly of phospholipids forming a bilayer with embedded peripheral and integral proteins (**Figure 3.52**). These membrane components move within the plane of the membrane according to the fluid mosaic model. However, unlike the prokaryotic membrane, eukaryotic membranes contain sterols, including cholesterol, that alter membrane fluidity. Additionally, many eukaryotic cells contain some specialized lipids, including sphingolipids, which are thought to play a role in maintaining membrane stability as well as being involved in signal transduction pathways and cell-to-cell communication.

---

28. N. Yarlett, J.H.P. Hackstein. "Hydrogenosomes: One Organelle, Multiple Origins." *BioScience* 55 no. 8 (2005):657–658.




**Figure 3.52** The eukaryotic plasma membrane is composed of a lipid bilayer with many embedded or associated proteins. It contains cholesterol for the maintenance of membrane, as well as glycoproteins and glycolipids that are important in the recognition other cells or pathogens.

### Membrane Transport Mechanisms

The processes of simple diffusion, facilitated diffusion, and active transport are used in both eukaryotic and prokaryotic cells. However, eukaryotic cells also have the unique ability to perform various types of **endocytosis**, the uptake of matter through plasma membrane invagination and vacuole/vesicle formation (**Figure 3.53**). A type of endocytosis involving the engulfment of large particles through membrane invagination is called **phagocytosis**, which means “cell eating.” In phagocytosis, particles (or other cells) are enclosed in a pocket within the membrane, which then pinches off from the membrane to form a vacuole that completely surrounds the particle. Another type of endocytosis is called **pinocytosis**, which means “cell drinking.” In pinocytosis, small, dissolved materials and liquids are taken into the cell through small vesicles. Saprophytic fungi, for example, obtain their nutrients from dead and decaying matter largely through pinocytosis.

**Receptor-mediated endocytosis** is a type of endocytosis that is initiated by specific molecules called ligands when they bind to cell surface receptors on the membrane. Receptor-mediated endocytosis is the mechanism that peptide and amine-derived hormones use to enter cells and is also used by various viruses and bacteria for entry into host cells.





**Figure 3.53** Three variations of endocytosis are shown. (a) In phagocytosis, the cell membrane surrounds the particle and pinches off to form an intracellular vacuole. (b) In pinocytosis, the cell membrane surrounds a small volume of fluid and pinches off, forming a vesicle. (c) In receptor-mediated endocytosis, the uptake of substances is targeted to a specific substance (a ligand) that binds at the receptor on the external cell membrane.

The process by which secretory vesicles release their contents to the cell's exterior is called **exocytosis** (Figure 3.41). Vesicles move toward the plasma membrane and then meld with the membrane, ejecting their contents out of the cell. Exocytosis is used by cells to remove waste products and may also be used to release chemical signals that can be taken up by other cells.

## Cell Wall

In addition to a plasma membrane, some eukaryotic cells have a cell wall. Cells of fungi, algae, plants, and even some protists have cell walls. Depending upon the type of eukaryotic cell, cell walls can be made of a wide range of materials, including cellulose (fungi and plants); biogenic silica, calcium carbonate, agar, and carrageenan (protists and algae); or chitin (fungi). In general, all cell walls provide structural stability for the cell and protection from environmental stresses such as desiccation, changes in osmotic pressure, and traumatic injury.<sup>[29]</sup>

## Extracellular Matrix


Cells of animals and some protozoans do not have cell walls to help maintain shape and provide structural stability. Instead, these types of eukaryotic cells produce an **extracellular matrix** for this purpose. They secrete a sticky mass of carbohydrates and proteins into the spaces between adjacent cells (Figure 3.54). Some protein components assemble into a basement membrane to which the remaining extracellular matrix components adhere. Proteoglycans typically form the bulky mass of the extracellular matrix while fibrous proteins, like collagen, provide strength. Both proteoglycans and collagen are attached to fibronectin proteins, which, in turn, are attached to integrin proteins. These integrin proteins interact with transmembrane proteins in the plasma membranes of eukaryotic cells that lack cell walls.

In animal cells, the extracellular matrix allows cells within tissues to withstand external stresses and transmits signals from the outside of the cell to the inside. The amount of extracellular matrix is quite extensive in various types of

29. M. Dudzick. "Protists." OpenStax CNX. November 27, 2013. <http://cnx.org/contents/f7048bb6-e462-459b-805c-ef291cf7049c@1>



connective tissues, and variations in the extracellular matrix can give different types of tissues their distinct properties. In addition, a host cell's extracellular matrix is often the site where microbial pathogens attach themselves to establish infection. For example, *Streptococcus pyogenes*, the bacterium that causes strep throat and various other infections, binds to fibronectin in the extracellular matrix of the cells lining the oropharynx (upper region of the throat).




**Figure 3.54** The extracellular matrix is composed of protein and carbohydrate components. It protects cells from physical stresses and transmits signals arriving at the outside edges of the tissue to cells deeper within the tissue.

## Flagella and Cilia

Some eukaryotic cells use **flagella** for locomotion; however, eukaryotic flagella are structurally distinct from those found in prokaryotic cells. Whereas the prokaryotic flagellum is a stiff, rotating structure, a eukaryotic flagellum is more like a flexible whip composed of nine parallel pairs of microtubules surrounding a central pair of microtubules. This arrangement is referred to as a 9+2 array (**Figure 3.55**). The parallel microtubules use **dynein** motor proteins to move relative to each other, causing the flagellum to bend.

**Cilia** (singular: **cilium**) are a similar external structure found in some eukaryotic cells. Unique to eukaryotes, cilia are shorter than flagella and often cover the entire surface of a cell; however, they are structurally similar to flagella (a 9+2 array of microtubules) and use the same mechanism for movement. A structure called a **basal body** is found at the base of each cilium and flagellum. The basal body, which attaches the cilium or flagellum to the cell, is composed of an array of triplet microtubules similar to that of a centriole but embedded in the plasma membrane. Because of their shorter length, cilia use a rapid, flexible, waving motion. In addition to motility, cilia may have other functions such as sweeping particles past or into cells. For example, ciliated protozoans use the sweeping of cilia to move food

particles into their mouthparts, and ciliated cells in the mammalian respiratory tract beat in synchrony to sweep mucus and debris up and out of the lungs (**Figure 3.55**).



**Figure 3.55** (a) Eukaryotic flagella and cilia are composed of a 9+2 array of microtubules, as seen in this transmission electron micrograph cross-section. (b) The sliding of these microtubules relative to each other causes a flagellum to bend. (c) An illustration of *Trichomonas vaginalis*, a flagellated protozoan parasite that causes vaginitis. (d) Many protozoans, like this *Paramecium*, have numerous cilia that aid in locomotion as well as in feeding. Note the mouth opening shown here.

## Clinical Focus

### Resolution

Since amoxicillin has not resolved Barbara's case of pneumonia, the PA prescribes another antibiotic, azithromycin, which targets bacterial ribosomes rather than peptidoglycan. After taking the azithromycin as

directed, Barbara's symptoms resolve and she finally begins to feel like herself again. Presuming no drug resistance to amoxicillin was involved, and given the effectiveness of azithromycin, the causative agent of Barbara's pneumonia is most likely *Mycoplasma pneumoniae*. Even though this bacterium is a prokaryotic cell, it is not inhibited by amoxicillin because it does not have a cell wall and, therefore, does not make peptidoglycan.

Go back to the [previous Clinical Focus box](#).



## Check Your Understanding

- Explain how the cellular envelope of eukaryotic cells compares to that of prokaryotic cells.
- Explain the difference between eukaryotic and prokaryotic flagella.

## Summary

### 3.1 Spontaneous Generation

- The theory of **spontaneous generation** states that life arose from nonliving matter. It was a long-held belief dating back, within the written record, to Aristotle and the ancient Greeks, and persisted through the Middle Ages.
- Experimentation by Francesco Redi in the 17th century presented the first significant evidence refuting spontaneous generation by showing that flies must have access to meat for maggots to develop on the meat. Prominent scientists designed experiments and argued both in support of (John Needham) and against (Lazzaro Spallanzani) spontaneous generation.
- Louis Pasteur is credited with conclusively ending the controversy about the theory of spontaneous generation, with his famous swan-neck flask experiment. He subsequently proposed that “life only comes from life.”

### 3.2 Foundations of Modern Cell Theory

- Although cells were first observed in the 1660s by Robert Hooke, **cell theory** was not well accepted for another 200 years. The work of scientists such as Schleiden, Schwann, Remak, and Virchow contributed to its acceptance.
- **Endosymbiotic theory** states that mitochondria and chloroplasts, organelles found in many types of organisms, have their origins in bacteria. Significant structural and genetic information support this theory.
- The miasma theory of disease was widely accepted until the 19th century, when it was replaced by the **germ theory of disease** thanks to the work of Semmelweis, Snow, Pasteur, Lister, and Koch, and others.

### 3.3 Unique Characteristics of Prokaryotic Cells

- Prokaryotic cells differ from eukaryotic cells in that their genetic material is contained in a **nucleoid** rather than a membrane-bound nucleus. In addition, prokaryotic cells generally lack membrane-bound organelles.
- Prokaryotic cells of the same species typically share a similar **cell morphology** and **cellular arrangement**.
- Most prokaryotic cells have a **cell wall** that helps the organism maintain cellular morphology and protects it against changes in osmotic pressure.
- Outside of the nucleoid, prokaryotic cells may contain extrachromosomal DNA in **plasmids**.
- Prokaryotic **ribosomes** that are found in the cytoplasm have a size of 70S.
- Some prokaryotic cells have **inclusions** that store nutrients or chemicals for other uses.

- Some prokaryotic cells are able to form **endospores** through **sporulation** to survive in a dormant state when conditions are unfavorable. Endospores can **germinate**, transforming back into **vegetative cells** when conditions improve.
- In prokaryotic cells, the **cell envelope** includes a **plasma membrane** and usually a cell wall.
- Bacterial membranes are composed of phospholipids with integral or peripheral proteins. The fatty acid components of these phospholipids are ester-linked and are often used to identify specific types of bacteria. The proteins serve a variety of functions, including transport, cell-to-cell communication, and sensing environmental conditions. Archaeal membranes are distinct in that they are composed of fatty acids that are ether-linked to phospholipids.
- Some molecules can move across the bacterial membrane by simple diffusion, but most large molecules must be actively transported through membrane structures using cellular energy.
- Prokaryotic cell walls may be composed of **peptidoglycan** (bacteria) or **pseudopeptidoglycan** (archaea).
- Gram-positive bacterial cells are characterized by a thick **peptidoglycan** layer, whereas gram-negative bacterial cells are characterized by a thin peptidoglycan layer surrounded by an outer membrane.
- Some prokaryotic cells produce **glycocalyx** coatings, such as **capsules** and **slime layers**, that aid in attachment to surfaces and/or evasion of the host immune system.
- Some prokaryotic cells have **fimbriae** or **pili**, filamentous appendages that aid in attachment to surfaces. Pili are also used in the transfer of genetic material between cells.
- Some prokaryotic cells use one or more **flagella** to move through water. **Peritrichous** bacteria, which have numerous flagella, use **runs** and **tumbles** to move purposefully in the direction of a chemical attractant.

### 3.4 Unique Characteristics of Eukaryotic Cells

- Eukaryotic cells are defined by the presence of a **nucleus** containing the DNA genome and bound by a **nuclear membrane** (or **nuclear envelope**) composed of two lipid bilayers that regulate transport of materials into and out of the nucleus through nuclear pores.
- Eukaryotic cell morphologies vary greatly and may be maintained by various structures, including the cytoskeleton, the cell membrane, and/or the cell wall
- The **nucleolus**, located in the nucleus of eukaryotic cells, is the site of ribosomal synthesis and the first stages of ribosome assembly.
- Eukaryotic cells contain **80S ribosomes** in the rough endoplasmic reticulum (**membrane bound-ribosomes**) and cytoplasm (**free ribosomes**). They contain 70s ribosomes in mitochondria and chloroplasts.
- Eukaryotic cells have evolved an **endomembrane** system, containing membrane-bound organelles involved in transport. These include vesicles, the endoplasmic reticulum, and the Golgi apparatus.
- The **smooth endoplasmic reticulum** plays a role in lipid biosynthesis, carbohydrate metabolism, and detoxification of toxic compounds. The **rough endoplasmic reticulum** contains membrane-bound 80S ribosomes that synthesize proteins destined for the cell membrane
- The **Golgi apparatus** processes proteins and lipids, typically through the addition of sugar molecules, producing glycoproteins or glycolipids, components of the plasma membrane that are used in cell-to-cell communication.
- **Lysosomes** contain digestive enzymes that break down small particles ingested by **endocytosis**, large particles or cells ingested by **phagocytosis**, and damaged intracellular components.
- The **cytoskeleton**, composed of **microfilaments**, **intermediate filaments**, and **microtubules**, provides structural support in eukaryotic cells and serves as a network for transport of intracellular materials.
- **Centrosomes** are microtubule-organizing centers important in the formation of the mitotic spindle in mitosis.
- **Mitochondria** are the site of cellular respiration. They have two membranes: an outer membrane and an inner membrane with cristae. The mitochondrial matrix, within the inner membrane, contains the mitochondrial DNA, 70S ribosomes, and metabolic enzymes.



- The plasma membrane of eukaryotic cells is structurally similar to that found in prokaryotic cells, and membrane components move according to the fluid mosaic model. However, eukaryotic membranes contain sterols, which alter membrane fluidity, as well as glycoproteins and glycolipids, which help the cell recognize other cells and infectious particles.
- In addition to active transport and passive transport, eukaryotic cell membranes can take material into the cell via **endocytosis**, or expel matter from the cell via **exocytosis**.
- Cells of fungi, algae, plants, and some protists have a **cell wall**, whereas cells of animals and some protozoans have a sticky **extracellular matrix** that provides structural support and mediates cellular signaling.
- Eukaryotic flagella are structurally distinct from prokaryotic flagella but serve a similar purpose (locomotion). **Cilia** are structurally similar to eukaryotic flagella, but shorter; they may be used for locomotion, feeding, or movement of extracellular particles.

## Review Questions

### Multiple Choice

- Which of the following individuals argued in favor of the theory of spontaneous generation?
  - Francesco Redi
  - Louis Pasteur
  - John Needham
  - Lazzaro Spallanzani
- Which of the following individuals is credited for definitively refuting the theory of spontaneous generation using broth in swan-neck flask?
  - Aristotle
  - Jan Baptista van Helmont
  - John Needham
  - Louis Pasteur
- Which of the following experimented with raw meat, maggots, and flies in an attempt to disprove the theory of spontaneous generation.
  - Aristotle
  - Lazzaro Spallanzani
  - Antonie van Leeuwenhoek
  - Francesco Redi
- Which of the following individuals did not contribute to the establishment of cell theory?
  - Girolamo Fracastoro
  - Matthias Schleiden
  - Robert Remak
  - Robert Hooke
- Whose proposal of the endosymbiotic theory of mitochondrial and chloroplast origin was ultimately accepted by the greater scientific community?
  - Rudolf Virchow
  - Ignaz Semmelweis
  - Lynn Margulis
  - Theodor Schwann
- Which of the following developed a set of postulates for determining whether a particular disease is caused by a particular pathogen?
  - John Snow
  - Robert Koch
  - Joseph Lister
  - Louis Pasteur
- Which of the following terms refers to a prokaryotic cell that is comma shaped?
  - coccus
  - coccobacilli
  - vibrio
  - spirillum
- Which bacterial structures are important for adherence to surfaces? (Select all that apply.)
  - endospores
  - cell walls
  - fimbriae
  - capsules
  - flagella
- Which of the following cell wall components is unique to gram-negative cells?
  - lipopolysaccharide
  - teichoic acid
  - mycolic acid
  - peptidoglycan
- Which of the following terms refers to a bacterial cell having a single tuft of flagella at one end?
  - monotrichous
  - amphitrichous
  - peritrichous
  - lophotrichous



11. Bacterial cell walls are primarily composed of which of the following?

- a. phospholipid
- b. protein
- c. carbohydrate
- d. peptidoglycan

12. Which of the following organelles is not part of the endomembrane system?

- a. endoplasmic reticulum
- b. Golgi apparatus
- c. lysosome
- d. peroxisome

13. Which type of cytoskeletal fiber is important in the formation of the nuclear lamina?

- a. microfilaments
- b. intermediate filaments
- c. microtubules
- d. fibronectin

14. Sugar groups may be added to proteins in which of the following?

- a. smooth endoplasmic reticulum
- b. rough endoplasmic reticulum
- c. Golgi apparatus
- d. lysosome

15. Which of the following structures of a eukaryotic cell is not likely derived from endosymbiotic bacterium?

- a. mitochondrial DNA
- b. mitochondrial ribosomes
- c. inner membrane
- d. outer membrane

16. Which type of nutrient uptake involves the engulfment of small dissolved molecules into vesicles?

- a. active transport
- b. pinocytosis
- c. receptor-mediated endocytosis
- d. facilitated diffusion

17. Which of the following is not composed of microtubules?

- a. desmosomes
- b. centrioles
- c. eukaryotic flagella
- d. eukaryotic cilia

**True/False**

18. Exposure to air is necessary for microbial growth.

19. Bacteria have 80S ribosomes each composed of a 60S large subunit and a 40S small subunit.

20. Mitochondria in eukaryotic cells contain ribosomes that are structurally similar to those found in prokaryotic cells.

### Fill in the Blank

21. The assertion that “life only comes from life” was stated by Louis Pasteur in regard to his experiments that definitively refuted the theory of \_\_\_\_\_.

22. John Snow is known as the Father of \_\_\_\_\_.

23. The \_\_\_\_\_ theory states that disease may originate from proximity to decomposing matter and is not due to person-to-person contact.

24. The scientist who first described cells was \_\_\_\_\_.

25. Prokaryotic cells that are rod-shaped are called \_\_\_\_\_.

26. The type of inclusion containing polymerized inorganic phosphate is called \_\_\_\_\_.

27. Peroxisomes typically produce \_\_\_\_\_, a harsh chemical that helps break down molecules.

28. Microfilaments are composed of \_\_\_\_\_ monomers.

### Short Answer

29. Explain in your own words Pasteur’s swan-neck flask experiment.

30. Explain why the experiments of Needham and Spallanzani yielded in different results even though they used similar methodologies.

31. How did the explanation of Virchow and Remak for the origin of cells differ from that of Schleiden and Schwann?

32. What evidence exists that supports the endosymbiotic theory?

33. What were the differences in mortality rates due to puerperal fever that Ignaz Semmelweis observed? How did he propose to reduce the occurrence of puerperal fever? Did it work?

34. What is the direction of water flow for a bacterial cell living in a hypotonic environment? How do cell walls help bacteria living in such environments?

35. How do bacterial flagella respond to a chemical gradient of an attractant to move toward a higher concentration of the chemical?

36. What existing evidence supports the theory that mitochondria are of prokaryotic origin?

37. Why do eukaryotic cells require an endomembrane system?

38. Name at least two ways that prokaryotic flagella are different from eukaryotic flagella.


### Critical Thinking

39. What would the results of Pasteur’s swan-neck flask experiment have looked like if they supported the theory of spontaneous generation?

40. Why are mitochondria and chloroplasts unable to multiply outside of a host cell?

41. Why was the work of Snow so important in supporting the germ theory?

42. Label the parts of the prokaryotic cell.



43. Which of the following slides is a good example of staphylococci?




A



B



C




D

44. Provide some examples of bacterial structures that might be used as antibiotic targets and explain why.

45. The causative agent of botulism, a deadly form of food poisoning, is an endospore-forming bacterium called *Clostridium botulinum*. Why might it be difficult to kill this bacterium in contaminated food?


46. Label the following parts of this eukaryotic cell.



47. How are peroxisomes more like mitochondria than like the membrane-bound organelles of the endomembrane system? How do they differ from mitochondria?
48. Why must the functions of both lysosomes and peroxisomes be compartmentalized?

## Chapter 4

# Prokaryotic Diversity



**Figure 4.1** The bacterium *Shewanella* lives in the deep sea, where there is little oxygen diffused in the water. It is able to survive in this harsh environment by attaching to the sea floor and using long appendages, called “nanocables,” to sense oxygen. (credit b: modification of work by Liza Gross)

### Chapter Outline

- 4.1 Prokaryote Habitats, Relationships, and Microbiomes
- 4.2 Proteobacteria
- 4.3 Nonproteobacteria Gram-Negative Bacteria and Phototrophic Bacteria
- 4.4 Gram-Positive Bacteria
- 4.5 Deeply Branching Bacteria
- 4.6 Archaea

## Introduction

Scientists have studied prokaryotes for centuries, but it wasn't until 1966 that scientist Thomas Brock (1926–) discovered that certain bacteria can live in boiling water. This led many to wonder whether prokaryotes may also live in other extreme environments, such as at the bottom of the ocean, at high altitudes, or inside volcanoes, or even on other planets.

Prokaryotes have an important role in changing, shaping, and sustaining the entire biosphere. They can produce proteins and other substances used by molecular biologists in basic research and in medicine and industry. For example, the bacterium *Shewanella* lives in the deep sea, where oxygen is scarce. It grows long appendages, which have special sensors used to seek the limited oxygen in its environment. It can also digest toxic waste and generate electricity. Other species of prokaryotes can produce more oxygen than the entire Amazon rainforest, while still others supply plants, animals, and humans with usable forms of nitrogen; and inhabit our body, protecting us from harmful microorganisms and producing some vitally important substances. This chapter will examine the diversity, structure, and function of prokaryotes.



## 4.1 Prokaryote Habitats, Relationships, and Microbiomes

### Learning Objectives

- Identify and describe unique examples of prokaryotes in various habitats on earth
- Identify and describe symbiotic relationships
- Compare normal/commensal/resident microbiota to transient microbiota
- Explain how prokaryotes are classified

All living organisms are classified into three domains of life: Archaea, Bacteria, and Eukarya. In this chapter, we will focus on the domains Archaea and Bacteria. Archaea and bacteria are unicellular prokaryotic organisms. Unlike eukaryotes, they have no nuclei or any other membrane-bound organelles.

### Prokaryote Habitats and Functions

Prokaryotes are ubiquitous. They can be found everywhere on our planet, even in hot springs, in the Antarctic ice shield, and under extreme pressure two miles under water. One bacterium, *Paracoccus denitrificans*, has even been shown to survive when scientists removed it from its native environment (soil) and used a centrifuge to subject it to forces of gravity as strong as those found on the surface of Jupiter.

Prokaryotes also are abundant on and within the human body. According to a report by National Institutes of Health, prokaryotes, especially bacteria, outnumber human cells 10:1.<sup>[1]</sup> More recent studies suggest the ratio could be closer to 1:1,<sup>[2]</sup> but even that ratio means that there are a great number of bacteria within the human body. Bacteria thrive in the human mouth, nasal cavity, throat, ears, gastrointestinal tract, and vagina. Large colonies of bacteria can be found on healthy human skin, especially in moist areas (armpits, navel, and areas behind ears). However, even drier areas of the skin are not free from bacteria.

### Clinical Focus

#### Part 1

Marsha, a 20-year-old university student, recently returned to the United States from a trip to Nigeria, where she had interned as a medical assistant for an organization working to improve access to laboratory services for tuberculosis testing. When she returned, Marsha began to feel fatigue, which she initially attributed to jet lag. However, the fatigue persisted, and Marsha soon began to experience other bothersome symptoms, such as occasional coughing, night sweats, loss of appetite, and a low-grade fever of 37.4 °C (99.3 °F).

Marsha expected her symptoms would subside in a few days, but instead, they gradually became more severe. About two weeks after returning home, she coughed up some sputum and noticed that it contained blood and small whitish clumps resembling cottage cheese. Her fever spiked to 38.2 °C (100.8 °F), and she began feeling sharp pains in her chest when breathing deeply. Concerned that she seemed to be getting worse, Marsha scheduled an appointment with her physician.

- Could Marsha's symptoms be related to her overseas travel, even several weeks after returning home?

Jump to the **next** Clinical Focus box.

1. Medical Press. "Mouth Bacteria Can Change Their Diet, Supercomputers Reveal." August 12, 2014. <http://medicalxpress.com/news/2014-08-mouth-bacteria-diet-supercomputers-reveal.html>. Accessed February 24, 2015.

2. A. Abbott. "Scientists Bust Myth That Our Bodies Have More Bacteria Than Human Cells: Decades-Old Assumption about Microbiota Revisited." *Nature*. <http://www.nature.com/news/scientists-bust-myth-that-our-bodies-have-more-bacteria-than-human-cells-1.19136>. Accessed June 3, 2016.

The existence of prokaryotes is very important for the stability and thriving of ecosystems. For example, they are a necessary part of soil formation and stabilization processes through the breakdown of organic matter and development of biofilms. One gram of soil contains up to 10 billion microorganisms (most of them prokaryotic) belonging to about 1,000 species. Many species of bacteria use substances released from plant roots, such as acids and carbohydrates, as nutrients. The bacteria metabolize these plant substances and release the products of bacterial metabolism back to the soil, forming humus and thus increasing the soil's fertility. In salty lakes such as the Dead Sea (**Figure 4.2**), salt-loving halobacteria decompose dead brine shrimp and nourish young brine shrimp and flies with the products of bacterial metabolism.




**Figure 4.2** (a) Some prokaryotes, called halophiles, can thrive in extremely salty environments such as the Dead Sea, pictured here. (b) The archaeon *Halobacterium salinarum*, shown here in an electron micrograph, is a halophile that lives in the Dead Sea. (credit a: modification of work by Julien Menichini)

In addition to living in the ground and the water, prokaryotic microorganisms are abundant in the air, even high in the atmosphere. There may be up to 2,000 different kinds of bacteria in the air, similar to their diversity in the soil.

Prokaryotes can be found everywhere on earth because they are extremely resilient and adaptable. They are often metabolically flexible, which means that they might easily switch from one energy source to another, depending on the availability of the sources, or from one metabolic pathway to another. For example, certain prokaryotic cyanobacteria can switch from a conventional type of lipid metabolism, which includes production of fatty aldehydes, to a different type of lipid metabolism that generates biofuel, such as fatty acids and wax esters. Groundwater bacteria store complex high-energy carbohydrates when grown in pure groundwater, but they metabolize these molecules when the groundwater is enriched with phosphates. Some bacteria get their energy by reducing sulfates into sulfides, but can switch to a different metabolic pathway when necessary, producing acids and free hydrogen ions.

Prokaryotes perform functions vital to life on earth by capturing (or “fixing”) and recycling elements like carbon and nitrogen. Organisms such as animals require organic carbon to grow, but, unlike prokaryotes, they are unable to use inorganic carbon sources like carbon dioxide. Thus, animals rely on prokaryotes to convert carbon dioxide into organic carbon products that they can use. This process of converting carbon dioxide to organic carbon products is called carbon fixation.

Plants and animals also rely heavily on prokaryotes for nitrogen fixation, the conversion of atmospheric nitrogen into ammonia, a compound that some plants can use to form many different biomolecules necessary to their survival (**Figure 4.3**). Bacteria in the genus *Rhizobium*, for example, are nitrogen-fixing bacteria; they live in the roots of legume plants such as clover, alfalfa, and peas (**Figure 4.3**). Ammonia produced by *Rhizobium* helps these plants to survive by enabling them to make building blocks of nucleic acids. In turn, these plants may be eaten by animals—sustaining their growth and survival—or they may die, in which case the products of nitrogen fixation will enrich the soil and be used by other plants.



**Figure 4.3** (a) Nitrogen-fixing bacteria such as *Rhizobium* live in the root nodules of legumes such as clover. (b) This micrograph of the root nodule shows bacteroids (bacterium-like cells or modified bacterial cells) within the plant cells. The bacteroids are visible as darker ovals within the larger plant cell.

Another positive function of prokaryotes is in cleaning up the environment. Recently, some researchers focused on the diversity and functions of prokaryotes in manmade environments. They found that some bacteria play a unique role in degrading toxic chemicals that pollute water and soil.<sup>[3]</sup>

Despite all of the positive and helpful roles prokaryotes play, some are human pathogens that may cause illness or infection when they enter the body. In addition, some bacteria can contaminate food, causing spoilage or foodborne illness, which makes them subjects of concern in food preparation and safety. Less than 1% of prokaryotes (all of them bacteria) are thought to be human pathogens, but collectively these species are responsible for a large number of the diseases that afflict humans.

Besides pathogens, which have a direct impact on human health, prokaryotes also affect humans in many indirect ways. For example, prokaryotes are now thought to be key players in the processes of climate change. In recent years, as temperatures in the earth's polar regions have risen, soil that was formerly frozen year-round (permafrost) has begun to thaw. Carbon trapped in the permafrost is gradually released and metabolized by prokaryotes. This produces massive amounts of carbon dioxide and methane, greenhouse gases that escape into the atmosphere and contribute to the greenhouse effect.



### Check Your Understanding

1. In what types of environments can prokaryotes be found?
2. Name some ways that plants and animals rely on prokaryotes.

## Symbiotic Relationships

As we have learned, prokaryotic microorganisms can associate with plants and animals. Often, this association results in unique relationships between organisms. For example, bacteria living on the roots or leaves of a plant get nutrients from the plant and, in return, produce substances that protect the plant from pathogens. On the other hand, some bacteria are plant pathogens that use mechanisms of infection similar to bacterial pathogens of animals and humans.

Prokaryotes live in a **community**, or a group of interacting populations of organisms. A population is a group of individual organisms belonging to the same biological species and limited to a certain geographic area. Populations

3. A.M. Kravetz "Unique Bacteria Fights Man-Made Chemical Waste." 2012. <http://www.livescience.com/25181-bacteria-strain-cleans-up-toxins-nsf-bts.html>. Accessed March 9, 2015.

can have **cooperative interactions**, which benefit the populations, or **competitive interactions**, in which one population competes with another for resources. The study of these interactions between populations is called **microbial ecology**.

Any interaction between different species within a community is called **symbiosis**. Such interactions fall along a continuum between opposition and cooperation. Interactions in a symbiotic relationship may be beneficial or harmful, or have no effect on one or both of the species involved. **Table 4.1** summarizes the main types of symbiotic interactions among prokaryotes.

**Types of Symbiotic Relationships**

Type	Population A	Population B
Mutualism	Benefitted	Benefitted
Amensalism	Harmed	Unaffected
Commensalism	Benefitted	Unaffected
Neutralism	Unaffected	Unaffected
Parasitism	Benefitted	Harmed

**Table 4.1**

When two species benefit from each other, the symbiosis is called **mutualism** (or syntropy, or crossfeeding). For example, humans have a mutualistic relationship with the bacterium *Bacteroides thetaiotetraiotamicron*, which lives in the intestinal tract. *B. thetaiotetraiotamicron* digests complex polysaccharide plant materials that human digestive enzymes cannot break down, converting them into monosaccharides that can be absorbed by human cells. Humans also have a mutualistic relationship with certain strains of *Escherichia coli*, another bacterium found in the gut. *E. coli* relies on intestinal contents for nutrients, and humans derive certain vitamins from *E. coli*, particularly vitamin K, which is required for the formation of blood clotting factors. (This is only true for some strains of *E. coli*, however. Other strains are pathogenic and do not have a mutualistic relationship with humans.)

A type of symbiosis in which one population harms another but remains unaffected itself is called **amensalism**. In the case of bacteria, some amensalist species produce bactericidal substances that kill other species of bacteria. For example, the bacterium *Lucilia sericata* produces a protein that destroys *Staphylococcus aureus*, a bacterium commonly found on the surface of the human skin. Too much handwashing can affect this relationship and lead to *S. aureus* diseases and transmission.

In another type of symbiosis, called **commensalism**, one organism benefits while the other is unaffected. This occurs when the bacterium *Staphylococcus epidermidis* uses the dead cells of the human skin as nutrients. Billions of these bacteria live on our skin, but in most cases (especially when our immune system is healthy), we do not react to them in any way.

If neither of the symbiotic organisms is affected in any way, we call this type of symbiosis **neutralism**. An example of neutralism is the coexistence of metabolically active (vegetating) bacteria and endospores (dormant, metabolically passive bacteria). For example, the bacterium *Bacillus anthracis* typically forms endospores in soil when conditions are unfavorable. If the soil is warmed and enriched with nutrients, some endospores germinate and remain in symbiosis with other endospores that have not germinated.

A type of symbiosis in which one organism benefits while harming the other is called **parasitism**. The relationship between humans and many pathogenic prokaryotes can be characterized as parasitic because these organisms invade the body, producing toxic substances or infectious diseases that cause harm. Diseases such as tetanus, diphtheria, pertussis, tuberculosis, and leprosy all arise from interactions between bacteria and humans.

Scientists have coined the term **microbiome** to refer to all prokaryotic and eukaryotic microorganisms that are associated with a certain organism. Within the human microbiome, there are **resident microbiota** and **transient**

**microbiota.** The resident microbiota consists of microorganisms that constantly live in or on our bodies. The term transient microbiota refers to microorganisms that are only temporarily found in the human body, and these may include pathogenic microorganisms. Hygiene and diet can alter both the resident and transient microbiota.

The resident microbiota is amazingly diverse, not only in terms of the variety of species but also in terms of the preference of different microorganisms for different areas of the human body. For example, in the human mouth, there are thousands of commensal or mutualistic species of bacteria. Some of these bacteria prefer to inhabit the surface of the tongue, whereas others prefer the internal surface of the cheeks, and yet others prefer the front or back teeth or gums. The inner surface of the cheek has the least diverse microbiota because of its exposure to oxygen. By contrast, the crypts of the tongue and the spaces between teeth are two sites with limited oxygen exposure, so these sites have more diverse microbiota, including bacteria living in the absence of oxygen (e.g., *Bacteroides*, *Fusobacterium*). Differences in the oral microbiota between randomly chosen human individuals are also significant. Studies have shown, for example, that the prevalence of such bacteria as *Streptococcus*, *Haemophilus*, *Neisseria*, and others was dramatically different when compared between individuals.<sup>[4]</sup>

There are also significant differences between the microbiota of different sites of the same human body. The inner surface of the cheek has a predominance of *Streptococcus*, whereas in the throat, the palatine tonsil, and saliva, there are two to three times fewer *Streptococcus*, and several times more *Fusobacterium*. In the plaque removed from gums, the predominant bacteria belong to the genus *Fusobacterium*. However, in the intestine, both *Streptococcus* and *Fusobacterium* disappear, and the genus *Bacteroides* becomes predominant.

Not only can the microbiota vary from one body site to another, the microbiome can also change over time within the same individual. Humans acquire their first inoculations of normal flora during natural birth and shortly after birth. Before birth, there is a rapid increase in the population of *Lactobacilli* spp. in the vagina, and this population serves as the first colonization of microbiota during natural birth. After birth, additional microbes are acquired from health-care providers, parents, other relatives, and individuals who come in contact with the baby. This process establishes a microbiome that will continue to evolve over the course of the individual's life as new microbes colonize and are eliminated from the body. For example, it is estimated that within a 9-hour period, the microbiota of the small intestine can change so that half of the microbial inhabitants will be different.<sup>[5]</sup> The importance of the initial *Lactobacilli* colonization during vaginal child birth is highlighted by studies demonstrating a higher incidence of diseases in individuals born by cesarean section, compared to those born vaginally. Studies have shown that babies born vaginally are predominantly colonized by vaginal *Lactobacilli*, whereas babies born by cesarean section are more frequently colonized by microbes of the normal skin microbiota, including common hospital-acquired pathogens.

Throughout the body, resident microbiotas are important for human health because they occupy niches that might be otherwise taken by pathogenic microorganisms. For instance, *Lactobacilli* spp. are the dominant bacterial species of the normal vaginal microbiota for most women. *Lactobacilli* produce lactic acid, contributing to the acidity of the vagina and inhibiting the growth of pathogenic yeasts. However, when the population of the resident microbiota is decreased for some reason (e.g., because of taking antibiotics), the pH of the vagina increases, making it a more favorable environment for the growth of yeasts such as *Candida albicans*. Antibiotic therapy can also disrupt the microbiota of the intestinal tract and respiratory tract, increasing the risk for secondary infections and/or promoting the long-term carriage and shedding of pathogens.



### Check Your Understanding

1. Explain the difference between cooperative and competitive interactions in microbial communities.
2. List the types of symbiosis and explain how each population is affected.

---

4. E.M. Bik et al. "Bacterial Diversity in the Oral Cavity of 10 Healthy Individuals." *The ISME Journal* 4 no. 8 (2010):962–974.  
5. C.C. Booijink et al. "High Temporal and Intra-Individual Variation Detected in the Human Ileal Microbiota." *Environmental Microbiology* 12 no. 12 (2010):3213–3227.



## Taxonomy and Systematics

Assigning prokaryotes to a certain species is challenging. They do not reproduce sexually, so it is not possible to classify them according to the presence or absence of interbreeding. Also, they do not have many morphological features. Traditionally, the classification of prokaryotes was based on their shape, staining patterns, and biochemical or physiological differences. More recently, as technology has improved, the nucleotide sequences in genes have become an important criterion of microbial classification.

In 1923, American microbiologist David Hendricks Bergey (1860–1937) published *A Manual in Determinative Bacteriology*. With this manual, he attempted to summarize the information about the kinds of bacteria known at that time, using Latin binomial classification. Bergey also included the morphological, physiological, and biochemical properties of these organisms. His manual has been updated multiple times to include newer bacteria and their properties. It is a great aid in bacterial taxonomy and methods of characterization of bacteria. A more recent sister publication, the five-volume *Bergey's Manual of Systematic Bacteriology*, expands on Bergey's original manual. It includes a large number of additional species, along with up-to-date descriptions of the taxonomy and biological properties of all named prokaryotic taxa. This publication incorporates the approved names of bacteria as determined by the List of Prokaryotic Names with Standing in Nomenclature (LPSN).

### Link to Learning



*Bergey's Manual of Determinative Bacteriology* is now **available** (<https://openstaxcollege.org//22mandeterbact>) online. You can also access a searchable **database** (<https://openstaxcollege.org//22datamicrofst>) of microbial reference strains, published by the American Type Culture Collection (ATCC).

## Classification by Staining Patterns

According to their staining patterns, which depend on the properties of their cell walls, bacteria have traditionally been classified into gram-positive, gram-negative, and “atypical,” meaning neither gram-positive nor gram-negative. As explained in **Staining Microscopic Specimens**, gram-positive bacteria possess a thick peptidoglycan cell wall that retains the primary stain (crystal violet) during the decolorizing step; they remain purple after the gram-stain procedure because the crystal violet dominates the light red/pink color of the secondary counterstain, safranin. In contrast, gram-negative bacteria possess a thin peptidoglycan cell wall that does not prevent the crystal violet from washing away during the decolorizing step; therefore, they appear light red/pink after staining with the safranin. Bacteria that cannot be stained by the standard Gram stain procedure are called atypical bacteria. Included in the atypical category are species of *Mycoplasma* and *Chlamydia*, which lack a cell wall and therefore cannot retain the gram-stain reagents. *Rickettsia* are also considered atypical because they are too small to be evaluated by the Gram stain.

More recently, scientists have begun to further classify gram-negative and gram-positive bacteria. They have added a special group of deeply branching bacteria based on a combination of physiological, biochemical, and genetic features. They also now further classify gram-negative bacteria into Proteobacteria, *Cytophaga-Flavobacterium-Bacteroides* (CFB), and spirochetes.

The deeply branching bacteria are thought to be a very early evolutionary form of bacteria (see **Deeply Branching Bacteria**). They live in hot, acidic, ultraviolet-light-exposed, and anaerobic (deprived of oxygen) conditions. Proteobacteria is a phylum of very diverse groups of gram-negative bacteria; it includes some important human pathogens (e.g., *E. coli* and *Bordetella pertussis*). The CFB group of bacteria includes components of the normal human gut microbiota, like *Bacteroides*. The spirochetes are spiral-shaped bacteria and include the pathogen *Treponema pallidum*, which causes syphilis. We will characterize these groups of bacteria in more detail later in the chapter.

Based on their prevalence of guanine and cytosine nucleotides, gram-positive bacteria are also classified into low G+C and high G+C gram-positive bacteria. The low G+C gram-positive bacteria have less than 50% of guanine and cytosine nucleotides in their DNA. They include human pathogens, such as those that cause anthrax (*Bacillus anthracis*), tetanus (*Clostridium tetani*), and listeriosis (*Listeria monocytogenes*). High G+C gram-positive bacteria, which have more than 50% guanine and cytosine nucleotides in their DNA, include the bacteria that cause diphtheria (*Corynebacterium diphtheriae*), tuberculosis (*Mycobacterium tuberculosis*), and other diseases.

The classifications of prokaryotes are constantly changing as new species are being discovered. We will describe them in more detail, along with the diseases they cause, in later sections and chapters.



## Check Your Understanding

1. How do scientists classify prokaryotes?

### Micro Connections

#### Human Microbiome Project

The Human Microbiome Project was launched by the National Institutes of Health (NIH) in 2008. One main goal of the project is to create a large repository of the gene sequences of important microbes found in humans, helping biologists and clinicians understand the dynamics of the human microbiome and the relationship between the human microbiota and diseases. A network of labs working together has been compiling the data from swabs of several areas of the skin, gut, and mouth from hundreds of individuals.

One of the challenges in understanding the human microbiome has been the difficulty of culturing many of the microbes that inhabit the human body. It has been estimated that we are only able to culture 1% of the bacteria in nature and that we are unable to grow the remaining 99%. To address this challenge, researchers have used metagenomic analysis, which studies genetic material harvested directly from microbial communities, as opposed to that of individual species grown in a culture. This allows researchers to study the genetic material of all microbes in the microbiome, rather than just those that can be cultured.<sup>[6]</sup>

One important achievement of the Human Microbiome Project is establishing the first reference database on microorganisms living in and on the human body. Many of the microbes in the microbiome are beneficial, but some are not. It was found, somewhat unexpectedly, that all of us have some serious microbial pathogens in our microbiota. For example, the conjunctiva of the human eye contains 24 genera of bacteria and numerous pathogenic species.<sup>[7]</sup> A healthy human mouth contains a number of species of the genus *Streptococcus*, including pathogenic species *S. pyogenes* and *S. pneumoniae*.<sup>[8]</sup> This raises the question of why certain prokaryotic organisms exist commensally in certain individuals but act as deadly pathogens in others. Also unexpected was the number of organisms that had never been cultured. For example, in one metagenomic study of the human gut microbiota, 174 new species of bacteria were identified.<sup>[9]</sup>

Another goal for the near future is to characterize the human microbiota in patients with different diseases and to find out whether there are any relationships between the contents of an individual's microbiota and risk for or susceptibility to specific diseases. Analyzing the microbiome in a person with a specific disease may reveal new ways to fight diseases.

6. National Institutes of Health. "Human Microbiome Project. Overview." <http://commonfund.nih.gov/hmp/overview>. Accessed June 7, 2016.

7. Q. Dong et al. "Diversity of Bacteria at Healthy Human Conjunctiva." *Investigative Ophthalmology & Visual Science* 52 no. 8 (2011):5408–5413.

8. F.E. Dewhirst et al. "The Human Oral Microbiome." *Journal of Bacteriology* 192 no. 19 (2010):5002–5017.

9. J.C. Lagier et al. "Microbial Culturomics: Paradigm Shift in the Human Gut Microbiome Study." *Clinical Microbiology and Infection* 18 no. 12 (2012):1185–1193.

## 4.2 Proteobacteria

### Learning Objectives

- Describe the unique features of each class within the phylum Proteobacteria: Alphaproteobacteria, Betaproteobacteria, Gammaproteobacteria, Deltaproteobacteria, and Epsilonproteobacteria
- Give an example of a bacterium in each class of Proteobacteria


In 1987, the American microbiologist Carl Woese (1928–2012) suggested that a large and diverse group of bacteria that he called “purple bacteria and their relatives” should be defined as a separate phylum within the domain Bacteria based on the similarity of the nucleotide sequences in their genome.<sup>[10]</sup> This phylum of gram-negative bacteria subsequently received the name **Proteobacteria**. It includes many bacteria that are part of the normal human microbiota as well as many pathogens. The Proteobacteria are further divided into five classes: Alphaproteobacteria, Betaproteobacteria, Gammaproteobacteria, Deltaproteobacteria, and Epsilonproteobacteria (**Appendix D** (<http://cnx.org/content/m58949/latest/>)).

### Alphaproteobacteria

The first class of Proteobacteria is the **Alphaproteobacteria**. The unifying characteristic of this class is that they are **oligotrophs**, organisms capable of living in low-nutrient environments such as deep oceanic sediments, glacial ice, or deep undersurface soil.

Among the Alphaproteobacteria are two taxa, chlamydias and rickettsias, that are **obligate intracellular pathogens**, meaning that part of their life cycle must occur inside other cells called host cells. When not growing inside a host cell, *Chlamydia* and *Rickettsia* are metabolically inactive outside of the host cell. They cannot synthesize their own adenosine triphosphate (ATP), and, therefore, rely on cells for their energy needs.

*Rickettsia* spp. include a number of serious human pathogens. For example, *R. rickettsii* causes Rocky Mountain spotted fever, a life-threatening form of meningoencephalitis (inflammation of the membranes that wrap the brain). *R. rickettsii* infects ticks and can be transmitted to humans via a bite from an infected tick (**Figure 4.4**).



**Figure 4.4** Rickettsias require special staining methods to see them under a microscope. Here, *R. rickettsii*, which causes Rocky Mountain spotted fever, is shown infecting the cells of a tick.


Another species of *Rickettsia*, *R. prowazekii*, is spread by lice. It causes epidemic typhus, a severe infectious disease common during warfare and mass migrations of people. *R. prowazekii* infects human endothelium cells, causing inflammation of the inner lining of blood vessels, high fever, abdominal pain, and sometimes delirium. A relative, *R. typhi*, causes a less severe disease known as murine or endemic typhus, which is still observed in the southwestern United States during warm seasons.

10. C.R. Woese. “Bacterial Evolution.” *Microbiological Review* 51 no. 2 (1987):221–271.

*Chlamydia* is another taxon of the Alphaproteobacteria. Members of this genus are extremely resistant to the cellular defenses, giving them the ability to spread from host to host rapidly via elementary bodies. The metabolically and reproductively inactive **elementary bodies** are the endospore-like form of intracellular bacteria that enter an epithelial cell, where they become active. **Figure 4.5** illustrates the life cycle of *Chlamydia*.

*C. trachomatis* is a human pathogen that causes trachoma, a disease of the eyes, often leading to blindness. *C. trachomatis* also causes the sexually transmitted disease lymphogranuloma venereum (LGV). This disease is often mildly symptomatic, manifesting as regional lymph node swelling, or it may be asymptomatic, but it is extremely contagious and is common on college campuses.

**Table 4.2** summarizes the characteristics of important genera of Alphaproteobacteria.



**Figure 4.5** *Chlamydia* begins infection of a host when the metabolically inactive elementary bodies enter an epithelial cell. Once inside the host cell, the elementary bodies turn into active reticulate bodies. The reticulate bodies multiply and release more elementary bodies when the cell dies after the *Chlamydia* uses all of the host cell's ATP.

### Class Alphaproteobacteria

Genus	Microscopic Morphology	Unique Characteristics
<i>Agrobacterium</i>	Gram-negative bacillus	Plant pathogen; one species, <i>A. tumefaciens</i> , causes tumors in plants

**Table 4.2**

### Class Alphaproteobacteria

Genus	Microscopic Morphology	Unique Characteristics
<i>Bartonella</i>	Gram-negative, pleomorphic, flagellated coccobacillus	Facultative intracellular bacteria, transmitted by lice and fleas, cause trench fever and cat scratch disease in humans
<i>Brucella</i>	Gram-negative, small, flagellated coccobacillus	Facultative intracellular bacteria, transmitted by contaminated milk from infected cows, cause brucellosis in cattle and humans
<i>Caulobacter</i>	Gram-negative bacillus	Used in studies on cellular adaptation and differentiation because of its peculiar life cycle (during cell division, forms "swarm" cells and "stalked" cells)
<i>Chlamydia</i>	Gram-negative, coccoid or ovoid bacterium	Obligatory intracellular bacteria; some cause chlamydia, trachoma, and pneumonia
<i>Coxiella</i>	Small, gram-negative bacillus	Obligatory intracellular bacteria; cause Q fever; potential for use as biological weapon
<i>Ehrlichia</i>	Very small, gram-negative, coccoid or ovoid bacteria	Obligatory intracellular bacteria; can be transported from cell to cell; transmitted by ticks; cause ehrlichiosis (destruction of white blood cells and inflammation) in humans and dogs
<i>Hyphomicrobium</i>	Gram-negative bacilli; grows from a stalk	Similar to <i>Caulobacter</i>
<i>Methylocystis</i>	Gram-negative, coccoid or short bacilli	Nitrogen-fixing aerobic bacteria
<i>Rhizobium</i>	Gram-negative, rectangular bacilli with rounded ends forming clusters	Nitrogen-fixing bacteria that live in soil and form symbiotic relationship with roots of legumes (e.g., clover, alfalfa, and beans)
<i>Rickettsia</i>	Gram-negative, highly pleomorphic bacteria (may be cocci, rods, or threads)	Obligate intracellular bacteria; transmitted by ticks; may cause Rocky Mountain spotted fever and typhus

Table 4.2



### Check Your Understanding

1. What characteristic do all Alphaproteobacteria share?

### Betaproteobacteria

Unlike Alphaproteobacteria, which survive on a minimal amount of nutrients, the class **Betaproteobacteria** are **eutrophs** (or copiotrophs), meaning that they require a copious amount of organic nutrients. Betaproteobacteria often grow between aerobic and anaerobic areas (e.g., in mammalian intestines). Some genera include species that are human pathogens, able to cause severe, sometimes life-threatening disease. The genus *Neisseria*, for example, includes the bacteria *N. gonorrhoeae*, the causative agent of the STI gonorrhea, and *N. meningitidis*, the causative agent of bacterial meningitis.



*Neisseria* are cocci that live on mucosal surfaces of the human body. They are fastidious, or difficult to culture, and they require high levels of moisture, nutrient supplements, and carbon dioxide. Also, *Neisseria* are microaerophilic, meaning that they require low levels of oxygen. For optimal growth and for the purposes of identification, *Neisseria* spp. are grown on chocolate agar (i.e., agar supplemented by partially hemolyzed red blood cells). Their characteristic pattern of growth in culture is diplococcal: pairs of cells resembling coffee beans (**Figure 4.6**).



**Figure 4.6** *Neisseria meningitidis* growing in colonies on a chocolate agar plate.

The pathogen responsible for pertussis (whooping cough) is also a member of Betaproteobacteria. The bacterium *Bordetella pertussis*, from the order Burkholderiales, produces several toxins that paralyze the movement of cilia in the human respiratory tract and directly damage cells of the respiratory tract, causing a severe cough.

**Table 4.3** summarizes the characteristics of important genera of Betaproteobacteria.

#### Class Betaproteobacteria

Example Genus	Microscopic Morphology	Unique Characteristics
<i>Bordetella</i>	A small, gram-negative coccobacillus	Aerobic, very fastidious; <i>B. pertussis</i> causes pertussis (whooping cough)
<i>Burkholderia</i>	Gram-negative bacillus	Aerobic, aquatic, cause diseases in horses and humans (especially patients with cystic fibrosis); agents of nosocomial infections
<i>Leptothrix</i>	Gram-negative, sheathed, filamentous bacillus	Aquatic; oxidize iron and manganese; can live in wastewater treatment plants and clog pipes
<i>Neisseria</i>	Gram-negative, coffee bean-shaped coccus forming pairs	Require moisture and high concentration of carbon dioxide; oxidase positive, grow on chocolate agar; pathogenic species cause gonorrhea and meningitis
<i>Thiobacillus</i>	Gram-negative bacillus	Thermophilic, acidophilic, strictly aerobic bacteria; oxidize iron and sulfur

**Table 4.3**



### Check Your Understanding


1. What characteristic do all Betaproteobacteria share?

## Clinical Focus

### Part 2

When Marsha finally went to the doctor's office, the physician listened to her breathing through a stethoscope. He heard some crepitation (a crackling sound) in her lungs, so he ordered a chest radiograph and asked the nurse to collect a sputum sample for microbiological evaluation and cytology. The radiologic evaluation found cavities, opacities, and a particular pattern of distribution of abnormal material (**Figure 4.7**).

- What are some possible diseases that could be responsible for Marsha's radiograph results?



**Figure 4.7** This anteroposterior radiograph shows the presence of bilateral pulmonary infiltrate (white triangles) and “caving formation” (black arrows) present in the right apical region.

Jump to the **next** Clinical Focus box. Go back to the **previous** Clinical Focus box.


## Gammaproteobacteria

The most diverse class of gram-negative bacteria is **Gammaproteobacteria**, and it includes a number of human pathogens. For example, a large and diverse family, *Pseudomonaceae*, includes the genus *Pseudomonas*. Within this genus is the species *P. aeruginosa*, a pathogen responsible for diverse infections in various regions of the body. *P. aeruginosa* is a strictly aerobic, nonfermenting, highly motile bacterium. It often infects wounds and burns, can be the cause of chronic urinary tract infections, and can be an important cause of respiratory infections in patients with cystic fibrosis or patients on mechanical ventilators. Infections by *P. aeruginosa* are often difficult to treat because the bacterium is resistant to many antibiotics and has a remarkable ability to form biofilms. Other representatives of *Pseudomonas* include the fluorescent (glowing) bacterium *P. fluorescens* and the soil bacteria *P. putida*, which is known for its ability to degrade xenobiotics (substances not naturally produced or found in living organisms).

The *Pasteurellaceae* also includes several clinically relevant genera and species. This family includes several bacteria that are human and/or animal pathogens. For example, *Pasteurella haemolytica* causes severe pneumonia in sheep and goats. *P. multocida* is a species that can be transmitted from animals to humans through bites, causing infections of the skin and deeper tissues. The genus *Haemophilus* contains two human pathogens, *H. influenzae* and *H. ducreyi*. Despite its name, *H. influenzae* does not cause influenza (which is a viral disease). *H. influenzae* can cause both upper and lower respiratory tract infections, including sinusitis, bronchitis, ear infections, and pneumonia. Before the development of effective vaccination, strains of *H. influenzae* were a leading cause of more invasive diseases, like meningitis in children. *H. ducreyi* causes the STI known as chancroid.


The order Vibrionales includes the human pathogen *Vibrio cholerae*. This comma-shaped aquatic bacterium thrives in highly alkaline environments like shallow lagoons and sea ports. A toxin produced by *V. cholerae* causes

hypersecretion of electrolytes and water in the large intestine, leading to profuse watery diarrhea and dehydration. *V. parahaemolyticus* is also a cause of gastrointestinal disease in humans, whereas *V. vulnificus* causes serious and potentially life-threatening cellulitis (infection of the skin and deeper tissues) and blood-borne infections. Another representative of Vibrionales, *Aliivibrio fischeri*, engages in a symbiotic relationship with squid. The squid provides nutrients for the bacteria to grow and the bacteria produce bioluminescence that protects the squid from predators (**Figure 4.8**).



**Figure 4.8** (a) *Aliivibrio fischeri* is a bioluminescent bacterium. (b) *A. fischeri* colonizes and lives in a mutualistic relationship with the Hawaiian bobtail squid (*Euprymna scolopes*). (credit b: modification of work by Margaret McFall-Ngai)

The genus *Legionella* also belongs to the Gammaproteobacteria. *L. pneumophila*, the pathogen responsible for Legionnaires disease, is an aquatic bacterium that tends to inhabit pools of warm water, such as those found in the tanks of air conditioning units in large buildings (**Figure 4.9**). Because the bacteria can spread in aerosols, outbreaks of Legionnaires disease often affect residents of a building in which the water has become contaminated with *Legionella*. In fact, these bacteria derive their name from the first known outbreak of Legionnaires disease, which occurred in a hotel hosting an American Legion veterans' association convention in Philadelphia in 1976.




**Figure 4.9** (a) *Legionella pneumophila*, the causative agent of Legionnaires disease, thrives in warm water. (b) Outbreaks of Legionnaires disease often originate in the air conditioning units of large buildings when water in or near the system becomes contaminated with *L. pneumophila*.

Enterobacteriaceae is a large family of **enteric** (intestinal) bacteria belonging to the Gammaproteobacteria. They are facultative anaerobes and are able to ferment carbohydrates. Within this family, microbiologists recognize two distinct categories. The first category is called the coliforms, after its prototypical bacterium species, *Escherichia coli*. Coliforms are able to ferment lactose completely (i.e., with the production of acid and gas). The second category, noncoliforms, either cannot ferment lactose or can only ferment it incompletely (producing either acid or gas, but not

both). The noncoliforms include some notable human pathogens, such as *Salmonella* spp., *Shigella* spp., and *Yersinia pestis*.

*E. coli* has been perhaps the most studied bacterium since it was first described in 1886 by Theodor Escherich (1857–1911). Many strains of *E. coli* are in mutualistic relationships with humans. However, some strains produce a potentially deadly toxin called Shiga toxin, which perforates cellular membranes in the large intestine, causing bloody diarrhea and peritonitis (inflammation of the inner linings of the abdominal cavity). Other *E. coli* strains may cause traveler’s diarrhea, a less severe but very widespread disease.

The genus *Salmonella*, which belongs to the noncoliform group of *Enterobacteriaceae*, is interesting in that there is still no consensus about how many species it includes. Scientists have reclassified many of the groups they once thought to be species as **serotypes** (also called serovars), which are strains or variations of the same species of bacteria. Their classification is based on patterns of reactivity by animal antisera against molecules on the surface of the bacterial cells. A number of serotypes of *Salmonella* can cause salmonellosis, characterized by inflammation of the small and the large intestine, accompanied by fever, vomiting, and diarrhea. The species *S. enterobacterica* (serovar *typhi*) causes typhoid fever, with symptoms including fever, abdominal pain, and skin rashes (**Figure 4.10**).



**Figure 4.10** *Salmonella typhi* is the causative agent of typhoid fever.

**Table 4.4** summarizes the characteristics of important genera of Gammaproteobacteria.

#### Class Gammaproteobacteria

Example Genus	Microscopic Morphology	Unique Characteristics
<i>Beggiatoa</i>	Gram-negative bacteria; disc-shaped or cylindrical	Aquatic, live in water with high content of hydrogen disulfide; can cause problems for sewage treatment
<i>Enterobacter</i>	Gram-negative bacillus	Facultative anaerobe; cause urinary and respiratory tract infections in hospitalized patients; implicated in the pathogenesis of obesity
<i>Erwinia</i>	Gram-negative bacillus	Plant pathogen causing leaf spots and discoloration; may digest cellulose; prefer relatively low temperatures (25–30 °C)
<i>Escherichia</i>	Gram-negative bacillus	Facultative anaerobe; inhabit the gastrointestinal tract of warm-blooded animals; some strains are mutualists, producing vitamin K; others, like serotype <i>E. coli</i> O157:H7, are pathogens; <i>E. coli</i> has been a model organism for many studies in genetics and molecular biology

**Table 4.4**

### Class Gammaproteobacteria

Example Genus	Microscopic Morphology	Unique Characteristics
<i>Hemophilus</i>	Gram-negative bacillus	Pleomorphic, may appear as coccobacillus, aerobe, or facultative anaerobe; grow on blood agar; pathogenic species can cause respiratory infections, chancroid, and other diseases
<i>Klebsiella</i>	Gram-negative bacillus; appears rounder and thicker than other members of the <i>Enterobacteriaceae</i>	Facultative anaerobe, encapsulated, nonmotile; pathogenic species may cause pneumonia, especially in people with alcoholism
<i>Legionella</i>	Gram-negative bacillus	Fastidious, grow on charcoal-buffered yeast extract; <i>L. pneumophila</i> causes Legionnaires disease
<i>Methylobacter</i>	Gram-negative bacillus	Use methane as source of carbon and energy
<i>Proteus</i>	Gram-negative bacillus (pleomorphic)	Common inhabitants of the human gastrointestinal tract; motile; produce urease; opportunistic pathogens; may cause urinary tract infections and sepsis
<i>Pseudomonas</i>	Gram-negative bacillus	Aerobic; versatile; produce yellow and blue pigments, making them appear green in culture; opportunistic, antibiotic-resistant pathogens may cause wound infections, hospital-acquired infections, and secondary infections in patients with cystic fibrosis
<i>Serratia</i>	Gram-negative bacillus	Motile; may produce red pigment; opportunistic pathogens responsible for a large number of hospital-acquired infections
<i>Shigella</i>	Gram-negative bacillus	Nonmotile; dangerously pathogenic; produce Shiga toxin, which can destroy cells of the gastrointestinal tract; can cause dysentery
<i>Vibrio</i>	Gram-negative, comma- or curved rod-shaped bacteria	Inhabit seawater; flagellated, motile; may produce toxin that causes hypersecretion of water and electrolytes in the gastrointestinal tract; some species may cause serious wound infections
<i>Yersinia</i>	Gram-negative bacillus	Carried by rodents; human pathogens; <i>Y. pestis</i> causes bubonic plague and pneumonic plague; <i>Y. enterocolitica</i> can be a pathogen causing diarrhea in humans

Table 4.4



### Check Your Understanding

- List two families of Gammaproteobacteria.

### Deltaproteobacteria


The **Deltaproteobacteria** is a small class of gram-negative Proteobacteria that includes sulfate-reducing bacteria (SRBs), so named because they use sulfate as the final electron acceptor in the electron transport chain. Few SRBs are pathogenic. However, the SRB *Desulfovibrio orale* is associated with periodontal disease (disease of the gums).

Deltaproteobacteria also includes the genus *Bdellovibrio*, species of which are parasites of other gram-negative bacteria. *Bdellovibrio* invades the cells of the host bacterium, positioning itself in the periplasm, the space between



the plasma membrane and the cell wall, feeding on the host's proteins and polysaccharides. The infection is lethal for the host cells.

Another type of Deltaproteobacteria, myxobacteria, lives in the soil, scavenging inorganic compounds. Motile and highly social, they interact with other bacteria within and outside their own group. They can form multicellular, macroscopic “fruiting bodies” (**Figure 4.11**), structures that are still being studied by biologists and bacterial ecologists.<sup>[11]</sup> These bacteria can also form metabolically inactive myxospores.



**Figure 4.11** Myxobacteria form fruiting bodies. (credit: modification of work by Michiel Vos)

**Table 4.5** summarizes the characteristics of several important genera of Deltaproteobacteria.

#### Class Deltaproteobacteria

Genus	Microscopic Morphology	Unique characteristics
<i>Bdellovibrio</i>	Gram-negative, comma-shaped rod	Obligate aerobes; motile; parasitic (infecting other bacteria)
<i>Desulfovibrio</i> (formerly <i>Desulfuromonas</i> )	Gram-negative, comma-shaped rod	Reduce sulfur; can be used for removal of toxic and radioactive waste
<i>Myxobacterium</i>	Gram-negative, coccoid bacteria forming colonies (swarms)	Live in soil; can move by gliding; used as a model organism for studies of intercellular communication (signaling)

**Table 4.5**



### Check Your Understanding

1. What type of Deltaproteobacteria forms fruiting bodies?

11. H. Reichenbach. “Myxobacteria, Producers of Novel Bioactive Substances.” *Journal of Industrial Microbiology & Biotechnology* 27 no. 3 (2001):149–156.

## Epsilonproteobacteria

The smallest class of Proteobacteria is **Epsilonproteobacteria**, which are gram-negative microaerophilic bacteria (meaning they only require small amounts of oxygen in their environment). Two clinically relevant genera of Epsilonproteobacteria are *Campylobacter* and *Helicobacter*, both of which include human pathogens. *Campylobacter* can cause food poisoning that manifests as severe enteritis (inflammation in the small intestine). This condition, caused by the species *C. jejuni*, is rather common in developed countries, usually because of eating contaminated poultry products. Chickens often harbor *C. jejuni* in their gastrointestinal tract and feces, and their meat can become contaminated during processing.

Within the genus *Helicobacter*, the helical, flagellated bacterium *H. pylori* has been identified as a beneficial member of the stomach microbiota, but it is also the most common cause of chronic gastritis and ulcers of the stomach and duodenum (**Figure 4.12**). Studies have also shown that *H. pylori* is linked to stomach cancer.<sup>[12]</sup> *H. pylori* is somewhat unusual in its ability to survive in the highly acidic environment of the stomach. It produces urease and other enzymes that modify its environment to make it less acidic.



**Figure 4.12** *Helicobacter pylori* can cause chronic gastritis, which can lead to ulcers and stomach cancer.

**Table 4.6** summarizes the characteristics of the most clinically relevant genera of Epsilonproteobacteria.

### Class Epsilonproteobacteria

Example Genus	Microscopic Morphology	Unique Characteristics
<i>Campylobacter</i>	Gram-negative, spiral-shaped rod	Aerobic (microaerophilic); often infects chickens; may infect humans via undercooked meat, causing severe enteritis
<i>Helicobacter</i>	Gram-negative, spiral-shaped rod	Aerobic (microaerophilic) bacterium; can damage the inner lining of the stomach, causing chronic gastritis, peptic ulcers, and stomach cancer

**Table 4.6**



### Check Your Understanding

1. Name two Epsilonproteobacteria that cause gastrointestinal disorders.

12. S. Suerbaum, P. Michetti. “*Helicobacter pylori* infection.” *New England Journal of Medicine* 347 no. 15 (2002):1175–1186.

## 4.3 Nonproteobacteria Gram-Negative Bacteria and Phototrophic Bacteria

---


### Learning Objectives

- Describe the unique features of nonproteobacteria gram-negative bacteria
- Give an example of a nonproteobacteria bacterium in each category
- Describe the unique features of phototrophic bacteria
- Identify phototrophic bacteria

The majority of the gram-negative bacteria belong to the phylum Proteobacteria, discussed in the previous section. Those that do not are called the nonproteobacteria. In this section, we will describe three classes of gram-negative nonproteobacteria: the spirochetes, the CFB group, and the Planctomycetes. A diverse group of phototrophic bacteria that includes Proteobacteria and nonproteobacteria will be discussed at the end of this section.

### Spirochetes

Spirochetes are characterized by their long (up to 250  $\mu\text{m}$ ), spiral-shaped bodies. Most **spirochetes** are also very thin, which makes it difficult to examine gram-stained preparations under a conventional brightfield microscope. Darkfield fluorescent microscopy is typically used instead (**Figure 4.13**). Spirochetes are also difficult or even impossible to culture. They are highly motile, using their axial filament to propel themselves. The axial filament is similar to a flagellum, but it wraps around the cell and runs inside the cell body of a spirochete in the periplasmic space between the outer membrane and the plasma membrane (**Figure 4.13**).



**Figure 4.13** Spirochetes are typically observed using darkfield microscopy (left). However, electron microscopy (top center, bottom center) provides a more detailed view of their cellular morphology. The flagella found between the inner and outer membranes of spirochetes wrap around the bacterium, causing a twisting motion used for locomotion. (credit: modification of work by Guyard C, Raffel SJ, Schrumf ME, Dahlstrom E, Sturdevant D, Ricklefs SM, Martens C, Hayes SF, Fischer ER, Hansen BT, Porcella SF, Schwan TG)

Several genera of spirochetes include human pathogens. For example, the genus *Treponema* includes a species *T. pallidum*, which is further classified into four subspecies: *T. pallidum pallidum*, *T. pallidum pertenuis*, *T. pallidum carateum*, and *T. pallidum endemicum*. The subspecies *T. pallidum pallidum* causes the sexually transmitted infection known as syphilis, the third most prevalent sexually transmitted bacterial infection in the United States, after chlamydiosis and gonorrhea. The other subspecies of *T. pallidum* cause tropical infectious diseases of the skin, bones, and joints.

Another genus of spirochete, *Borrelia*, contains a number of pathogenic species. *B. burgdorferi* causes Lyme disease, which is transmitted by several genera of ticks (notably *Ixodes* and *Amblyomma*) and often produces a “bull’s eye” rash, fever, fatigue, and, sometimes, debilitating arthritis. *B. recurrentis* causes a condition known as relapsing fever. **Appendix D** (<http://cnx.org/content/m58949/latest/>) lists the genera, species, and related diseases for spirochetes.



### Check Your Understanding

1. Why do scientists typically use darkfield fluorescent microscopy to visualize spirochetes?

## Cytophaga, Fusobacterium, and Bacteroides

The gram-negative nonproteobacteria genera *Cytophaga*, *Fusobacterium*, and *Bacteroides* are classified together as a phylum and called the **CFB group**. Although they are phylogenetically diverse, bacteria of the CFB group share some similarities in the sequence of nucleotides in their DNA. They are rod-shaped bacteria adapted to anaerobic environments, such as the tissue of the gums, gut, and rumen of ruminating animals. CFB bacteria are avid fermenters, able to process cellulose in rumen, thus enabling ruminant animals to obtain carbon and energy from grazing.

*Cytophaga* are motile aquatic bacteria that glide. *Fusobacteria* inhabit the human mouth and may cause severe infectious diseases. The largest genus of the CFB group is *Bacteroides*, which includes dozens of species that are prevalent inhabitants of the human large intestine, making up about 30% of the entire gut microbiome (**Figure 4.14**). One gram of human feces contains up to 100 billion *Bacteroides* cells. Most *Bacteroides* are mutualistic. They benefit from nutrients they find in the gut, and humans benefit from their ability to prevent pathogens from colonizing the large intestine. Indeed, when populations of *Bacteroides* are reduced in the gut—as often occurs when a patient takes antibiotics—the gut becomes a more favorable environment for pathogenic bacteria and fungi, which can cause secondary infections.



**Figure 4.14** *Bacteroides* comprise up to 30% of the normal microbiota in the human gut.

Only a few species of *Bacteroides* are pathogenic. *B. melaninogenicus*, for example, can cause wound infections in patients with weakened immune systems.




### Check Your Understanding

1. Why are *Cytophaga*, *Fusobacterium*, and *Bacteroides* classified together as the CFB group?

## Planctomycetes

The Planctomycetes are found in aquatic environments, inhabiting freshwater, saltwater, and brackish water. Planctomycetes are unusual in that they reproduce by budding, meaning that instead of one maternal cell splitting into two equal daughter cells in the process of binary fission, the mother cell forms a bud that detaches from the mother cell and lives as an independent cell. These so-called swarmer cells are motile and not attached to a surface. However, they will soon differentiate into sessile (immobile) cells with an appendage called a holdfast that allows them to attach to surfaces in the water (**Figure 4.15**). Only the sessile cells are able to reproduce.





**Figure 4.15** (a) Sessile Planctomycetes have a holdfast that allows them to adhere to surfaces in aquatic environments. (b) Swimmers are motile and lack a holdfast.

**Table 4.7** summarizes the characteristics of some of the most clinically relevant genera of nonproteobacteria.

### Nonproteobacteria

Example Genus	Microscopic Morphology	Unique Characteristics
<i>Bacteroides</i>	Gram-negative bacillus	Obligate anaerobic bacteria; abundant in the human gastrointestinal tract; usually mutualistic, although some species are opportunistic pathogens
<i>Cytophaga</i>	Gram-negative bacillus	Motile by gliding; live in soil or water; decompose cellulose; may cause disease in fish
<i>Fusobacterium</i>	Gram-negative bacillus with pointed ends	Anaerobic; form; biofilms; some species cause disease in humans (periodontitis, ulcers)
<i>Leptospira</i>	Spiral-shaped bacterium (spirochetes); gram negative-like (better viewed by darkfield microscopy); very thin	Aerobic, abundant in shallow water reservoirs; infect rodents and domestic animals; can be transmitted to humans by infected animals' urine; may cause severe disease
<i>Sphingobacterium</i>	Gram-negative bacillus	Oxidase positive; nonmotile; contain high amounts of sphingophospholipids; rarely cause disease in humans
<i>Treponema</i>	Gram-negative-like spirochete; very thin; better viewed by darkfield microscopy	Motile; do not grow in culture; <i>T. pallidum</i> (subspecies <i>T. pallidum pallidum</i> ) causes syphilis

**Table 4.7**



### Check Your Understanding

1. How do Planctomycetes reproduce?

### Phototrophic Bacteria

The **phototrophic bacteria** are a large and diverse category of bacteria that do not represent a taxon but, rather, a group of bacteria that use sunlight as their primary source of energy. This group contains both Proteobacteria and Nonproteobacteria. They use solar energy to synthesize ATP through photosynthesis. When they produce oxygen,

they perform oxygenic photosynthesis. When they do not produce oxygen, they perform anoxygenic photosynthesis. With the exception of some cyanobacteria, the majority of phototrophic bacteria perform anoxygenic photosynthesis.

One large group of phototrophic bacteria includes the purple or green bacteria that perform photosynthesis with the help of **bacteriochlorophylls**, which are green, purple, or blue pigments similar to chlorophyll in plants. Some of these bacteria have a varying amount of red or orange pigments called carotenoids. Their color varies from orange to red to purple to green (**Figure 4.16**), and they are able to absorb light of various wavelengths. Traditionally, these bacteria are classified into sulfur and nonsulfur bacteria; they are further differentiated by color.



**Figure 4.16** Purple and green sulfur bacteria use bacteriochlorophylls to perform photosynthesis.

The sulfur bacteria perform anoxygenic photosynthesis, using sulfites as electron donors and releasing free elemental sulfur. Nonsulfur bacteria use organic substrates, such as succinate and malate, as donors of electrons.

The **purple sulfur bacteria** oxidize hydrogen sulfide into elemental sulfur and sulfuric acid and get their purple color from the pigments bacteriochlorophylls and carotenoids. Bacteria of the genus *Chromatium* are purple sulfur Gammaproteobacteria. These microorganisms are strict anaerobes and live in water. They use carbon dioxide as their only source of carbon, but their survival and growth are possible only in the presence of sulfites, which they use as electron donors. *Chromatium* has been used as a model for studies of bacterial photosynthesis since the 1950s.<sup>[13]</sup>

The **green sulfur bacteria** use sulfide for oxidation and produce large amounts of green bacteriochlorophyll. The genus *Chlorobium* is a green sulfur bacterium that is implicated in climate change because it produces methane, a greenhouse gas. These bacteria use at least four types of chlorophyll for photosynthesis. The most prevalent of these, bacteriochlorophyll, is stored in special vesicle-like organelles called chlorosomes.


Purple nonsulfur bacteria are similar to purple sulfur bacteria, except that they use hydrogen rather than hydrogen sulfide for oxidation. Among the **purple nonsulfur bacteria** is the genus *Rhodospirillum*. These microorganisms are facultative anaerobes, which are actually pink rather than purple, and can metabolize (“fix”) nitrogen. They may be valuable in the field of biotechnology because of their potential ability to produce biological plastic and hydrogen fuel.<sup>[14]</sup>

The **green nonsulfur bacteria** are similar to green sulfur bacteria but they use substrates other than sulfides for oxidation. *Chloroflexus* is an example of a green nonsulfur bacterium. It often has an orange color when it grows in the dark, but it becomes green when it grows in sunlight. It stores bacteriochlorophyll in chlorosomes, similar to *Chlorobium*, and performs anoxygenic photosynthesis, using organic sulfites (low concentrations) or molecular hydrogen as electron donors, so it can survive in the dark if oxygen is available. *Chloroflexus* does not have flagella but can glide, like *Cytophaga*. It grows at a wide range of temperatures, from 35 °C to 70 °C, thus can be thermophilic.

13. R.C. Fuller et al. “Carbon Metabolism in *Chromatium*.” *Journal of Biological Chemistry* 236 (1961):2140–2149.

14. T.T. Selao et al. “Comparative Proteomic Studies in *Rhodospirillum rubrum* Grown Under Different Nitrogen Conditions.” *Journal of Proteome Research* 7 no. 8 (2008):3267–3275.

Another large, diverse group of phototrophic bacteria compose the phylum **Cyanobacteria**; they get their blue-green color from the chlorophyll contained in their cells (**Figure 4.17**). Species of this group perform oxygenic photosynthesis, producing megatons of gaseous oxygen. Scientists hypothesize that cyanobacteria played a critical role in the change of our planet's anoxic atmosphere 1–2 billion years ago to the oxygen-rich environment we have today.<sup>[15]</sup>



**Figure 4.17** (a) *Microcystis aeruginosa* is a type of cyanobacteria commonly found in freshwater environments. (b) In warm temperatures, *M. aeruginosa* and other cyanobacteria can multiply rapidly and produce neurotoxins, resulting in blooms that are harmful to fish and other aquatic animals. (credit: modification of work by Dr. Barry H. Rosen)

Cyanobacteria have other remarkable properties. Amazingly adaptable, they thrive in many habitats, including marine and freshwater environments, soil, and even rocks. They can live at a wide range of temperatures, even in the extreme temperatures of the Antarctic. They can live as unicellular organisms or in colonies, and they can be filamentous, forming sheaths or biofilms. Many of them fix nitrogen, converting molecular nitrogen into nitrites and nitrates that other bacteria, plants, and animals can use. The reactions of nitrogen fixation occur in specialized cells called heterocysts.

Photosynthesis in Cyanobacteria is oxygenic, using the same type of chlorophyll a found in plants and algae as the primary photosynthetic pigment. Cyanobacteria also use phycocyanin and cyanophycin, two secondary photosynthetic pigments that give them their characteristic blue color. They are located in special organelles called phycobilisomes and in folds of the cellular membrane called thylakoids, which are remarkably similar to the photosynthetic apparatus of plants. Scientists hypothesize that plants originated from endosymbiosis of ancestral eukaryotic cells and ancestral photosynthetic bacteria.<sup>[16]</sup> Cyanobacteria are also an interesting object of research in biochemistry,<sup>[17]</sup> with studies investigating their potential as biosorbents<sup>[18]</sup> and products of human nutrition.<sup>[19]</sup>

Unfortunately, cyanobacteria can sometimes have a negative impact on human health. Genera such as *Microcystis* can form harmful cyanobacterial blooms, forming dense mats on bodies of water and producing large quantities of toxins that can harm wildlife and humans. These toxins have been implicated in tumors of the liver and diseases of the nervous system in animals and humans.<sup>[20]</sup>

15. A. De los Rios et al. "Ultrastructural and Genetic Characteristics of Endolithic Cyanobacterial Biofilms Colonizing Antarctic Granite Rocks." *FEMS Microbiology Ecology* 59 no. 2 (2007):386–395.

16. T. Cavalier-Smith. "Membrane Heredity and Early Chloroplast Evolution." *Trends in Plant Science* 5 no. 4 (2000):174–182.

17. S. Zhang, D.A. Bryant. "The Tricarboxylic Acid Cycle in Cyanobacteria." *Science* 334 no. 6062 (2011):1551–1553.

18. A. Cain et al. "Cyanobacteria as a Biosorbent for Mercuric Ion." *Bioresource Technology* 99 no. 14 (2008):6578–6586.

19. C.S. Ku et al. "Edible Blue-Green Algae Reduce the Production of Pro-Inflammatory Cytokines by Inhibiting NF-κB Pathway in Macrophages and Splenocytes." *Biochimica et Biophysica Acta* 1830 no. 4 (2013):2981–2988.

20. I. Stewart et al. Cyanobacterial Poisoning in Livestock, Wild Mammals and Birds – an Overview. *Advances in Experimental Medicine and Biology* 619 (2008):613–637.

**Table 4.8** summarizes the characteristics of important phototrophic bacteria.

**Phototrophic Bacteria**

Phylum	Class	Example Genus or Species	Common Name	Oxygenic or Anoxygenic	Sulfur Deposition
Cyanobacteria	Cyanophyceae	<i>Microcystis aeruginosa</i>	Blue-green bacteria	Oxygenic	None
Chlorobi	Chlorobia	<i>Chlorobium</i>	Green sulfur bacteria	Anoxygenic	Outside the cell
Chloroflexi (Division)	Chloroflexi	<i>Chloroflexus</i>	Green nonsulfur bacteria	Anoxygenic	None
Proteobacteria	Alphaproteobacteria	<i>Rhodospirillum</i>	Purple nonsulfur bacteria	Anoxygenic	None
	Betaproteobacteria	<i>Rhodocyclus</i>	Purple nonsulfur bacteria	Anoxygenic	None
	Gammaproteobacteria	<i>Chromatium</i>	Purple sulfur bacteria	Anoxygenic	Inside the cell

**Table 4.8**



### Check Your Understanding

1. What characteristic makes phototrophic bacteria different from other prokaryotes?

## 4.4 Gram-Positive Bacteria

### Learning Objectives

- Describe the unique features of each category of high G+C and low G+C gram-positive bacteria
- Identify similarities and differences between high G+C and low G+C bacterial groups
- Give an example of a bacterium of high G+C and low G+C group commonly associated with each category


Prokaryotes are identified as gram-positive if they have a multiple layer matrix of peptidoglycan forming the cell wall. Crystal violet, the primary stain of the Gram stain procedure, is readily retained and stabilized within this matrix, causing gram-positive prokaryotes to appear purple under a brightfield microscope after Gram staining. For many years, the retention of Gram stain was one of the main criteria used to classify prokaryotes, even though some prokaryotes did not readily stain with either the primary or secondary stains used in the Gram stain procedure.

Advances in nucleic acid biochemistry have revealed additional characteristics that can be used to classify gram-positive prokaryotes, namely the guanine to cytosine ratios (G+C) in DNA and the composition of 16S rRNA subunits. Microbiologists currently recognize two distinct groups of gram-positive, or weakly staining gram-positive, prokaryotes. The class Actinobacteria comprises the **high G+C gram-positive bacteria**, which have more than 50% guanine and cytosine nucleotides in their DNA. The class Bacilli comprises **low G+C gram-positive bacteria**, which have less than 50% of guanine and cytosine nucleotides in their DNA.

## Actinobacteria: High G+C Gram-Positive Bacteria

The name Actinobacteria comes from the Greek words for *rays* and *small rod*, but Actinobacteria are very diverse. Their microscopic appearance can range from thin filamentous branching rods to coccobacilli. Some Actinobacteria are very large and complex, whereas others are among the smallest independently living organisms. Most Actinobacteria live in the soil, but some are aquatic. The vast majority are aerobic. One distinctive feature of this group is the presence of several different peptidoglycans in the cell wall.

The genus *Actinomyces* is a much studied representative of Actinobacteria. *Actinomyces* spp. play an important role in soil ecology, and some species are human pathogens. A number of *Actinomyces* spp. inhabit the human mouth and are opportunistic pathogens, causing infectious diseases like periodontitis (inflammation of the gums) and oral abscesses. The species *A. israelii* is an anaerobe notorious for causing endocarditis (inflammation of the inner lining of the heart) (**Figure 4.18**).



**Figure 4.18** (a) *Actinomyces israelii* (false-color scanning electron micrograph [SEM]) has a branched structure. (b) *Corynebacterium diphtheria* causes the deadly disease diphtheria. Note the distinctive palisades. (c) The gram-variable bacterium *Gardnerella vaginalis* causes bacterial vaginosis in women. This micrograph shows a Pap smear from a woman with vaginosis. (credit a: modification of work by “GrahamColm”/Wikimedia Commons; credit c: modification of work by Mwakigonja AR, Torres LM, Mwakyoma HA, Kaaya EE)

The genus *Mycobacterium* is represented by bacilli covered with a mycolic acid coat. This waxy coat protects the bacteria from some antibiotics, prevents them from drying out, and blocks penetration by Gram stain reagents (see **Staining Microscopic Specimens**). Because of this, a special acid-fast staining procedure is used to visualize these bacteria. The genus *Mycobacterium* is an important cause of a diverse group of infectious diseases. *M. tuberculosis* is the causative agent of tuberculosis, a disease that primarily impacts the lungs but can infect other parts of the body as well. It has been estimated that one-third of the world’s population has been infected with *M. tuberculosis* and millions of new infections occur each year. Treatment of *M. tuberculosis* is challenging and requires patients to take a combination of drugs for an extended time. Complicating treatment even further is the development and spread of multidrug-resistant strains of this pathogen.

Another pathogenic species, *M. leprae*, is the cause of Hansen’s disease (leprosy), a chronic disease that impacts peripheral nerves and the integrity of the skin and mucosal surface of the respiratory tract. Loss of pain sensation and the presence of skin lesions increase susceptibility to secondary injuries and infections with other pathogens.

Bacteria in the genus *Corynebacterium* contain diaminopimelic acid in their cell walls, and microscopically often form *palisades*, or pairs of rod-shaped cells resembling the letter V. Cells may contain metachromatic granules, intracellular storage of inorganic phosphates that are useful for identification of *Corynebacterium*. The vast majority of *Corynebacterium* spp. are nonpathogenic; however, *C. diphtheria* is the causative agent of diphtheria, a disease that can be fatal, especially in children (**Figure 4.18**). *C. diphtheria* produces a toxin that forms a pseudomembrane in the patient’s throat, causing swelling, difficulty breathing, and other symptoms that can become serious if untreated.

The genus *Bifidobacterium* consists of filamentous anaerobes, many of which are commonly found in the gastrointestinal tract, vagina, and mouth. In fact, *Bifidobacterium* spp. constitute a substantial part of the human gut microbiota and are frequently used as probiotics and in yogurt production.



The genus *Gardnerella*, contains only one species, *G. vaginalis*. This species is defined as “gram-variable” because its small coccobacilli do not show consistent results when Gram stained (**Figure 4.18**). Based on its genome, it is placed into the high G+C gram-positive group. *G. vaginalis* can cause bacterial vaginosis in women; symptoms are typically mild or even undetectable, but can lead to complications during pregnancy.

**Table 4.9** summarizes the characteristics of some important genera of Actinobacteria. Additional information on Actinobacteria appears in **Appendix D** (<http://cnx.org/content/m58949/latest/>).

#### Actinobacteria: High G+C Gram-Positive

Example Genus	Microscopic Morphology	Unique Characteristics
<i>Actinomyces</i>	Gram-positive bacillus; in colonies, shows fungus-like threads (hyphae)	Facultative anaerobes; in soil, decompose organic matter; in the human mouth, may cause gum disease
<i>Arthrobacter</i>	Gram-positive bacillus (at the exponential stage of growth) or coccus (in stationary phase)	Obligate aerobes; divide by “snapping,” forming V-like pairs of daughter cells; degrade phenol, can be used in bioremediation
<i>Bifidobacterium</i>	Gram-positive, filamentous actinobacterium	Anaerobes commonly found in human gut microbiota
<i>Corynebacterium</i>	Gram-positive bacillus	Aerobes or facultative anaerobes; form palisades; grow slowly; require enriched media in culture; <i>C. diphtheriae</i> causes diphtheria
<i>Frankia</i>	Gram-positive, fungus-like (filamentous) bacillus	Nitrogen-fixing bacteria; live in symbiosis with legumes
<i>Gardnerella</i>	Gram-variable coccobacillus	Colonize the human vagina, may alter the microbial ecology, thus leading to vaginosis
<i>Micrococcus</i>	Gram-positive coccus, form microscopic clusters	Ubiquitous in the environment and on the human skin; oxidase-positive (as opposed to morphologically similar <i>S. aureus</i> ); some are opportunistic pathogens
<i>Mycobacterium</i>	Gram-positive, acid-fast bacillus	Slow growing, aerobic, resistant to drying and phagocytosis; covered with a waxy coat made of mycolic acid; <i>M. tuberculosis</i> causes tuberculosis; <i>M. leprae</i> causes leprosy
<i>Nocardia</i>	Weakly gram-positive bacillus; forms acid-fast branches	May colonize the human gingiva; may cause severe pneumonia and inflammation of the skin
<i>Propionibacterium</i>	Gram-positive bacillus	Aerotolerant anaerobe; slow-growing; <i>P. acnes</i> reproduces in the human sebaceous glands and may cause or contribute to acne
<i>Rhodococcus</i>	Gram-positive bacillus	Strict aerobe; used in industry for biodegradation of pollutants; <i>R. fascians</i> is a plant pathogen, and <i>R. equi</i> causes pneumonia in foals
<i>Streptomyces</i>	Gram-positive, fungus-like (filamentous) bacillus	Very diverse genus (>500 species); aerobic, spore-forming bacteria; scavengers, decomposers found in soil (give the soil its “earthy” odor); used in pharmaceutical industry as antibiotic producers (more than two-thirds of clinically useful antibiotics)

**Table 4.9**



## Check Your Understanding

1. What is one distinctive feature of Actinobacteria?

## Low G+C Gram-positive Bacteria

The low G+C gram-positive bacteria have less than 50% guanine and cytosine in their DNA, and this group of bacteria includes a number of genera of bacteria that are pathogenic.

### Clinical Focus

#### Part 3

Based on her symptoms, Marsha's doctor suspected that she had a case of tuberculosis. Although less common in the United States, tuberculosis is still extremely common in many parts of the world, including Nigeria. Marsha's work there in a medical lab likely exposed her to *Mycobacterium tuberculosis*, the bacterium that causes tuberculosis.

Marsha's doctor ordered her to stay at home, wear a respiratory mask, and confine herself to one room as much as possible. He also said that Marsha had to take one semester off school. He prescribed isoniazid and rifampin, antibiotics used in a drug cocktail to treat tuberculosis, which Marsha was to take three times a day for at least three months.


- Why did the doctor order Marsha to stay home for three months?

Jump to the **next** Clinical Focus box. Go back to the **previous** Clinical Focus box.

## Clostridia

One large and diverse class of low G+C gram-positive bacteria is Clostridia. The best studied genus of this class is *Clostridium*. These rod-shaped bacteria are generally obligate anaerobes that produce endospores and can be found in anaerobic habitats like soil and aquatic sediments rich in organic nutrients. The endospores may survive for many years.

*Clostridium* spp. produce more kinds of protein toxins than any other bacterial genus, and several species are human pathogens. *C. perfringens* is the third most common cause of food poisoning in the United States and is the causative agent of an even more serious disease called gas gangrene. Gas gangrene occurs when *C. perfringens* endospores enter a wound and germinate, becoming viable bacterial cells and producing a toxin that can cause the necrosis (death) of tissue. *C. tetani*, which causes tetanus, produces a neurotoxin that is able to enter neurons, travel to regions of the central nervous system where it blocks the inhibition of nerve impulses involved in muscle contractions, and cause a life-threatening spastic paralysis. *C. botulinum* produces botulinum neurotoxin, the most lethal biological toxin known. Botulinum toxin is responsible for rare but frequently fatal cases of botulism. The toxin blocks the release of acetylcholine in neuromuscular junctions, causing flaccid paralysis. In very small concentrations, botulinum toxin has been used to treat muscle pathologies in humans and in a cosmetic procedure to eliminate wrinkles. *C. difficile* is a common source of hospital-acquired infections (**Figure 4.19**) that can result in serious and even fatal cases of colitis (inflammation of the large intestine). Infections often occur in patients who are immunosuppressed or undergoing antibiotic therapy that alters the normal microbiota of the gastrointestinal tract. **Appendix D (<http://cnx.org/content/m58949/latest/>)** lists the genera, species, and related diseases for Clostridia.




**Figure 4.19** *Clostridium difficile*, a gram-positive, rod-shaped bacterium, causes severe colitis and diarrhea, often after the normal gut microbiota is eradicated by antibiotics.

### Lactobacillales

The order Lactobacillales comprises low G+C gram-positive bacteria that include both bacilli and cocci in the genera *Lactobacillus*, *Leuconostoc*, *Enterococcus*, and *Streptococcus*. Bacteria of the latter three genera typically are spherical or ovoid and often form chains.

*Streptococcus*, the name of which comes from the Greek word for *twisted chain*, is responsible for many types of infectious diseases in humans. Species from this genus, often referred to as streptococci, are usually classified by serotypes called Lancefield groups, and by their ability to lyse red blood cells when grown on blood agar.

*S. pyogenes* belongs to the Lancefield group A,  $\beta$ -hemolytic *Streptococcus*. This species is considered a pyogenic pathogen because of the associated pus production observed with infections it causes (**Figure 4.20**). *S. pyogenes* is the most common cause of bacterial pharyngitis (strep throat); it is also an important cause of various skin infections that can be relatively mild (e.g., impetigo) or life threatening (e.g., necrotizing fasciitis, also known as flesh eating disease), life threatening.



**Figure 4.20** (a) A gram-stained specimen of *Streptococcus pyogenes* shows the chains of cocci characteristic of this organism's morphology. (b) *S. pyogenes* on blood agar shows characteristic lysis of red blood cells, indicated by the halo of clearing around colonies.

The nonpyogenic (i.e., not associated with pus production) streptococci are a group of streptococcal species that are not a taxon but are grouped together because they inhabit the human mouth. The nonpyogenic streptococci do not belong to any of the Lancefield groups. Most are commensals, but a few, such as *S. mutans*, are implicated in the development of dental caries.


*S. pneumoniae* (commonly referred to as pneumococcus), is a *Streptococcus* species that also does not belong to any Lancefield group. *S. pneumoniae* cells appear microscopically as diplococci, pairs of cells, rather than the long chains typical of most streptococci. Scientists have known since the 19th century that *S. pneumoniae* causes pneumonia and other respiratory infections. However, this bacterium can also cause a wide range of other diseases, including meningitis, septicemia, osteomyelitis, and endocarditis, especially in newborns, the elderly, and patients with immunodeficiency.

## Bacilli

The name of the class Bacilli suggests that it is made up of bacteria that are bacillus in shape, but it is a morphologically diverse class that includes bacillus-shaped and coccus-shaped genera. Among the many genera in this class are two that are very important clinically: *Bacillus* and *Staphylococcus*.

Bacteria in the genus *Bacillus* are bacillus in shape and can produce endospores. They include aerobes or facultative anaerobes. A number of *Bacillus* spp. are used in various industries, including the production of antibiotics (e.g., barnase), enzymes (e.g., alpha-amylase, BamH1 restriction endonuclease), and detergents (e.g., subtilisin).


Two notable pathogens belong to the genus *Bacillus*. *B. anthracis* is the pathogen that causes anthrax, a severe disease that affects wild and domesticated animals and can spread from infected animals to humans. Anthrax manifests in humans as charcoal-black ulcers on the skin, severe enterocolitis, pneumonia, and brain damage due to swelling. If untreated, anthrax is lethal. *B. cereus*, a closely related species, is a pathogen that may cause food poisoning. It is a rod-shaped species that forms chains. Colonies appear milky white with irregular shapes when cultured on blood agar (**Figure 4.21**). One other important species is *B. thuringiensis*. This bacterium produces a number of substances used as insecticides because they are toxic for insects.



**Figure 4.21** (a) In this gram-stained specimen, the violet rod-shaped cells forming chains are the gram-positive bacteria *Bacillus cereus*. The small, pink cells are the gram-negative bacteria *Escherichia coli*. (b) In this culture, white colonies of *B. cereus* have been grown on sheep blood agar. (credit a: modification of work by “Bibliomaniac 15”/Wikimedia Commons)

The genus *Staphylococcus* also belongs to the class Bacilli, even though its shape is coccus rather than a bacillus. The name *Staphylococcus* comes from a Greek word for *bunches of grapes*, which describes their microscopic appearance in culture (**Figure 4.22**). *Staphylococcus* spp. are facultative anaerobic, halophilic, and nonmotile. The two best-studied species of this genus are *S. epidermidis* and *S. aureus*.





**Figure 4.22** This SEM of *Staphylococcus aureus* illustrates the typical “grape-like” clustering of cells.

*S. epidermidis*, whose main habitat is the human skin, is thought to be nonpathogenic for humans with healthy immune systems, but in patients with immunodeficiency, it may cause infections in skin wounds and prostheses (e.g., artificial joints, heart valves). *S. epidermidis* is also an important cause of infections associated with intravenous catheters. This makes it a dangerous pathogen in hospital settings, where many patients may be immunocompromised.

Strains of *S. aureus* cause a wide variety of infections in humans, including skin infections that produce boils, carbuncles, cellulitis, or impetigo. Certain strains of *S. aureus* produce a substance called enterotoxin, which can cause severe enteritis, often called staph food poisoning. Some strains of *S. aureus* produce the toxin responsible for toxic shock syndrome, which can result in cardiovascular collapse and death.

Many strains of *S. aureus* have developed resistance to antibiotics. Some antibiotic-resistant strains are designated as methicillin-resistant *S. aureus* (MRSA) and vancomycin-resistant *S. aureus* (VRSA). These strains are some of the most difficult to treat because they exhibit resistance to nearly all available antibiotics, not just methicillin and vancomycin. Because they are difficult to treat with antibiotics, infections can be lethal. MRSA and VRSA are also contagious, posing a serious threat in hospitals, nursing homes, dialysis facilities, and other places where there are large populations of elderly, bedridden, and/or immunocompromised patients. **Appendix D** (<http://cnx.org/content/m58949/latest/>) lists the genera, species, and related diseases for bacilli.

## Mycoplasmas

Although *Mycoplasma* spp. do not possess a cell wall and, therefore, are not stained by Gram-stain reagents, this genus is still included with the low G+C gram-positive bacteria. The genus *Mycoplasma* includes more than 100 species, which share several unique characteristics. They are very small cells, some with a diameter of about 0.2  $\mu\text{m}$ , which is smaller than some large viruses. They have no cell walls and, therefore, are **pleomorphic**, meaning that they may take on a variety of shapes and can even resemble very small animal cells. Because they lack a characteristic shape, they can be difficult to identify. One species, *M. pneumoniae*, causes the mild form of pneumonia known as “walking pneumonia” or “atypical pneumonia.” This form of pneumonia is typically less severe than forms caused by other bacteria or viruses.

**Table 4.10** summarizes the characteristics of notable genera low G+C Gram-positive bacteria.



### Bacilli: Low G+C Gram-Positive Bacteria

Example Genus	Microscopic Morphology	Unique Characteristics
<i>Bacillus</i>	Large, gram-positive bacillus	Aerobes or facultative anaerobes; form endospores; <i>B. anthracis</i> causes anthrax in cattle and humans, <i>B. cereus</i> may cause food poisoning
<i>Clostridium</i>	Gram-positive bacillus	Strict anaerobes; form endospores; all known species are pathogenic, causing tetanus, gas gangrene, botulism, and colitis
<i>Enterococcus</i>	Gram-positive coccus; forms microscopic pairs in culture (resembling <i>Streptococcus pneumoniae</i> )	Anaerobic aerotolerant bacteria, abundant in the human gut, may cause urinary tract and other infections in the nosocomial environment
<i>Lactobacillus</i>	Gram-positive bacillus	Facultative anaerobes; ferment sugars into lactic acid; part of the vaginal microbiota; used as probiotics
<i>Leuconostoc</i>	Gram-positive coccus; may form microscopic chains in culture	Fermenter, used in food industry to produce sauerkraut and kefir
<i>Mycoplasma</i>	The smallest bacteria; appear pleomorphic under electron microscope	Have no cell wall; classified as low G+C Gram-positive bacteria because of their genome; <i>M. pneumoniae</i> causes “walking” pneumonia
<i>Staphylococcus</i>	Gram-positive coccus; forms microscopic clusters in culture that resemble bunches of grapes	Tolerate high salt concentration; facultative anaerobes; produce catalase; <i>S. aureus</i> can also produce coagulase and toxins responsible for local (skin) and generalized infections
<i>Streptococcus</i>	Gram-positive coccus; forms chains or pairs in culture	Diverse genus; classified into groups based on sharing certain antigens; some species cause hemolysis and may produce toxins responsible for human local (throat) and generalized disease
<i>Ureaplasma</i>	Similar to <i>Mycoplasma</i>	Part of the human vaginal and lower urinary tract microbiota; may cause inflammation, sometimes leading to internal scarring and infertility

Table 4.10



### Check Your Understanding

1. Name some ways in which streptococci are classified.
2. Name one pathogenic low G+C gram-positive bacterium and a disease it causes.


## Clinical Focus

### Resolution

Marsha's sputum sample was sent to the microbiology lab to confirm the identity of the microorganism causing her infection. The lab also performed antimicrobial susceptibility testing (AST) on the sample to confirm that the physician has prescribed the correct antimicrobial drugs.

Direct microscopic examination of the sputum revealed acid-fast bacteria (AFB) present in Marsha's sputum. When placed in culture, there were no signs of growth for the first 8 days, suggesting that microorganism was either dead or growing very slowly. Slow growth is a distinctive characteristic of *M. tuberculosis*.

After four weeks, the lab microbiologist observed distinctive colorless granulated colonies (**Figure 4.23**). The colonies contained AFB showing the same microscopic characteristics as those revealed during the direct microscopic examination of Marsha's sputum. To confirm the identification of the AFB, samples of the colonies were analyzed using nucleic acid hybridization, or direct nucleic acid amplification (NAA) testing. When a bacterium is acid-fast, it is classified in the family *Mycobacteriaceae*. DNA sequencing of variable genomic regions of the DNA extracted from these bacteria revealed that it was high G+C. This fact served to finalize Marsha's diagnosis as infection with *M. tuberculosis*. After nine months of treatment with the drugs prescribed by her doctor, Marsha made a full recovery.



**Figure 4.23** *M. tuberculosis* grows on Löwenstein-Jensen (LJ) agar in distinct colonies.

Go back to the [previous](#) Clinical Focus box.

## Eye on Ethics



### Biopiracy and Bioprospecting

In 1969, an employee of a Swiss pharmaceutical company was vacationing in Norway and decided to collect some soil samples. He took them back to his lab, and the Swiss company subsequently used the fungus *Tolypocladium inflatum* in those samples to develop cyclosporine A, a drug widely used in patients who undergo tissue or organ transplantation. The Swiss company earns more than \$1 billion a year for production of cyclosporine A, yet Norway receives nothing in return—no payment to the government or benefit for the

Norwegian people. Despite the fact the cyclosporine A saves numerous lives, many consider the means by which the soil samples were obtained to be an act of “biopiracy,” essentially a form of theft. Do the ends justify the means in a case like this?

Nature is full of as-yet-undiscovered bacteria and other microorganisms that could one day be used to develop new life-saving drugs or treatments.<sup>[21]</sup> Pharmaceutical and biotechnology companies stand to reap huge profits from such discoveries, but ethical questions remain. To whom do biological resources belong? Should companies who invest (and risk) millions of dollars in research and development be required to share revenue or royalties for the right to access biological resources?

Compensation is not the only issue when it comes to bioprospecting. Some communities and cultures are philosophically opposed to bioprospecting, fearing unforeseen consequences of collecting genetic or biological material. Native Hawaiians, for example, are very protective of their unique biological resources.

For many years, it was unclear what rights government agencies, private corporations, and citizens had when it came to collecting samples of microorganisms from public land. Then, in 1993, the Convention on Biological Diversity granted each nation the rights to any genetic and biological material found on their own land. Scientists can no longer collect samples without a prior arrangement with the land owner for compensation. This convention now ensures that companies act ethically in obtaining the samples they use to create their products.

## 4.5 Deeply Branching Bacteria

---


### Learning Objectives

- Describe the unique features of deeply branching bacteria
- Give examples of significant deeply branching bacteria

On a phylogenetic tree (see **A Systematic Approach**), the trunk or root of the tree represents a common ancient evolutionary ancestor, often called the last universal common ancestor (LUCA), and the branches are its evolutionary descendants. Scientists consider the **deeply branching bacteria**, such as the genus *Acetothermus*, to be the first of these non-LUCA forms of life produced by evolution some 3.5 billion years ago. When placed on the phylogenetic tree, they stem from the common root of life, deep and close to the LUCA root—hence the name “deeply branching” (**Figure 4.24**).

---

21. J. Andre. *Bioethics as Practice*. Chapel Hill, NC: University of North Carolina Press, 2002.



**Figure 4.24** The star on this phylogenetic tree of life shows the position of the deeply branching bacteria *Acetothermus*.


The deeply branching bacteria may provide clues regarding the structure and function of ancient and now extinct forms of life. We can hypothesize that ancient bacteria, like the deeply branching bacteria that still exist, were thermophiles or hyperthermophiles, meaning that they thrived at very high temperatures. *Acetothermus paucivorans*, a gram-negative anaerobic bacterium discovered in 1988 in sewage sludge, is a thermophile growing at an optimal temperature of 58 °C.<sup>[22]</sup> Scientists have determined it to be the deepest branching bacterium, or the closest evolutionary relative of the LUCA (**Figure 4.24**).

The class Aquificae includes deeply branching bacteria that are adapted to the harshest conditions on our planet, resembling the conditions thought to dominate the earth when life first appeared. Bacteria from the genus *Aquifex* are hyperthermophiles, living in hot springs at a temperature higher than 90 °C. The species *A. pyrophilus* thrives near underwater volcanoes and thermal ocean vents, where the temperature of water (under high pressure) can reach 138 °C. *Aquifex* bacteria use inorganic substances as nutrients. For example, *A. pyrophilus* can reduce oxygen, and it is able to reduce nitrogen in anaerobic conditions. They also show a remarkable resistance to ultraviolet light and ionizing radiation. Taken together, these observations support the hypothesis that the ancient ancestors of deeply branching bacteria began evolving more than 3 billion years ago, when the earth was hot and lacked an atmosphere, exposing the bacteria to nonionizing and ionizing radiation.

The class Thermotogae is represented mostly by hyperthermophilic, as well as some mesophilic (preferring moderate temperatures), anaerobic gram-negative bacteria whose cells are wrapped in a peculiar sheath-like outer membrane called a toga. The thin layer of peptidoglycan in their cell wall has an unusual structure; it contains diaminopimelic acid and D-lysine. These bacteria are able to use a variety of organic substrates and produce molecular hydrogen, which can be used in industry. The class contains several genera, of which the best known is the genus *Thermotoga*. One species of this genus, *T. maritima*, lives near the thermal ocean vents and thrives in temperatures of 90 °C; another species, *T. subterranea*, lives in underground oil reservoirs.

Finally, the deeply branching bacterium *Deinococcus radiodurans* belongs to a genus whose name is derived from a Greek word meaning *terrible berry*. Nicknamed “Conan the Bacterium,” *D. radiodurans* is considered a polyextremophile because of its ability to survive under the many different kinds of extreme conditions—extreme heat, drought, vacuum, acidity, and radiation. It owes its name to its ability to withstand doses of ionizing radiation that kill all other known bacteria; this special ability is attributed to some unique mechanisms of DNA repair.

22. G. Dietrich et al. “*Acetothermus paucivorans*, gen. nov., sp. Nov., a Strictly Anaerobic, Thermophilic Bacterium From Sewage Sludge, Fermenting Hexoses to Acetate, CO<sub>2</sub>, and H<sub>2</sub>.” *Systematic and Applied Microbiology* 10 no. 2 (1988):174–179.



**Figure 4.25** *Deinococcus radiodurans*, or “Conan the Bacterium,” survives in the harshest conditions on earth.

## 4.6 Archaea

### Learning Objectives

- Describe the unique features of each category of Archaea
- Explain why archaea might not be associated with human microbiomes or pathology
- Give common examples of archaea commonly associated with unique environmental habitats

Like organisms in the domain Bacteria, organisms of the domain **Archaea** are all unicellular organisms. However, archaea differ structurally from bacteria in several significant ways, as discussed in **Unique Characteristics of Prokaryotic Cells**. To summarize:

- The archaeal cell membrane is composed of ether linkages with branched isoprene chains (as opposed to the bacterial cell membrane, which has ester linkages with unbranched fatty acids).
- Archaeal cell walls lack peptidoglycan, but some contain a structurally similar substance called pseudopeptidoglycan or pseudomurein.
- The genomes of Archaea are larger and more complex than those of bacteria.

Domain Archaea is as diverse as domain Bacteria, and its representatives can be found in any habitat. Some archaea are mesophiles, and many are extremophiles, preferring extreme hot or cold, extreme salinity, or other conditions that are hostile to most other forms of life on earth. Their metabolism is adapted to the harsh environments, and they can perform methanogenesis, for example, which bacteria and eukaryotes cannot.

The size and complexity of the archaeal genome makes it difficult to classify. Most taxonomists agree that within the Archaea, there are currently five major phyla: Crenarchaeota, Euryarchaeota, Korarchaeota, Nanoarchaeota, and Thaumarchaeota. There are likely many other archaeal groups that have not yet been systematically studied and classified.


With few exceptions, archaea are not present in the human microbiota, and none are currently known to be associated with infectious diseases in humans, animals, plants, or microorganisms. However, many play important roles in the environment and may thus have an indirect impact on human health.



## Crenarchaeota

Crenarchaeota is a class of Archaea that is extremely diverse, containing genera and species that differ vastly in their morphology and requirements for growth. All Crenarchaeota are aquatic organisms, and they are thought to be the most abundant microorganisms in the oceans. Most, but not all, Crenarchaeota are hyperthermophiles; some of them (notably, the genus *Pyrolobus*) are able to grow at temperatures up to 113 °C.<sup>[23]</sup>

Archaea of the genus *Sulfolobus* (Figure 4.26) are thermophiles that prefer temperatures around 70–80°C and acidophiles that prefer a pH of 2–3.<sup>[24]</sup> *Sulfolobus* can live in aerobic or anaerobic environments. In the presence of oxygen, *Sulfolobus* spp. use metabolic processes similar to those of heterotrophs. In anaerobic environments, they oxidize sulfur to produce sulfuric acid, which is stored in granules. *Sulfolobus* spp. are used in biotechnology for the production of thermostable and acid-resistant proteins called affitins.<sup>[25]</sup> Affitins can bind and neutralize various antigens (molecules found in toxins or infectious agents that provoke an immune response from the body).



**Figure 4.26** *Sulfolobus*, an archaeon of the class Crenarchaeota, oxidizes sulfur and stores sulfuric acid in its granules.

Another genus, *Thermoproteus*, is represented by strictly anaerobic organisms with an optimal growth temperature of 85 °C. They have flagella and, therefore, are motile. *Thermoproteus* has a cellular membrane in which lipids form a monolayer rather than a bilayer, which is typical for archaea. Its metabolism is autotrophic. To synthesize ATP, *Thermoproteus* spp. reduce sulfur or molecular hydrogen and use carbon dioxide or carbon monoxide as a source of carbon. *Thermoproteus* is thought to be the deepest-branching genus of Archaea, and thus is a living example of some of our planet's earliest forms of life.



### Check Your Understanding

1. What types of environments do Crenarchaeota prefer?

23. E. Blochl et al. "Pyrolobus fumani, gen. and sp. nov., represents a novel group of Archaea, extending the upper temperature limit for life to 113°C." *Extremophiles* 1 (1997):14–21.

24. T.D. Brock et al. "Sulfolobus: A New Genus of Sulfur-Oxidizing Bacteria Living at Low pH and High Temperature." *Archiv für Mikrobiologie* 84 no. 1 (1972):54–68.

25. S. Pacheco et al. "Affinity Transfer to the Archaeal Extremophilic Sac7d Protein by Insertion of a CDR." *Protein Engineering Design and Selection* 27 no. 10 (2014):431-438.

## Euryarchaeota

The phylum Euryarchaeota includes several distinct classes. Species in the classes Methanobacteria, Methanococci, and Methanomicrobia represent Archaea that can be generally described as methanogens. Methanogens are unique in that they can reduce carbon dioxide in the presence of hydrogen, producing methane. They can live in the most extreme environments and can reproduce at temperatures varying from below freezing to boiling. Methanogens have been found in hot springs as well as deep under ice in Greenland. Some scientists have even hypothesized that **methanogens** may inhabit the planet Mars because the mixture of gases produced by methanogens resembles the makeup of the Martian atmosphere.<sup>[26]</sup>

Methanogens are thought to contribute to the formation of anoxic sediments by producing hydrogen sulfide, making “marsh gas.” They also produce gases in ruminants and humans. Some genera of methanogens, notably *Methanosarcina*, can grow and produce methane in the presence of oxygen, although the vast majority are strict anaerobes.

The class Halobacteria (which was named before scientists recognized the distinction between Archaea and Bacteria) includes halophilic (“salt-loving”) archaea. Halobacteria require a very high concentrations of sodium chloride in their aquatic environment. The required concentration is close to saturation, at 36%; such environments include the Dead Sea as well as some salty lakes in Antarctica and south-central Asia. One remarkable feature of these organisms is that they perform photosynthesis using the protein bacteriorhodopsin, which gives them, and the bodies of water they inhabit, a beautiful purple color (**Figure 4.27**).



**Figure 4.27** Halobacteria growing in these salt ponds gives them a distinct purple color. (credit: modification of work by Tony Hisgett)

Notable species of Halobacteria include *Halobacterium salinarum*, which may be the oldest living organism on earth; scientists have isolated its DNA from fossils that are 250 million years old.<sup>[27]</sup> Another species, *Haloferax volcanii*, shows a very sophisticated system of ion exchange, which enables it to balance the concentration of salts at high temperatures.

---

26. R.R. Britt “Crater Critters: Where Mars Microbes Might Lurk.” <http://www.space.com/1880-crater-critters-mars-microbes-lurk.html>. Accessed April 7, 2015.

27. H. Vreeland et al. “Fatty acid and DA Analyses of Permian Bacterium Isolated From Ancient Salt Crystals Reveal Differences With Their Modern Relatives.” *Extremophiles* 10 (2006):71–78.



## Check Your Understanding

1. Where do Halobacteria live?

### Micro Connections

#### Finding a Link Between Archaea and Disease

Archaea are not known to cause any disease in humans, animals, plants, bacteria, or in other archaea. Although this makes sense for the extremophiles, not all archaea live in extreme environments. Many genera and species of Archaea are mesophiles, so they can live in human and animal microbiomes, although they rarely do. As we have learned, some methanogens exist in the human gastrointestinal tract. Yet we have no reliable evidence pointing to any archaean as the causative agent of any human disease.

Still, scientists have attempted to find links between human disease and archaea. For example, in 2004, Lepp et al. presented evidence that an archaean called *Methanobrevibacter oralis* inhabits the gums of patients with periodontal disease. The authors suggested that the activity of these methanogens causes the disease.<sup>[28]</sup> However, it was subsequently shown that there was no causal relationship between *M. oralis* and periodontitis. It seems more likely that periodontal disease causes an enlargement of anaerobic regions in the mouth that are subsequently populated by *M. oralis*.<sup>[29]</sup>

There remains no good answer as to why archaea do not seem to be pathogenic, but scientists continue to speculate and hope to find the answer.

## Summary

### 4.1 Prokaryote Habitats, Relationships, and Microbiomes

- Prokaryotes are unicellular microorganisms whose cells have no nucleus.
- Prokaryotes can be found everywhere on our planet, even in the most extreme environments.
- Prokaryotes are very flexible metabolically, so they are able to adjust their feeding to the available natural resources.
- Prokaryotes live in **communities** that interact among themselves and with large organisms that they use as hosts (including humans).
- The totality of forms of prokaryotes (particularly bacteria) living on the human body is called the human microbiome, which varies between regions of the body and individuals, and changes over time.
- The totality of forms of prokaryotes (particularly bacteria) living in a certain region of the human body (e.g., mouth, throat, gut, eye, vagina) is called the **microbiota** of this region.
- Prokaryotes are classified into domains Archaea and Bacteria.
- In recent years, the traditional approaches to classification of prokaryotes have been supplemented by approaches based on molecular genetics.

### 4.2 Proteobacteria

- **Proteobacteria** is a phylum of gram-negative bacteria discovered by Carl Woese in the 1980s based on nucleotide sequence homology.

28. P.W. Lepp et al. "Methanogenic Archaea and Human Gum Disease." *Proceedings of the National Academies of Science of the United States of America* 101 no. 16 (2004):6176–6181.

29. R.I. Aminov. "Role of Archaea in Human Disease." *Frontiers in Cellular and Infection Microbiology* 3 (2013):42.

- Proteobacteria are further classified into the classes alpha-, beta-, gamma-, delta- and epsilonproteobacteria, each class having separate orders, families, genera, and species.
- **Alphaproteobacteria** are **oligotrophs**. The taxa chlamydias and rickettsias are **obligate intracellular pathogens**, feeding on cells of host organisms; they are metabolically inactive outside of the host cell. Some Alphaproteobacteria can convert atmospheric nitrogen to nitrites, making nitrogen usable by other forms of life.
- **Betaproteobacteria** are **eutrophs**. They include human pathogens of the genus *Neisseria* and the species *Bordetella pertussis*.
- **Gammaproteobacteria** are the largest and the most diverse group of Proteobacteria. Many are human pathogens that are aerobes or facultative anaerobes. Some Gammaproteobacteria are **enteric** bacteria that may be coliform or noncoliform. *Escherichia coli*, a member of Gammaproteobacteria, is perhaps the most studied bacterium.
- **Deltaproteobacteria** make up a small group able to reduce sulfate or elemental sulfur. Some are scavengers and form myxospores, with multicellular fruiting bodies.
- **Epsilonproteobacteria** make up the smallest group of Proteobacteria. The genera *Campylobacter* and *Helicobacter* are human pathogens.

#### 4.3 Nonproteobacteria Gram-Negative Bacteria and Phototrophic Bacteria

- Gram-negative nonproteobacteria include the taxa **spirochetes**; the *Cytophaga*, *Fusobacterium*, *Bacteroides* group; Planctomycetes; and many representatives of **phototrophic bacteria**.
- Spirochetes are motile, spiral bacteria with a long, narrow body; they are difficult or impossible to culture.
- Several genera of spirochetes contain human pathogens that cause such diseases as syphilis and Lyme disease.
- *Cytophaga*, *Fusobacterium*, and *Bacteroides* are classified together as a phylum called the **CFB group**. They are rod-shaped anaerobic organoheterotrophs and avid fermenters. *Cytophaga* are aquatic bacteria with the gliding motility. *Fusobacteria* inhabit the human mouth and may cause severe infectious diseases. *Bacteroides* are present in vast numbers in the human gut, most of them being mutualistic but some are pathogenic.
- Planctomycetes are aquatic bacteria that reproduce by budding; they may form large colonies, and develop a holdfast.
- Phototrophic bacteria are not a taxon but, rather, a group categorized by their ability to use the energy of sunlight. They include Proteobacteria and nonproteobacteria, as well as sulfur and nonsulfur bacteria colored purple or green.
- Sulfur bacteria perform anoxygenic photosynthesis, using sulfur compounds as donors of electrons, whereas nonsulfur bacteria use organic compounds (succinate, malate) as donors of electrons.
- Some phototrophic bacteria are able to fix nitrogen, providing the usable forms of nitrogen to other organisms.
- **Cyanobacteria** are oxygen-producing bacteria thought to have played a critical role in the forming of the earth's atmosphere.

#### 4.4 Gram-Positive Bacteria

- Gram-positive bacteria are a very large and diverse group of microorganisms. Understanding their taxonomy and knowing their unique features is important for diagnostics and treatment of infectious diseases.
- Gram-positive bacteria are classified into **high G+C gram-positive** and **low G+C gram-positive** bacteria, based on the prevalence of guanine and cytosine nucleotides in their genome
- Actinobacteria is the taxonomic name of the class of high G+C gram-positive bacteria. This class includes the genera *Actinomyces*, *Arthrobacter*, *Corynebacterium*, *Frankia*, *Gardnerella*, *Micrococcus*, *Mycobacterium*, *Nocardia*, *Propionibacterium*, *Rhodococcus*, and *Streptomyces*. Some representatives of these genera are used in industry; others are human or animal pathogens.
- Examples of high G+C gram-positive bacteria that are human pathogens include *Mycobacterium tuberculosis*, which causes tuberculosis; *M. leprae*, which causes leprosy (Hansen's disease); and *Corynebacterium diphtheriae*, which causes diphtheria.

- *Clostridia* spp. are low G+C gram-positive bacteria that are generally obligate anaerobes and can form endospores. Pathogens in this genus include *C. perfringens* (gas gangrene), *C. tetani* (tetanus), and *C. botulinum* (botulism).
- Lactobacillales include the genera *Enterococcus*, *Lactobacillus*, *Leuconostoc*, and *Streptococcus*. *Streptococcus* is responsible for many human diseases, including pharyngitis (strep throat), scarlet fever, rheumatic fever, glomerulonephritis, pneumonia, and other respiratory infections.
- Bacilli is a taxonomic class of low G+C gram-positive bacteria that include rod-shaped and coccus-shaped species, including the genera *Bacillus* and *Staphylococcus*. *B. anthracis* causes anthrax, *B. cereus* may cause opportunistic infections of the gastrointestinal tract, and *S. aureus* strains can cause a wide range of infections and diseases, many of which are highly resistant to antibiotics.
- *Mycoplasma* spp. are very small, **pleomorphic** low G+C gram-positive bacteria that lack cell walls. *M. pneumoniae* causes atypical pneumonia.

#### 4.5 Deeply Branching Bacteria

- **Deeply branching bacteria** are phylogenetically the most ancient forms of life, being the closest to the last universal common ancestor.
- Deeply branching bacteria include many species that thrive in extreme environments that are thought to resemble conditions on earth billions of years ago
- Deeply branching bacteria are important for our understanding of evolution; some of them are used in industry

#### 4.6 Archaea

- **Archaea** are unicellular, prokaryotic microorganisms that differ from bacteria in their genetics, biochemistry, and ecology.
- Some archaea are extremophiles, living in environments with extremely high or low temperatures, or extreme salinity.
- Only archaea are known to produce methane. Methane-producing archaea are called **methanogens**.
- Halophilic archaea prefer a concentration of salt close to saturation and perform photosynthesis using bacteriorhodopsin.
- Some archaea, based on fossil evidence, are among the oldest organisms on earth.
- Archaea do not live in great numbers in human microbiomes and are not known to cause disease.

## Review Questions

### Multiple Choice

- The term prokaryotes refers to which of the following?
  - very small organisms
  - unicellular organisms that have no nucleus
  - multicellular organisms
  - cells that resemble animal cells more than plant cells
- The term microbiota refers to which of the following?
  - all microorganisms of the same species
  - all of the microorganisms involved in a symbiotic relationship
  - all microorganisms in a certain region of the human body
  - all microorganisms in a certain geographic region
- Which of the following refers to the type of interaction between two prokaryotic populations in which one population benefits and the other is not affected?
  - mutualism
  - commensalism
  - parasitism
  - neutralism
- Which of the following describes Proteobacteria in domain Bacteria?
  - phylum
  - class
  - species
  - genus



5. All Alphaproteobacteria are which of the following?
- oligotrophs
  - intracellular
  - pathogenic
  - all of the above
  - none of the above
6. Class Betaproteobacteria includes all but which of the following genera?
- Neisseria*.
  - Bordetella*.
  - Leptothrix*.
  - Campylobacter*.
7. *Haemophilus influenzae* is a common cause of which of the following?
- influenza
  - dysentery
  - upper respiratory tract infections
  - hemophilia
8. Which of the following is the organelle that spirochetes use to propel themselves?
- plasma membrane
  - axial filament
  - pilum
  - fimbria
9. Which of the following bacteria are the most prevalent in the human gut?
- cyanobacteria
  - staphylococci
  - Borrelia*
  - Bacteroides*
10. Which of the following refers to photosynthesis performed by bacteria with the use of water as the donor of electrons?
- oxygenic
  - anoxygenic
  - heterotrophic
  - phototrophic
11. Which of the following bacterial species is classified as high G+C gram-positive?
- Corynebacterium diphtheriae*
  - Staphylococcus aureus*
  - Bacillus anthracis*
  - Streptococcus pneumoniae*
12. The term “deeply branching” refers to which of the following?
- the cellular shape of deeply branching bacteria
  - the position in the evolutionary tree of deeply branching bacteria
  - the ability of deeply branching bacteria to live in deep ocean waters
  - the pattern of growth in culture of deeply branching bacteria
13. Which of these deeply branching bacteria is considered a polyextremophile?
- Aquifex pyrophilus*
  - Deinococcus radiodurans*
  - Staphylococcus aureus*
  - Mycobacterium tuberculosis*
14. Archaea and Bacteria are most similar in terms of their \_\_\_\_\_.
- genetics
  - cell wall structure
  - ecology
  - unicellular structure
15. Which of the following is true of archaea that produce methane?
- They reduce carbon dioxide in the presence of nitrogen.
  - They live in the most extreme environments.
  - They are always anaerobes.
  - They have been discovered on Mars.

**True/False**

16. Among prokaryotes, there are some that can live in every environment on earth.

**Fill in the Blank**

17. When prokaryotes live as interacting communities in which one population benefits to the harm of the other, the type of symbiosis is called \_\_\_\_\_.

18. The domain \_\_\_\_\_ does not include prokaryotes.

19. Pathogenic bacteria that are part of the transient microbiota can sometimes be eliminated by \_\_\_\_\_ therapy.

20. Nitrogen-fixing bacteria provide other organisms with usable nitrogen in the form of \_\_\_\_\_.

21. Rickettsias are \_\_\_\_\_ intracellular bacteria.

22. The species \_\_\_\_\_, which belongs to Epsilonproteobacteria, causes peptic ulcers of the stomach and duodenum.

23. The genus *Salmonella* belongs to the class \_\_\_\_\_ and includes pathogens that cause salmonellosis and typhoid fever.

24. The bacterium that causes syphilis is called \_\_\_\_\_.

25. Bacteria in the genus *Rhodospirillum* that use hydrogen for oxidation and fix nitrogen are \_\_\_\_\_ bacteria.

26. *Streptococcus* is the \_\_\_\_\_ of bacteria that is responsible for many human diseases.

27. One species of *Streptococcus*, *S. pyogenes*, is classified as a \_\_\_\_\_ pathogen due to the characteristic production of pus in infections it causes.

28. *Propionibacterium* belongs to \_\_\_\_\_ G+C gram-positive bacteria. One of its species is used in the food industry and another causes acne.

29. The length of the branches of the evolutionary tree characterizes the evolutionary \_\_\_\_\_ between organisms.

30. The deeply branching bacteria are thought to be the form of life closest to the last universal \_\_\_\_\_.

31. Many of the deeply branching bacteria are aquatic and hyperthermophilic, found near underwater volcanoes and thermal ocean \_\_\_\_\_.

32. The deeply branching bacterium *Deinococcus radiodurans* is able to survive exposure to high doses of \_\_\_\_\_.

33. \_\_\_\_\_ is a genus of Archaea. Its optimal environmental temperature ranges from 70 °C to 80 °C, and its optimal pH is 2–3. It oxidizes sulfur and produces sulfuric acid.

34. \_\_\_\_\_ was once thought to be the cause of periodontal disease, but, more recently, the causal relationship between this archaean and the disease was not confirmed.

**Short Answer**

35. Compare commensalism and amensalism.

36. Give an example of the changes of human microbiota that result from medical intervention.

37. What is the metabolic difference between coliforms and noncoliforms? Which category contains several species of intestinal pathogens?

38. Why are *Mycoplasma* and *Chlamydia* classified as obligate intracellular pathogens?

39. Explain the term CFB group and name the genera that this group includes.

40. Name and briefly describe the bacterium that causes Lyme disease.
41. Characterize the phylum Cyanobacteria.
42. Name and describe two types of *S. aureus* that show multiple antibiotic resistance.
43. Briefly describe the significance of deeply branching bacteria for basic science and for industry.
44. What is thought to account for the unique radiation resistance of *D. radiodurans*?
45. What accounts for the purple color in salt ponds inhabited by halophilic archaea?
46. What evidence supports the hypothesis that some archaea live on Mars?


### Critical Thinking

47. The cell shown is found in the human stomach and is now known to cause peptic ulcers. What is the name of this bacterium?
48. The microscopic growth pattern shown is characteristic of which genus of bacteria?
49. What is the connection between this methane bog and archaea?



## Chapter 5

# The Eukaryotes of Microbiology



**Figure 5.1** Malaria is a disease caused by a eukaryotic parasite transmitted to humans by mosquitos. Micrographs (left and center) show a sporozoite life stage, trophozoites, and a schizont in a blood smear. On the right is depicted a primary defense against mosquito-borne illnesses like malaria—mosquito netting. (credit left: modification of work by Ute Frevert; credit right: modification of work by Tjeerd Wiersma)

### Chapter Outline

- 5.1 Unicellular Eukaryotic Parasites
- 5.2 The Parasitic Helminths
- 5.3 Fungi
- 5.4 Algae
- 5.5 Lichens

## Introduction

Although bacteria and viruses account for a large number of the infectious diseases that afflict humans, many serious illnesses are caused by eukaryotic organisms. One example is malaria, which is caused by *Plasmodium*, a eukaryotic organism transmitted through mosquito bites. Malaria is a major cause of morbidity (illness) and mortality (death) that threatens 3.4 billion people worldwide.<sup>[1]</sup> In severe cases, organ failure and blood or metabolic abnormalities contribute to medical emergencies and sometimes death. Even after initial recovery, relapses may occur years later. In countries where malaria is endemic, the disease represents a major public health challenge that can place a tremendous strain on developing economies.

Worldwide, major efforts are underway to reduce malaria infections. Efforts include the distribution of insecticide-treated bed nets and the spraying of pesticides. Researchers are also making progress in their efforts to develop effective vaccines.<sup>[2]</sup> The President’s Malaria Initiative, started in 2005, supports prevention and treatment. The Bill and Melinda Gates Foundation has a large initiative to eliminate malaria. Despite these efforts, malaria continues to cause long-term morbidity (such as intellectual disabilities in children) and mortality (especially in children younger than 5 years), so we still have far to go.

1. Centers for Disease Control and Prevention. “Impact of Malaria.” September 22, 2015. [http://www.cdc.gov/malaria/malaria\\_worldwide/impact.html](http://www.cdc.gov/malaria/malaria_worldwide/impact.html). Accessed January 18, 2016.

2. RTS, S Clinical Trials Partnership. “Efficacy and safety of RTS,S/AS01 malaria vaccine with or without a booster dose in infants and children in Africa: final results of a phase 3, individually randomised, controlled trial.” *The Lancet* 23 April 2015. DOI: [http://dx.doi.org/10.1016/S0140-6736\(15\)60721-8](http://dx.doi.org/10.1016/S0140-6736(15)60721-8).

## 5.1 Unicellular Eukaryotic Parasites

### Learning Objectives

- Summarize the general characteristics of unicellular eukaryotic parasites
- Describe the general life cycles and modes of reproduction in unicellular eukaryotic parasites
- Identify challenges associated with classifying unicellular eukaryotes
- Explain the taxonomic scheme used for unicellular eukaryotes
- Give examples of infections caused by unicellular eukaryotes

Eukaryotic microbes are an extraordinarily diverse group, including species with a wide range of life cycles, morphological specializations, and nutritional needs. Although more diseases are caused by viruses and bacteria than by microscopic eukaryotes, these eukaryotes are responsible for some diseases of great public health importance. For example, the protozoal disease malaria was responsible for 584,000 deaths worldwide (primarily children in Africa) in 2013, according to the World Health Organization (WHO). The protist parasite *Giardia* causes a diarrheal illness (giardiasis) that is easily transmitted through contaminated water supplies. In the United States, *Giardia* is the most common human intestinal parasite (**Figure 5.3**). Although it may seem surprising, parasitic worms are included within the study of microbiology because identification depends on observation of microscopic adult worms or eggs. Even in developed countries, these worms are important parasites of humans and of domestic animals. There are fewer fungal pathogens, but these are important causes of illness, as well. On the other hand, fungi have been

### Clinical Focus

#### Part 1

Upon arriving home from school, 7-year-old Sarah complains that a large spot on her arm will not stop itching. She keeps scratching at it, drawing the attention of her parents. Looking more closely, they see that it is a red circular spot with a raised red edge (**Figure 5.2**). The next day, Sarah's parents take her to their doctor, who examines the spot using a Woods lamp. A Woods lamp produces ultraviolet light that causes the spot on Sarah's arm to fluoresce, which confirms what the doctor already suspected: Sarah has a case of ringworm.

Sarah's mother is mortified to hear that her daughter has a "worm." How could this happen?




**Figure 5.2** Ringworm presents as a raised, red ring on the skin.

- What are some likely ways that Sarah might have contracted ringworm?

Jump to the **next** Clinical Focus box.



important in producing antimicrobial substances such as penicillin. In this chapter, we will examine characteristics of protists, worms, and fungi while considering their roles in causing disease.




**Figure 5.3** (a) A scanning electron micrograph shows many *Giardia* parasites in the trophozoite, or feeding stage, in a gerbil intestine. (b) An individual trophozoite of *G. lamblia*, visualized here in a scanning electron micrograph. This waterborne protist causes severe diarrhea when ingested.

## Characteristics of Protists

The word *protist* is a historical term that is now used informally to refer to a diverse group of microscopic eukaryotic organisms. It is not considered a formal taxonomic term because the organisms it describes do not have a shared evolutionary origin. Historically, the protists were informally grouped into the “animal-like” protozoans, the “plant-like” algae, and the “fungus-like” protists such as water molds. These three groups of protists differ greatly in terms of their basic characteristics. For example, algae are photosynthetic organisms that can be unicellular or multicellular. Protozoa, on the other hand, are nonphotosynthetic, motile organisms that are always unicellular. Other informal terms may also be used to describe various groups of protists. For example, microorganisms that drift or float in water, moved by currents, are referred to as **plankton**. Types of plankton include **zooplankton**, which are motile and nonphotosynthetic, and **phytoplankton**, which are photosynthetic.

Protozoans inhabit a wide variety of habitats, both aquatic and terrestrial. Many are free-living, while others are parasitic, carrying out a life cycle within a host or hosts and potentially causing illness. There are also beneficial symbionts that provide metabolic services to their hosts. During the feeding and growth part of their life cycle, they are called **trophozoites**; these feed on small particulate food sources such as bacteria. While some types of protozoa exist exclusively in the trophozoite form, others can develop from trophozoite to an encapsulated cyst stage when environmental conditions are too harsh for the trophozoite. A **cyst** is a cell with a protective wall, and the process by which a trophozoite becomes a cyst is called **encystment**. When conditions become more favorable, these cysts are triggered by environmental cues to become active again through **excystment**.

One protozoan genus capable of encystment is *Eimeria*, which includes some human and animal pathogens. **Figure 5.4** illustrates the life cycle of *Eimeria*.



**Figure 5.4** In the sexual/asexual life cycle of *Eimeria*, oocysts (inset) are shed in feces and may cause disease when ingested by a new host.

Protozoans have a variety of reproductive mechanisms. Some protozoans reproduce asexually and others reproduce sexually; still others are capable of both sexual and asexual reproduction. In protozoans, asexual reproduction occurs by binary fission, budding, or schizogony. In **schizogony**, the nucleus of a cell divides multiple times before the cell divides into many smaller cells. The products of schizogony are called merozoites and they are stored in structures known as schizonts. Protozoans may also reproduce sexually, which increases genetic diversity and can lead to complex life cycles. Protozoans can produce haploid gametes that fuse through **syngamy**. However, they can also exchange genetic material by joining to exchange DNA in a process called conjugation. This is a different process than the conjugation that occurs in bacteria. The term protist conjugation refers to a true form of eukaryotic sexual reproduction between two cells of different mating types. It is found in **ciliates**, a group of protozoans, and is described later in this subsection.


All protozoans have a plasma membrane, or **plasmalemma**, and some have bands of protein just inside the membrane that add rigidity, forming a structure called the **pellicle**. Some protists, including protozoans, have distinct layers of cytoplasm under the membrane. In these protists, the outer gel layer (with microfilaments of actin) is called the **ectoplasm**. Inside this layer is a sol (fluid) region of cytoplasm called the **endoplasm**. These structures contribute to complex cell shapes in some protozoans, whereas others (such as amoebas) have more flexible shapes (**Figure 5.5**).

Different groups of protozoans have specialized feeding structures. They may have a specialized structure for taking in food through phagocytosis, called a **cytostome**, and a specialized structure for the exocytosis of wastes called a **cytoproct**. Oral grooves leading to cytosomes are lined with hair-like cilia to sweep in food particles. Protozoans

are heterotrophic. Protozoans that are **holozoic** ingest whole food particles through phagocytosis. Forms that are **saprozoic** ingest small, soluble food molecules.

Many protists have whip-like flagella or hair-like cilia made of microtubules that can be used for locomotion (**Figure 5.5**). Other protists use cytoplasmic extensions known as pseudopodia (“false feet”) to attach the cell to a surface; they then allow cytoplasm to flow into the extension, thus moving themselves forward.

Protozoans have a variety of unique organelles and sometimes lack organelles found in other cells. Some have **contractile vacuoles**, organelles that can be used to move water out of the cell for osmotic regulation (salt and water balance) (**Figure 5.5**). Mitochondria may be absent (as in *Giardia*) in parasites or altered to kinetoplasts (modified mitochondria) or hydrogenosomes (see **Unique Characteristics of Prokaryotic Cells** for more discussion of these structures).



**Figure 5.5** (a) *Paramecium* spp. have hair-like appendages called cilia for locomotion. (b) *Amoeba* spp. use lobe-like pseudopodia to anchor the cell to a solid surface and pull forward. (c) *Euglena* spp. use a whip-like structure called a flagellum to propel the cell.




### Check Your Understanding

What is the sequence of events in reproduction by schizogony and what are the cells produced called?

## Taxonomy of Protists

The protists are a **polyphyletic** group, meaning they lack a shared evolutionary origin. Since the current taxonomy is based on evolutionary history (as determined by biochemistry, morphology, and genetics), protists are scattered across many different taxonomic groups within the domain Eukarya. Eukarya is currently divided into six supergroups that are further divided into subgroups, as illustrated in (**Figure 5.6**). In this section, we will primarily be concerned with the supergroups Amoebozoa, Excavata, and Chromalveolata; these supergroups include many protozoans of clinical significance. The supergroups Opisthokonta and Rhizaria also include some protozoans, but few of clinical significance. In addition to protozoans, Opisthokonta also includes animals and fungi, some of which we will discuss in **Parasitic Helminths** and **Fungi**. Some examples of the Archaeplastida will be discussed in **Algae**. **Figure 5.7** and **Figure 5.8** summarize the characteristics of each supergroup and subgroup and list representatives of each.



**Figure 5.6** This tree shows a proposed classification of the domain Eukarya based on evolutionary relationships. Currently, the domain Eukarya is divided into six supergroups. Within each supergroup are multiple kingdoms. Dotted lines indicate suggested evolutionary relationships that remain under debate.



The Eukaryote Supergroups and Some Examples				
Supergroup	Subgroups	Distinguishing Features	Examples	Clinical Notes
Excavata	Fornicata	Form cysts Pair of equal nuclei No mitochondria Often parasitic Four free flagella	<i>Giardia lamblia</i>	Giardiasis
	Parabasalids	No mitochondria Four free flagella One attached flagellum No cysts Parasitic or symbiotic Basal bodies Kinetoplastids	<i>Trichomonas</i>	Trichomoniasis
	Euglenozoans	Photosynthetic or heterotrophic Flagella	<i>Euglena</i>	N/a
			<i>Trypanosoma</i>	African sleeping sickness, Chagas disease
			<i>Leishmania</i>	Leishmaniasis
	Chromalveolata	Dinoflagellates	Cellulose theca Two dissimilar flagella	<i>Gonyaulax</i>
<i>Alexandrium</i>				Paralytic shellfish poisoning
<i>Pfiesteria</i>				Harmful algal blooms
Apicomplexans		Intracellular parasite Apical organelles	<i>Plasmodium</i>	Malaria
			<i>Cryptosporidium</i>	Cryptosporidiosis
			<i>Theileria (Babesia)</i>	Babesiosis
			<i>Toxoplasma</i>	Toxoplasmosis
Ciliates		Cilia	<i>Balantidium</i>	Balantidiasis
			<i>Paramecium</i>	N/a
			<i>Stentor</i>	N/a
Öomycetes/ peronosporomy- cetes		"Water molds" Generally diploid Cellulose cell walls	<i>Phytophthora</i>	Diseases in crops

Figure 5.7



The Eukaryote Supergroups and Some Examples (continued)				
Supergroup	Subgroups	Distinguishing Features	Examples	Clinical Notes
Rhizaria	Foraminifera	Amoeboid Threadlike pseudopodia Calcium carbonate shells	<i>Astrodonche</i>	N/a
	Radiolaria	Amoeboid Threadlike pseudopodia Silica shells	<i>Actinomma</i>	N/a
	Cercozoa	Amoeboid Threadlike pseudopodia Complex shells Parasitic forms	<i>Spongospora subterranea</i>	Powdery scab (potato disease)
<i>Plasmodiophora brassicae</i>			Cabbage clubroot	
Archaeplastida	Red algae	Chlorophyll <i>a</i> Phycocerythrin Phycocyanin Floridean starch Agar in cell walls	<i>Gelidium</i>	Source of agar
			<i>Gracilaria</i>	Source of agar
	Chlorophytes	Chlorophyll <i>a</i> Chlorophyll <i>b</i> Cellulose cell walls Starch storage	<i>Acetabularia</i>	N/a
			<i>Ulva</i>	N/a
Amoebozoa	Slime molds	Plasmodial and cellular forms	<i>Dictyostelium</i>	N/a
	Entamoebas	Trophozoites Form cysts	<i>Entamoeba</i>	Amoebiasis
			<i>Naegleria</i>	Primary amoebic meningoencephalitis
			<i>Acanthamoeba</i>	Keratitis, granulomatous amoebic encephalitis
Opisthokonta	Fungi	Chitin cell walls Unicellular or multicellular Often hyphae	Zygomycetes	Zygomycosis
			Ascomycetes	Candidiasis
			Basidiomycetes	Cryptococcosis
			Microsporidia	Microsporidiosis
	Animals	Multicellular heterotrophs No cell walls	Nematoda	Trichinosis; hookworm and pinworm infections
			Trematoda	Schistosomiasis
			Cestoda	Tapeworm infections

Figure 5.8



### Check Your Understanding

Which supergroups contain the clinically significant protists?

## Amoebozoa

The supergroup Amoebozoa includes protozoans that use amoeboid movement. Actin microfilaments produce pseudopodia, into which the remainder of the protoplasm flows, thereby moving the organism. The genus *Entamoeba* includes commensal or parasitic species, including the medically important *E. histolytica*, which is transmitted by cysts in feces and is the primary cause of amoebic dysentery. The notorious “brain-eating amoeba,” *Naegleria fowleri*, is also classified within the Amoebozoa. This deadly parasite is found in warm, fresh water and causes primary amoebic meningoencephalitis (PAM). Another member of this group is *Acanthamoeba*, which can cause keratitis (corneal inflammation) and blindness.

The Eumycetozoa are an unusual group of organisms called slime molds, which have previously been classified as animals, fungi, and plants (**Figure 5.9**). Slime molds can be divided into two types: cellular slime molds and plasmodial slime molds. The cellular slime molds exist as individual amoeboid cells that periodically aggregate into a mobile slug. The aggregate then forms a fruiting body that produces haploid spores. Plasmodial slime molds exist as large, multinucleate amoeboid cells that form reproductive stalks to produce spores that divide into gametes. One cellular slime mold, *Dictyostelium discoideum*, has been an important study organism for understanding cell differentiation, because it has both single-celled and multicelled life stages, with the cells showing some degree of differentiation in the multicelled form. **Figure 5.10** and **Figure 5.11** illustrate the life cycles of cellular and plasmodial slime molds, respectively.




(a)



(b)


**Figure 5.9** (a) The cellular slime mold *Dictyostelium discoideum* can be grown on agar in a Petri dish. In this image, individual amoeboid cells (visible as small spheres) are streaming together to form an aggregation that is beginning to rise in the upper right corner of the image. The primitively multicellular aggregation consists of individual cells that each have their own nucleus. (b) *Fuligo septica* is a plasmodial slime mold. This brightly colored organism consists of a large cell with many nuclei.

### Haploid and Asexual Reproduction



**Figure 5.10** The life cycle of the cellular slime mold *Dictyostelium discoideum* primarily involves individual amoebas but includes the formation of a multinucleate plasmodium formed from a uninucleate zygote (the result of the fusion of two individual amoeboid cells). The plasmodium is able to move and forms a fruiting body that generates haploid spores.






**Figure 5.11** Plasmodial slime molds exist as large multinucleate amoeboid cells that form reproductive stalks to produce spores that divide into gametes.

## Chromalveolata

The supergroup Chromalveolata is united by similar origins of its members' plastids and includes the apicomplexans, ciliates, diatoms, and dinoflagellates, among other groups (we will cover the diatoms and dinoflagellates with the algae in **Algae**). The apicomplexans are intra- or extracellular parasites that have an apical complex at one end of the cell. The apical complex is a concentration of organelles, vacuoles, and microtubules that allows the parasite to enter host cells (**Figure 5.12**). Apicomplexans have complex life cycles that include an infective sporozoite that undergoes schizogony to make many merozoites (see the example in **Figure 5.4**). Many are capable of infecting a variety of animal cells, from insects to livestock to humans, and their life cycles often depend on transmission between multiple hosts. The genus *Plasmodium* is an example of this group.



**Figure 5.12** (a) Apicomplexans are parasitic protists. They have a characteristic apical complex that enables them to infect host cells. (b) A colorized electron microscope image of a *Plasmodium* sporozoite. (credit b: modification of work by Ute Frevert)

Other apicomplexans are also medically important. *Cryptosporidium parvum* causes intestinal symptoms and can cause epidemic diarrhea when the cysts contaminate drinking water. *Theileria (Babesia) microti*, transmitted by the tick *Ixodes scapularis*, causes recurring fever that can be fatal and is becoming a common transfusion-transmitted pathogen in the United States (*Theileria* and *Babesia* are closely related genera and there is some debate about the best classification). Finally, *Toxoplasma gondii* causes toxoplasmosis and can be transmitted from cat feces, unwashed fruit and vegetables, or from undercooked meat. Because toxoplasmosis can be associated with serious birth defects, pregnant women need to be aware of this risk and use caution if they are exposed to the feces of potentially infected cats. A national survey found the frequency of individuals with antibodies for toxoplasmosis (and thus who presumably have a current latent infection) in the United States to be 11%. Rates are much higher in other countries, including some developed countries.<sup>[3]</sup> There is also evidence and a good deal of theorizing that the parasite may be responsible for altering infected humans' behavior and personality traits.<sup>[4]</sup>

The ciliates (Ciliophora), also within the Chromalveolata, are a large, very diverse group characterized by the presence of cilia on their cell surface. Although the cilia may be used for locomotion, they are often used for feeding, as well, and some forms are nonmotile. *Balantidium coli* (Figure 5.13) is the only parasitic ciliate that affects humans by causing intestinal illness, although it rarely causes serious medical issues except in the immunocompromised (those having a weakened immune system). Perhaps the most familiar ciliate is *Paramecium*, a motile organism with a clearly visible cytosome and cytoproct that is often studied in biology laboratories (Figure 5.14). Another ciliate, *Stentor*, is sessile and uses its cilia for feeding (Figure 5.15). Generally, these organisms have a **micronucleus** that is diploid, somatic, and used for sexual reproduction by conjugation. They also have a **macronucleus** that is derived from the micronucleus; the macronucleus becomes polyploid (multiple sets of duplicate chromosomes), and has a reduced set of metabolic genes.

Ciliates are able to reproduce through conjugation, in which two cells attach to each other. In each cell, the diploid micronuclei undergo meiosis, producing eight haploid nuclei each. Then, all but one of the haploid micronuclei and the macronucleus disintegrate; the remaining (haploid) micronucleus undergoes mitosis. The two cells then exchange one micronucleus each, which fuses with the remaining micronucleus present to form a new, genetically different,

3. J. Flegr et al. "Toxoplasmosis—A Global Threat. Correlation of Latent Toxoplasmosis With Specific Disease Burden in a Set of 88 Countries." *PLoS ONE* 9 no. 3 (2014):e90203.

4. J. Flegr. "Effects of Toxoplasma on Human Behavior." *Schizophrenia Bull* 33, no. 3 (2007):757–760.




diploid micronucleus. The diploid micronucleus undergoes two mitotic divisions, so each cell has four micronuclei, and two of the four combine to form a new macronucleus. The chromosomes in the macronucleus then replicate repeatedly, the macronucleus reaches its polyploid state, and the two cells separate. The two cells are now genetically different from each other and from their previous versions.


### Link to Learning




Protozoan conjugation is illustrated in this [animation](https://openstaxcollege.org/l/22proconjanim) (<https://openstaxcollege.org/l/22proconjanim>).



**Figure 5.13** This specimen of the ciliate *Balantidium coli* is a trophozoite form isolated from the gut of a primate. *B. coli* is the only ciliate capable of parasitizing humans (scale bar = 5  $\mu\text{m}$ ). (credit: modification of work by Kouassi RYW, McGraw SW, Yao PK, Abou-Bacar A, Brunet J, Pesson B, Bonfoh B, N'goran EK & Candolfi E)



**Figure 5.14** *Paramecium* has a primitive mouth (called an oral groove) to ingest food, and an anal pore to excrete it. Contractile vacuoles allow the organism to excrete excess water. Cilia enable the organism to move.



**Figure 5.15** This differential interference contrast micrograph (magnification:  $\times 65$ ) of *Stentor roeselie* shows cilia present on the margins of the structure surrounding the cytostome; the cilia move food particles.

Öomycetes have similarities to fungi and were once classified with them. They are also called water molds. However, they differ from fungi in several important ways. Öomycetes have cell walls of cellulose (unlike the chitinous cell walls of fungi) and they are generally diploid, whereas the dominant life forms of fungi are typically haploid. *Phytophthora*, the plant pathogen found in the soil that caused the Irish potato famine, is classified within this group (**Figure 5.16**).



**Figure 5.16** A saprobic oomycete, or water mold, engulfs a dead insect. (credit: modification of work by Thomas Bresson)

## Link to Learning



Explore the procedures for detecting the presence of an apicomplexan in a public water supply, at [this \(https://openstaxcollege.org//22detpreapicom\)](https://openstaxcollege.org//22detpreapicom) website.


This [video \(https://openstaxcollege.org//22feedstentor\)](https://openstaxcollege.org//22feedstentor) shows the feeding of *Stentor*.

## Excavata


The third and final supergroup to be considered in this section is the Excavata, which includes primitive eukaryotes and many parasites with limited metabolic abilities. These organisms have complex cell shapes and structures, often including a depression on the surface of the cell called an excavate. The group Excavata includes the subgroups Fornicata, Parabasalia, and Euglenozoa. The Fornicata lack mitochondria but have flagella. This group includes *Giardia lamblia* (also known as *G. intestinalis* or *G. duodenalis*), a widespread pathogen that causes diarrheal illness and can be spread through cysts from feces that contaminate water supplies (**Figure 5.3**). Parabasalia are frequent animal endosymbionts; they live in the guts of animals like termites and cockroaches. They have basal bodies and modified mitochondria (kinetoplastids). They also have a large, complex cell structure with an undulating membrane and often have many flagella. The trichomonads (a subgroup of the Parabasalia) include pathogens such as *Trichomonas vaginalis*, which causes the human sexually transmitted disease trichomoniasis. Trichomoniasis often does not cause symptoms in men, but men are able to transmit the infection. In women, it causes vaginal discomfort and discharge and may cause complications in pregnancy if left untreated.

The Euglenozoa are common in the environment and include photosynthetic and nonphotosynthetic species. Members of the genus *Euglena* are typically not pathogenic. Their cells have two flagella, a pellicle, a **stigma** (eyespot) to sense light, and chloroplasts for photosynthesis (**Figure 5.17**). The pellicle of *Euglena* is made of a series of protein bands surrounding the cell; it supports the cell membrane and gives the cell shape.

The Euglenozoa also include the trypanosomes, which are parasitic pathogens. The genus *Trypanosoma* includes *T. brucei*, which causes African trypanosomiasis (African sleeping sickness) and *T. cruzi*, which causes American trypanosomiasis (Chagas disease). These tropical diseases are spread by insect bites. In African sleeping sickness, *T. brucei* colonizes the blood and the brain after being transmitted via the bite of a tsetse fly (*Glossina* spp.) (**Figure 5.18**). The early symptoms include confusion, difficulty sleeping, and lack of coordination. Left untreated, it is fatal.



**Figure 5.17** (a) This illustration of a *Euglena* shows the characteristic structures, such as the stigma and flagellum. (b) The pellicle, under the cell membrane, gives the cell its distinctive shape and is visible in this image as delicate parallel striations over the surface of the entire cell (especially visible over the grey contractile vacuole). (credit a: modification of work by Claudio Miklos; credit b: modification of work by David Shykind)



**Figure 5.18** *Trypanosoma brucei*, the causative agent of African trypanosomiasis, spends part of its life cycle in the tsetse fly and part in humans.



Chagas' disease originated and is most common in Latin America. The disease is transmitted by a species of *Triatoma*, insects often called “kissing bugs,” and affects either the heart tissue or tissues of the digestive system. Untreated cases can eventually lead to heart failure or significant digestive or neurological disorders.

The genus *Leishmania*, yet another trypanosome, includes microbes that cause disfiguring skin disease and sometimes systemic illness as well.

## Eye on Ethics



### Neglected Parasites

The Centers for Disease Control and Prevention (CDC) is responsible for identifying public health priorities in the United States and developing strategies to address areas of concern. As part of this mandate, the CDC has officially identified five parasitic diseases it considers to have been neglected (i.e., not adequately studied). These neglected parasitic infections (NPIs) include toxoplasmosis, Chagas disease, toxicariasis (a nematode infection transmitted primarily by infected dogs), cysticercosis (a disease caused by a tissue infection of the tapeworm *Taenia solium*), and trichomoniasis (a sexually transmitted disease caused by the parabasalid *Trichomonas vaginalis*).

The decision to name these specific diseases as NPIs means that the CDC will devote resources toward improving awareness and developing better diagnostic testing and treatment through studies of available data. The CDC may also advise on treatment of these diseases and assist in the distribution of medications that might otherwise be difficult to obtain.<sup>[5]</sup>

Of course, the CDC does not have unlimited resources, so by prioritizing these five diseases, it is effectively deprioritizing others. Given that many Americans have never heard of many of these NPIs, it is fair to ask what criteria the CDC used in prioritizing diseases. According to the CDC, the factors considered were the number of people infected, the severity of the illness, and whether the illness can be treated or prevented. Although several of these NPIs may seem to be more common outside the United States, the CDC argues that many cases in the United States likely go undiagnosed and untreated because so little is known about these diseases.<sup>[6]</sup>

What criteria should be considered when prioritizing diseases for purposes of funding or research? Are those identified by the CDC reasonable? What other factors could be considered? Should government agencies like the CDC have the same criteria as private pharmaceutical research labs? What are the ethical implications of deprioritizing other potentially neglected parasitic diseases such as leishmaniasis?

5. Centers for Disease Control and Prevention. “Neglected Parasitic Infections (NPIs) in the United States.” <http://www.cdc.gov/parasites/npi/>. Last updated July 10, 2014.

6. Centers for Disease Control and Prevention. “Fact Sheet: Neglected Parasitic Infections in the United States.” [http://www.cdc.gov/parasites/resources/pdf/npi\\_factsheet.pdf](http://www.cdc.gov/parasites/resources/pdf/npi_factsheet.pdf)



## 5.2 The Parasitic Helminths


### Learning Objectives

- Explain why we include the study of parasitic worms within the discipline of microbiology
- Compare the basic morphology of the major groups of parasitic helminths
- Describe the characteristics of parasitic nematodes, and give an example of infective eggs and infective larvae
- Describe the characteristics of parasitic trematodes and cestodes, and give examples of each
- Identify examples of the primary causes of infections due to nematodes, trematodes, and cestodes
- Classify parasitic worms according to major groups

Parasitic helminths are animals that are often included within the study of microbiology because many species of these worms are identified by their microscopic eggs and larvae. There are two major groups of parasitic helminths: the roundworms (Nematoda) and flatworms (Platyhelminthes). Of the many species that exist in these groups, about half are parasitic and some are important human pathogens. As animals, they are multicellular and have organ systems. However, the parasitic species often have limited digestive tracts, nervous systems, and locomotor abilities. Parasitic forms may have complex reproductive cycles with several different life stages and more than one type of host. Some are **monoecious**, having both male and female reproductive organs in a single individual, while others are **dioecious**, each having either male or female reproductive organs.

### Nematoda (Roundworms)

Phylum **Nematoda** (the roundworms) is a diverse group containing more than 15,000 species, of which several are important human parasites (**Figure 5.19**). These unsegmented worms have a full digestive system even when parasitic. Some are common intestinal parasites, and their eggs can sometimes be identified in feces or around the anus of infected individuals. *Ascaris lumbricoides* is the largest nematode intestinal parasite found in humans; females may reach lengths greater than 1 meter. *A. lumbricoides* is also very widespread, even in developed nations, although it is now a relatively uncommon problem in the United States. It may cause symptoms ranging from relatively mild (such as a cough and mild abdominal pain) to severe (such as intestinal blockage and impaired growth).



**Figure 5.19** A micrograph of the nematode *Enterobius vermicularis*, also known as the pinworm.

Of all nematode infections in the United States, pinworm (caused by *Enterobius vermicularis*) is the most common. Pinworm causes sleeplessness and itching around the anus, where the female worms lay their eggs during the night. *Toxocara canis* and *T. cati* are nematodes found in dogs and cats, respectively, that can be transmitted to humans,

causing toxocariasis. Antibodies to these parasites have been found in approximately 13.9% of the U.S. population, suggesting that exposure is common.<sup>[7]</sup> Infection can cause larval migrans, which can result in vision loss and eye inflammation, or fever, fatigue, coughing, and abdominal pain, depending on whether the organism infects the eye or the viscera. Another common nematode infection is hookworm, which is caused by *Necator americanus* (the New World or North American hookworm) and *Ancylostoma duodenale* (the Old World hookworm). Symptoms of hookworm infection can include abdominal pain, diarrhea, loss of appetite, weight loss, fatigue, and anemia.

Trichinellosis, also called trichinosis, caused by *Trichinella spiralis*, is contracted by consuming undercooked meat, which releases the larvae and allows them to encyst in muscles. Infection can cause fever, muscle pains, and digestive system problems; severe infections can lead to lack of coordination, breathing and heart problems, and even death. Finally, heartworm in dogs and other animals is caused by the nematode *Dirofilaria immitis*, which is transmitted by mosquitoes. Symptoms include fatigue and cough; when left untreated, death may result.

## Clinical Focus

### Part 2

The physician explains to Sarah's mother that ringworm can be transferred between people through touch. "It's common in school children, because they often come in close contact with each other, but anyone can become infected," he adds. "Because you can transfer it through objects, locker rooms and public pools are also a potential source of infection. It's very common among wrestlers and athletes in other contact sports."

Looking very uncomfortable, Sarah says to her mother "I want this worm out of me."

The doctor laughs and says, "Sarah, you're in luck because ringworm is just a name; it is not an actual worm. You have nothing wriggling around under your skin."

"Then what is it?" asks Sarah.

- What type of pathogen causes ringworm?

Jump to the **next** Clinical Focus box. Go back to the **previous** Clinical Focus box.



## Check Your Understanding

What is the most common nematode infection in the United States?

## Platyhelminths (Flatworms)

Phylum **Platyhelminthes** (the platyhelminths) are flatworms. This group includes the flukes, tapeworms, and the turbellarians, which include planarians. The flukes and tapeworms are medically important parasites (**Figure 5.20**).


The **flukes** (trematodes) are nonsegmented flatworms that have an oral sucker (**Figure 5.21**) (and sometimes a second ventral sucker) and attach to the inner walls of intestines, lungs, large blood vessels, or the liver. Trematodes have complex life cycles, often with multiple hosts. Several important examples are the liver flukes (*Clonorchis* and *Opisthorchis*), the intestinal fluke (*Fasciolopsis buski*), and the oriental lung fluke (*Paragonimus westermani*). Schistosomiasis is a serious parasitic disease, considered second in the scale of its impact on human populations only to malaria. The parasites *Schistosoma mansoni*, *S. haematobium*, and *S. japonicum*, which are found in freshwater snails, are responsible for schistosomiasis (**Figure 5.22**). Immature forms burrow through the skin into the blood.

7. Won K, Kruszon-Moran D, Schantz P, Jones J. "National seroprevalence and risk factors for zoonotic Toxocara spp. infection." In: Abstracts of the 56th American Society of Tropical Medicine and Hygiene; Philadelphia, Pennsylvania; 2007 Nov 4-8.


They migrate to the lungs, then to the liver and, later, other organs. Symptoms include anemia, malnutrition, fever, abdominal pain, fluid buildup, and sometimes death.



(a) Class Turbellaria



(b) Class Monogenea




(c) Class Trematoda




(d) Class Cestoda

**Figure 5.20** Phylum Platyhelminthes is divided into four classes. (a) Class Turbellaria includes the Bedford's flatworm (*Pseudobiceros bedfordi*), which is about 8–10 cm long. (b) The parasitic class Monogenea includes *Dactylogyrus* spp. Worms in this genus are commonly called gill flukes. The specimen pictured here is about 0.2 mm long and has two anchors, indicated by arrows, that it uses to latch onto the gills of host fish. (c) The Trematoda class includes the common liver fluke *Fasciola hepatica* and the giant liver fluke *Fascioloides magna* (right). The *F. magna* specimen shown here is about 7 cm long. (d) Class Cestoda includes tapeworms such as *Taenia saginata*, which infects both cattle and humans and can reach lengths of 4–10 meters; the specimen shown here is about 4 meters long.



**Figure 5.21** (a) The oral sucker is visible on the anterior end of this liver fluke, *Opisthorchis viverrini*. (b) This micrograph shows the scolex of the cestode *Taenia solium*, also known as the pork tapeworm. The visible suckers and hooks allow the worm to attach itself to the inner wall of the intestine. (credit a: modification of work by Sripa B, Kaewkes S, Sithithaworn P, Mairiang E, Laha T, and Smout M)






**Figure 5.22** The life cycle of *Schistosoma* spp. includes several species of water snails, which serve as secondary hosts. The parasite is transmitted to humans through contact with contaminated water and takes up residence in the veins of the digestive system. Eggs escape the host in the urine or feces and infect a snail to complete the life cycle.

The other medically important group of platyhelminths are commonly known as **tapeworms** (cestodes) and are segmented flatworms that may have suckers or hooks at the **scolex** (head region) (**Figure 5.21**). Tapeworms use these suckers or hooks to attach to the wall of the small intestine. The body of the worm is made up of segments called **proglottids** that contain reproductive structures; these detach when the gametes are fertilized, releasing gravid proglottids with eggs. Tapeworms often have an intermediate host that consumes the eggs, which then hatch into a larval form called an oncosphere. The oncosphere migrates to a particular tissue or organ in the intermediate host, where it forms cysticerci. After being eaten by the definitive host, the cysticerci develop into adult tapeworms in the host's digestive system (**Figure 5.23**). *Taenia saginata* (the beef tapeworm) and *T. solium* (the pork tapeworm) enter



humans through ingestion of undercooked, contaminated meat. The adult worms develop and reside in the intestine, but the larval stage may migrate and be found in other body locations such as skeletal and smooth muscle. The beef tapeworm is relatively benign, although it can cause digestive problems and, occasionally, allergic reactions. The pork tapeworm can cause more serious problems when the larvae leave the intestine and colonize other tissues, including those of the central nervous system. *Diphyllobothrium latum* is the largest human tapeworm and can be ingested in undercooked fish. It can grow to a length of 15 meters. *Echinococcus granulosus*, the dog tapeworm, can parasitize humans and uses dogs as an important host.



**Figure 5.23** Life cycle of a tapeworm.



## Check Your Understanding

What group of medically important flatworms is segmented and what group is unsegmented?

## Micro Connections

### Food for Worms?

For residents of temperate, developed countries, it may be difficult to imagine just how common helminth infections are in the human population. In fact, they are quite common and even occur frequently in the United States. Worldwide, approximately 807–1,221 million people are infected with *Ascaris lumbricoides* (perhaps one-sixth of the human population) and far more are infected if all nematode species are considered.<sup>[8]</sup> Rates of infection are relatively high even in industrialized nations. Approximately 604–795 million people are infected with whipworm (*Trichuris*) worldwide (*Trichuris* can also infect dogs), and 576–740 million people are infected with hookworm (*Necatur americanus* and *Ancylostoma duodenale*).<sup>[9]</sup> *Toxocara*, a nematode parasite of dogs and cats, is also able to infect humans. It is widespread in the United States, with about 10,000 symptomatic cases annually. However, one study found 14% of the population (more than 40 million Americans) was seropositive, meaning they had been exposed to the parasite at one time. More than 200 million people have schistosomiasis worldwide. Most of the World Health Organization (WHO) neglected tropical diseases are helminths. In some cases, helminths may cause subclinical illnesses, meaning the symptoms are so mild that that they go unnoticed. In other cases, the effects may be more severe or chronic, leading to fluid accumulation and organ damage. With so many people affected, these parasites constitute a major global public health concern.

## Micro Connections

### Eradicating the Guinea Worm

Dracunculiasis, or Guinea worm disease, is caused by a nematode called *Dracunculus medinensis*. When people consume contaminated water, water fleas (small crustaceans) containing the nematode larvae may be ingested. These larvae migrate out of the intestine, mate, and move through the body until females eventually emerge (generally through the feet). While Guinea worm disease is rarely fatal, it is extremely painful and can be accompanied by secondary infections and edema (Figure 5.24).



**Figure 5.24** The Guinea worm can be removed from a leg vein of an infected person by gradually winding it around a stick, like this matchstick.

An eradication campaign led by WHO, the CDC, the United Nations Children's Fund (UNICEF), and the Carter Center (founded by former U.S. president Jimmy Carter) has been extremely successful in reducing cases of dracunculiasis. This has been possible because diagnosis is straightforward, there is an inexpensive method of control, there is no animal reservoir, the water fleas are not airborne (they are restricted to still water),

8. Fenwick, A. "The global burden of neglected tropical diseases." *Public health* 126 no.3 (Mar 2012): 233–6.

9. de Silva, N., et al. (2003). "Soil-transmitted helminth infections: updating the global picture". *Trends in Parasitology* 19 (December 2003): 547–51.

the disease is geographically limited, and there has been a commitment from the governments involved. Additionally, no vaccines or medication are required for treatment and prevention. In 1986, 3.5 million people were estimated to be affected. After the eradication campaign, which included helping people in affected areas learn to filter water with cloth, only four countries continue to report the disease (Chad, Mali, South Sudan, and Ethiopia) with a total of 126 cases reported to WHO in 2014.<sup>[10]</sup>

## 5.3 Fungi

### Learning Objectives

- Explain why the study of fungi such as yeast and molds is within the discipline of microbiology
- Describe the unique characteristics of fungi
- Describe examples of asexual and sexual reproduction of fungi
- Compare the major groups of fungi in this chapter, and give examples of each
- Identify examples of the primary causes of infections due to yeasts and molds
- Identify examples of toxin-producing fungi
- Classify fungal organisms according to major groups


The fungi comprise a diverse group of organisms that are heterotrophic and typically saprozoic. In addition to the well-known macroscopic fungi (such as mushrooms and molds), many unicellular yeasts and spores of macroscopic fungi are microscopic. For this reason, fungi are included within the field of microbiology.

Fungi are important to humans in a variety of ways. Both microscopic and macroscopic fungi have medical relevance, with some pathogenic species that can cause **mycoses** (illnesses caused by fungi). Some pathogenic fungi are opportunistic, meaning that they mainly cause infections when the host's immune defenses are compromised and do not normally cause illness in healthy individuals. Fungi are important in other ways. They act as decomposers in the environment, and they are critical for the production of certain foods such as cheeses. Fungi are also major sources of antibiotics, such as penicillin from the fungus *Penicillium*.

### Characteristics of Fungi

Fungi have well-defined characteristics that set them apart from other organisms. Most multicellular fungal bodies, commonly called molds, are made up of filaments called **hyphae**. Hyphae can form a tangled network called a **mycelium** and form the **thallus** (body) of fleshy fungi. Hyphae that have walls between the cells are called **septate hyphae**; hyphae that lack walls and cell membranes between the cells are called nonseptate or **coenocytic hyphae**. (Figure 5.25).

10. World Health Organization. "South Sudan Reports Zero Cases of Guinea-Worm Disease for Seventh Consecutive Month." 2016. [http://www.who.int/dracunculiasis/no\\_new\\_case\\_for\\_seventh\\_consecutive\\_months/en/](http://www.who.int/dracunculiasis/no_new_case_for_seventh_consecutive_months/en/). Accessed May 2, 2016.




**Figure 5.25** Multicellular fungi (molds) form hyphae, which may be septate or nonseptate. Unicellular fungi (yeasts) cells form pseudohyphae from individual yeast cells.

In contrast to molds, yeasts are unicellular fungi. The **budding yeasts** reproduce asexually by budding off a smaller daughter cell; the resulting cells may sometimes stick together as a short chain or **pseudohypha** (**Figure 5.25**). *Candida albicans* is a common yeast that forms pseudohyphae; it is associated with various infections in humans, including vaginal yeast infections, oral thrush, and candidiasis of the skin.

Some fungi are dimorphic, having more than one appearance during their life cycle. These **dimorphic fungi** may be able to appear as yeasts or molds, which can be important for infectivity. They are capable of changing their appearance in response to environmental changes such as nutrient availability or fluctuations in temperature, growing as a mold, for example, at 25 °C (77 °F), and as yeast cells at 37 °C (98.6 °F). This ability helps dimorphic fungi to survive in diverse environments. *Histoplasma capsulatum*, the pathogen that causes histoplasmosis, a lung infection, is an example of a dimorphic fungus (**Figure 5.26**).






**Figure 5.26** *Histoplasma capsulatum* is a dimorphic fungus that grows in soil exposed to bird feces or bat feces (guano) (top left). It can change forms to survive at different temperatures. In the outdoors, it typically grows as a mycelium (as shown in the micrograph, bottom left), but when the spores are inhaled (right), it responds to the high internal temperature of the body (37 °C [98.6 °F]) by turning into a yeast that can multiply in the lungs, causing the chronic lung disease histoplasmosis.

There are notable unique features in fungal cell walls and membranes. Fungal cell walls contain **chitin**, as opposed to the cellulose found in the cell walls of plants and many protists. Additionally, whereas animals have cholesterol in their cell membranes, fungal cell membranes have different sterols called ergosterols. Ergosterols are often exploited as targets for antifungal drugs.

Fungal life cycles are unique and complex. Fungi reproduce sexually either through cross- or self-fertilization. Haploid fungi form hyphae that have gametes at the tips. Two different mating types (represented as “+ type” and “– type”) are involved. The cytoplasms of the + and – type gametes fuse (in an event called plasmogamy), producing a cell with two distinct nuclei (a **dikaryotic** cell). Later, the nuclei fuse (in an event called karyogamy) to create a diploid zygote. The zygote undergoes meiosis to form **spores** that germinate to start the haploid stage, which eventually creates more haploid mycelia (**Figure 5.27**). Depending on the taxonomic group, these sexually produced spores are known as zygospores (in Zygomycota), ascospores (in Ascomycota), or basidiospores (in Basidiomycota) (**Figure 5.28**).





**Figure 5.27** Zygomycetes have sexual and asexual life cycles. In the sexual life cycle, + and - mating types conjugate to form a zygosporangium.

Fungi may also exhibit asexual reproduction by mitosis, mitosis with budding, fragmentation of hyphae, and formation of asexual spores by mitosis. These spores are specialized cells that, depending on the organism, may have unique characteristics for survival, reproduction, and dispersal. Fungi exhibit several types of asexual spores and these can be important in classification (**Figure 5.28**).



**Figure 5.28** These images show asexually produced spores. (a) This brightfield micrograph shows the release of spores from a sporangium at the end of a hypha called a sporangiophore. The organism is a *Mucor* sp. fungus, a mold often found indoors. (b) Sporangia grow at the ends of stalks, which appear as the white fuzz seen on this bread mold, *Rhizopus stolonifer*. The tips of bread mold are the dark, spore-containing sporangia. (credit “spores of bread mold”: “Andrew”/Flickr)







### Check Your Understanding

Is a dimorphic fungus a yeast or a mold? Explain.

## Fungal Diversity

The fungi are very diverse, comprising seven major groups. Not all of the seven groups contain pathogens. Some of these groups are generally associated with plants and include plant pathogens. For example, Urediniomycetes and Ustilagomycetes include the plant rusts and smuts, respectively. These form reddish or dark masses, respectively, on plants as rusts (red) or smuts (dark). Some species have substantial economic impact because of their ability to reduce crop yields. Glomeromycota includes the mycorrhizal fungi, important symbionts with plant roots that can promote plant growth by acting like an extended root system. The Glomeromycota are obligate symbionts, meaning that they can only survive when associated with plant roots; the fungi receive carbohydrates from the plant and the plant benefits from the increased ability to take up nutrients and minerals from the soil. The Chytridiomycetes (chytrids) are small fungi, but are extremely ecologically important. Chytrids are generally aquatic and have flagellated, motile gametes; specific types are implicated in amphibian declines around the world. Because of their medical importance, we will focus on Zygomycota, Ascomycota, Basidiomycota, and Microsporidia (**Figure 5.29**).

Select Groups of Fungi				
Group	Characteristics	Examples	Medically Important Species	Image
Ascomycota	Septate hyphae Ascus with ascospores in ascocarp Conidiospores	Cup fungi Edible mushrooms Morels Truffles <i>Neurospora</i> <i>Penicillium</i>	<i>Aspergillus</i> spp. <i>Trichophyton</i> spp. <i>Microsporum</i> spp. <i>Epidermophyton</i> spp. <i>Blastomyces dermatitidis</i> <i>Histoplasma capsulatum</i>	 <i>Aspergillus niger</i>
Basidiomycota	Basidia produce basidiospores in a basidiocarp	Club fungi Rusts Stinkhorns Puffballs Mushrooms <i>Cryptococcus neoformans</i> <i>Amanita phalloides</i>	<i>Cryptococcus neoformans</i>	 <i>Amanita phalloides</i>
Microsporidia	Lack mitochondria, peroxisomes, centrioles Spores produce a polar tube	<i>Enterocystozoan bieneusi</i>	<i>Enterocystozoan bieneusi</i>	 Microsporidia (unidentified)
Zygomycota	Mainly saprophytes Coenocytic hyphae Haploid nuclei Zygospores	<i>Rhizopus stolonifera</i>	<i>Mucor</i> spp.	 <i>Rhizopus</i> sp.


**Figure 5.29**

The Zygomycota (zygomycetes) are mainly saprophytes with coenocytic hyphae and haploid nuclei. They use sporangiospores for asexual reproduction. The group name comes from the **zygospores** that they use for sexual reproduction (**Figure 5.27**), which have hard walls formed from the fusion of reproductive cells from two individuals. Zygomycetes are important for food science and as crop pathogens. One example is *Rhizopus stolonifera* (**Figure 5.28**), an important bread mold that also causes rice seedling blight. *Mucor* is a genus of fungi that can

potentially cause necrotizing infections in humans, although most species are intolerant of temperatures found in mammalian bodies (**Figure 5.28**).

The Ascomycota include fungi that are used as food (edible mushrooms, morels, and truffles), others that are common causes of food spoilage (bread molds and plant pathogens), and still others that are human pathogens. Ascomycota may have septate hyphae and cup-shaped fruiting bodies called **ascocarps**. Some genera of Ascomycota use sexually produced **ascospores** as well as asexual spores called **conidia**, but sexual phases have not been discovered or described for others. Some produce an **ascus** containing ascospores within an ascocarp (**Figure 5.31**).

Examples of the Ascomycota include several bread molds and minor pathogens, as well as species capable of causing more serious mycoses. Species in the genus *Aspergillus* are important causes of allergy and infection, and are useful in research and in the production of certain fermented alcoholic beverages such as Japanese *sake*. The fungus *Aspergillus flavus*, a contaminant of nuts and stored grains, produces an **aflatoxin** that is both a toxin and the most potent known natural carcinogen. *Neurospora crassa* is of particular use in genetics research because the spores produced by meiosis are kept inside the ascus in a row that reflects the cell divisions that produced them, giving a direct view of segregation and assortment of genes (**Figure 5.30**). *Penicillium* produces the antibiotic penicillin (**Figure 5.31**).




**Figure 5.30** These ascospores, lined up within an ascus, are produced sexually.

Many species of ascomycetes are medically important. A large number of species in the genera *Trichophyton*, *Microsporum*, and *Epidermophyton* are dermatophytes, pathogenic fungi capable of causing skin infections such as athlete's foot, jock itch, and ringworm. *Blastomyces dermatitidis* is a dimorphic fungus that can cause blastomycosis, a respiratory infection that, if left untreated, can become disseminated to other body sites, sometimes leading to death. Another important respiratory pathogen is the dimorphic fungus *Histoplasma capsulatum* (**Figure 5.26**), which is associated with birds and bats in the Ohio and Mississippi river valleys. *Coccidioides immitis* causes the serious lung disease Valley fever. *Candida albicans*, the most common cause of vaginal and other yeast infections, is also an ascomycete fungus; it is a part of the normal microbiota of the skin, intestine, genital tract, and ear (**Figure 5.31**). Ascomycetes also cause plant diseases, including ergot infections, Dutch elm disease, and powdery mildews.


*Saccharomyces* yeasts, including the baker's yeast *S. cerevisiae*, are unicellular ascomycetes with haploid and diploid stages (**Figure 5.32**). This and other *Saccharomyces* species are used for brewing beer.






**Figure 5.31** (a) This brightfield micrograph shows ascospores being released from asci in the fungus *Talaromyces flavus* var. *flavus*. (b) This electron micrograph shows the conidia (spores) borne on the conidiophore of *Aspergillus*, a type of toxic fungus found mostly in soil and plants. (c) This brightfield micrograph shows the yeast *Candida albicans*, the causative agent of candidiasis and thrush.





**Figure 5.32** The life cycle of an ascomycete is characterized by the production of asci during the sexual phase. The haploid phase is the predominant phase of the life cycle.

The Basidiomycota (basidiomycetes) are fungi that have **basidia** (club-shaped structures) that produce **basidiospores** (spores produced through budding) within fruiting bodies called **basidiocarps** (Figure 5.33). They are important as decomposers and as food. This group includes rusts, stinkhorns, puffballs, and mushrooms. Several species are of particular importance. *Cryptococcus neoformans*, a fungus commonly found as a yeast in the environment, can cause serious lung infections when inhaled by individuals with weakened immune systems. The edible meadow mushroom, *Agaricus campestris*, is a basidiomycete, as is the poisonous mushroom *Amanita phalloides*, known as the death cap. The deadly toxins produced by *A. phalloides* have been used to study transcription.



**Figure 5.33** The life cycle of a basidiomycete alternates a haploid generation with a prolonged stage in which two nuclei (dikaryon) are present in the hyphae.

Finally, the **Microsporidia** are unicellular fungi that are obligate intracellular parasites. They lack mitochondria, peroxisomes, and centrioles, but their spores release a unique **polar tubule** that pierces the host cell membrane to allow the fungus to gain entry into the cell. A number of microsporidia are human pathogens, and infections with microsporidia are called microsporidiosis. One pathogenic species is *Enterocystozoan bieneusi*, which can cause symptoms such as diarrhea, cholecystitis (inflammation of the gall bladder), and in rare cases, respiratory illness.



### Check Your Understanding

Which group of fungi appears to be associated with the greatest number of human diseases?

## Micro Connections

### Eukaryotic Pathogens in Eukaryotic Hosts

When we think about antimicrobial medications, antibiotics such as penicillin often come to mind. Penicillin and related antibiotics interfere with the synthesis of peptidoglycan cell walls, which effectively targets bacterial cells. These antibiotics are useful because humans (like all eukaryotes) do not have peptidoglycan cell walls.

Developing medications that are effective against eukaryotic cells but not harmful to human cells is more difficult. Despite huge morphological differences, the cells of humans, fungi, and protists are similar in terms of their ribosomes, cytoskeletons, and cell membranes. As a result, it is more challenging to develop medications that target protozoans and fungi in the same way that antibiotics target prokaryotes.

Fungicides have relatively limited modes of action. Because fungi have ergosterols (instead of cholesterol) in their cell membranes, the different enzymes involved in sterol production can be a target of some medications. The azole and morpholine fungicides interfere with the synthesis of membrane sterols. These are used widely in agriculture (fenpropimorph) and clinically (e.g., miconazole). Some antifungal medications target the chitin cell walls of fungi. Despite the success of these compounds in targeting fungi, antifungal medications for systemic infections still tend to have more toxic side effects than antibiotics for bacteria.

## Clinical Focus


### Part 3

Sarah is relieved the ringworm is not an actual worm, but wants to know what it really is. The physician explains that ringworm is a fungus. He tells her that she will not see mushrooms popping out of her skin, because this fungus is more like the invisible part of a mushroom that hides in the soil. He reassures her that they are going to get the fungus out of her too.

The doctor cleans and then carefully scrapes the lesion to place a specimen on a slide. By looking at it under a microscope, the physician is able to confirm that a fungal infection is responsible for Sarah's lesion. In **Figure 5.34**, it is possible to see macro- and microconidia in *Trichophyton rubrum*. Cell walls are also visible. Even if the pathogen resembled a helminth under the microscope, the presence of cell walls would rule out the possibility because animal cells lack cell walls.

The doctor prescribes an antifungal cream for Sarah's mother to apply to the ringworm. Sarah's mother asks, "What should we do if it doesn't go away?"

- Can all forms of ringworm be treated with the same antifungal medication?



**Figure 5.34** This micrograph shows hyphae (macroconidium) and microconidia of *Trichophyton rubrum*, a dermatophyte responsible for fungal infections of the skin.

Jump to the **next** Clinical Focus box. Go back to the **previous** Clinical Focus box.

## 5.4 Algae

### Learning Objectives

- Explain why algae are included within the discipline of microbiology
- Describe the unique characteristics of algae
- Identify examples of toxin-producing algae
- Compare the major groups of algae in this chapter, and give examples of each
- Classify algal organisms according to major groups

The **algae** are autotrophic protists that can be unicellular or multicellular. These organisms are found in the supergroups Chromalveolata (dinoflagellates, diatoms, golden algae, and brown algae) and Archaeplastida (red algae and green algae). They are important ecologically and environmentally because they are responsible for the production of approximately 70% of the oxygen and organic matter in aquatic environments. Some types of algae, even those that are microscopic, are regularly eaten by humans and other animals. Additionally, algae are the source for **agar**, agarose, and **carrageenan**, solidifying agents used in laboratories and in food production. Although algae are typically not pathogenic, some produce toxins. Harmful **algal blooms**, which occur when algae grow quickly and produce dense populations, can produce high concentrations of toxins that impair liver and nervous-system function in aquatic animals and humans.

Like protozoans, algae often have complex cell structures. For instance, algal cells can have one or more chloroplasts that contain structures called **pyrenoids** to synthesize and store starch. The chloroplasts themselves differ in their number of membranes, indicative of secondary or rare tertiary endosymbiotic events. Primary chloroplasts have two membranes—one from the original cyanobacteria that the ancestral eukaryotic cell engulfed, and one from the plasma membrane of the engulfing cell. Chloroplasts in some lineages appear to have resulted from secondary endosymbiosis, in which another cell engulfed a green or red algal cell that already had a primary chloroplast within it. The engulfing cell destroyed everything except the chloroplast and possibly the cell membrane of its original cell, leaving three or four membranes around the chloroplast. Different algal groups have different pigments, which are reflected in common names such as red algae, brown algae, and green algae.



Some algae, the seaweeds, are macroscopic and may be confused with plants. Seaweeds can be red, brown, or green, depending on their photosynthetic pigments. Green algae, in particular, share some important similarities with land plants; however, there are also important distinctions. For example, seaweeds do not have true tissues or organs like plants do. Additionally, seaweeds do not have a waxy cuticle to prevent desiccation. Algae can also be confused with cyanobacteria, photosynthetic bacteria that bear a resemblance to algae; however, cyanobacteria are prokaryotes (see **Nonproteobacteria Gram-negative Bacteria and Phototrophic Bacteria**).


Algae have a variety of life cycles. Reproduction may be asexual by mitosis or sexual using gametes.

## Algal Diversity

Although the algae and protozoa were formerly separated taxonomically, they are now mixed into supergroups. The algae are classified within the Chromalveolata and the Archaeplastida. Although the Euglenozoa (within the supergroup Excavata) include photosynthetic organisms, these are not considered algae because they feed and are motile.

The dinoflagellates and stramenopiles fall within the Chromalveolata. The **dinoflagellates** are mostly marine organisms and are an important component of plankton. They have a variety of nutritional types and may be phototrophic, heterotrophic, or mixotrophic. Those that are photosynthetic use chlorophyll *a*, chlorophyll *c*<sub>2</sub>, and other photosynthetic pigments (**Figure 5.35**). They generally have two flagella, causing them to whirl (in fact, the name dinoflagellate comes from the Greek word for “whirl”: *dini*). Some have cellulose plates forming a hard outer covering, or **theca**, as armor. Additionally, some dinoflagellates produce neurotoxins that can cause paralysis in humans or fish. Exposure can occur through contact with water containing the dinoflagellate toxins or by feeding on organisms that have eaten dinoflagellates.

When a population of dinoflagellates becomes particularly dense, a **red tide** (a type of harmful algal bloom) can occur. Red tides cause harm to marine life and to humans who consume contaminated marine life. Major toxin producers include *Gonyaulax* and *Alexandrium*, both of which cause paralytic shellfish poisoning. Another species, *Pfiesteria piscicida*, is known as a fish killer because, at certain parts of its life cycle, it can produce toxins harmful to fish and it appears to be responsible for a suite of symptoms, including memory loss and confusion, in humans exposed to water containing the species.




**Figure 5.35** The dinoflagellates exhibit great diversity in shape. Many are encased in cellulose armor and have two flagella that fit in grooves between the plates. Movement of these two perpendicular flagella causes a spinning motion. (credit: modification of work by CSIRO)

The **stramenopiles** include the golden algae (Chrysophyta), the brown algae (Phaeophyta), and the **diatoms** (Bacillariophyta). Stramenopiles have chlorophyll *a*, chlorophyll *c*<sub>1/c</sub><sub>2</sub>, and fucoxanthin as photosynthetic pigments. Their storage carbohydrate is chrysolaminarin. While some lack cell walls, others have scales. Diatoms have flagella and **frustules**, which are outer cell walls of crystallized silica; their fossilized remains are used to produce



diatomaceous earth, which has a range of uses such as filtration and insulation. Additionally, diatoms can reproduce sexually or asexually. One diatom genus, *Pseudo-nitzschia*, is known to be associated with harmful algal blooms.


Brown algae (Phaeophyta) are multicellular marine seaweeds. Some can be extremely large, such as the giant kelp (*Laminaria*). They have leaf-like blades, stalks, and structures called holdfasts that are used to attach to substrate. However, these are not true leaves, stems, or roots (**Figure 5.36**). Their photosynthetic pigments are chlorophyll *a*, chlorophyll *c*,  $\beta$ -carotene, and fucoxanthine. They use laminarin as a storage carbohydrate.



**Figure 5.36** (a) These large multicellular kelps are members of the brown algae. Note the “leaves” and “stems” that make them appear similar to green plants. (b) This is a species of red algae that is also multicellular. (c) The green alga *Halimeda incrassata*, shown here growing on the sea floor in shallow water, appears to have plant-like structures, but is not a true plant. (d) Bioluminescence, visible in the cresting wave in this picture, is a phenomenon of certain dinoflagellates. (e) Diatoms (pictured in this micrograph) produce siliceous tests (skeletons) that form diatomaceous earths. (f) Colonial green algae, like volvox in these three micrographs, exhibit simple cooperative associations of cells. (credit b: modification of work by Ed Bierman; credit c: modification of work by James St. John; credit d: modification of work by “catalano82”/Flickr; credit f: modification of work by Dr. Ralf Wagner)

The Archaeplastids include the green algae (Chlorophyta), the red algae (Rhodophyta), another group of green algae (Charophyta), and the land plants. The Charophyta are the most similar to land plants (**Figure 5.6**) because they share a mechanism of cell division and an important biochemical pathway, among other traits that the other groups do not have. Like land plants, the Charophyta and Chlorophyta have chlorophyll *a* and chlorophyll *b* as photosynthetic pigments, cellulose cell walls, and starch as a carbohydrate storage molecule. *Chlamydomonas* is a green alga that has

a single large chloroplast, two flagella, and a stigma (eyespot); it is important in molecular biology research (**Figure 5.37**).



**Figure 5.37** *Chlamydomonas* is a unicellular green alga.

*Chlorella* is a nonmotile, large, unicellular alga, and *Acetabularia* is an even larger unicellular green alga. The size of these organisms challenges the idea that all cells are small, and they have been used in genetics research since Joachim Hämmerling (1901–1980) began to work with them in 1943. *Volvox* is a colonial, unicellular alga (**Figure 5.37**). A larger, multicellular green alga is *Ulva*, also known as the sea lettuce because of its large, edible, green blades. The range of life forms within the Chlorophyta—from unicellular to various levels of coloniality to multicellular forms—has been a useful research model for understanding the evolution of multicellularity. The red algae are mainly multicellular but include some unicellular forms. They have rigid cell walls containing agar or carrageenan, which are useful as food solidifying agents and as a solidifier added to growth media for microbes.



### Check Your Understanding

Which groups of algae are associated with harmful algal blooms?

## 5.5 Lichens

### Learning Objectives

- Explain why lichens are included in the study of microbiology
- Describe the unique characteristics of a lichen and the role of each partner in the symbiotic relationship of a lichen
- Describe ways in which lichens are beneficial to the environment

No one has to worry about getting sick from a lichen infection, but lichens are interesting from a microbiological perspective and they are an important component of most terrestrial ecosystems. Lichens provide opportunities for study of close relationships between unrelated microorganisms. Lichens contribute to soil production by breaking down rock, and they are early colonizers in soilless environments such as lava flows. The cyanobacteria in some lichens can fix nitrogen and act as a nitrogen source in some environments. Lichens are also important soil stabilizers in some desert environments and they are an important winter food source for caribou and reindeer. Finally, lichens

produce compounds that have antibacterial effects, and further research may discover compounds that are medically useful to humans.


## Characteristics

A **lichen** (Figure 5.39) is a combination of two organisms, a green alga or cyanobacterium and an ascomycete fungus, living in a symbiotic relationship. Whereas algae normally grow only in aquatic or extremely moist environments, lichens can potentially be found on almost any surface (especially rocks) or as **epiphytes** (meaning that they grow on other plants).

In some ways, the symbiotic relationship between lichens and algae seems like a mutualism (a relationship in which both organisms benefit). The fungus can obtain photosynthates from the algae or cyanobacterium and the algae or cyanobacterium can grow in a drier environment than it could otherwise tolerate. However, most scientists consider this symbiotic relationship to be a controlled parasitism (a relationship in which one organism benefits and the other is harmed) because the photosynthetic organism grows less well than it would without the fungus. It is important to note that such symbiotic interactions fall along a continuum between conflict and cooperation.

Lichens are slow growing and can live for centuries. They have been used in foods and to extract chemicals as dyes or antimicrobial substances. Some are very sensitive to pollution and have been used as environmental indicators.

Lichens have a body called a thallus, an outer, tightly packed fungal layer called a **cortex**, and an inner, loosely packed fungal layer called a **medulla** (Figure 5.38). Lichens use hyphal bundles called **rhizines** to attach to the substrate.



**Figure 5.38** This cross-section of a lichen thallus shows its various components. The upper cortex of fungal hyphae provides protection. Photosynthesis occurs in the algal zone. The medulla consists of fungal hyphae. The lower cortex also provides protection. The rhizines anchor the thallus to the substrate.




### Check Your Understanding

What types of organisms are found in lichens?



## Lichen Diversity

Lichens are classified as fungi and the fungal partners belong to the Ascomycota and Basidiomycota. Lichens can also be grouped into types based on their morphology. There are three major types of lichens, although other types exist as well. Lichens that are tightly attached to the substrate, giving them a crusty appearance, are called **crustose lichens**. Those that have leaf-like lobes are **foliose lichens**; they may only be attached at one point in the growth form, and they also have a second cortex below the medulla. Finally, **fruticose lichens** have rounded structures and an overall branched appearance. **Figure 5.39** shows an example of each of the forms of lichens.



**Figure 5.39** Examples of the three types of lichens are shown here. (a) This is a crustose lichen found mostly on marine rocks, *Caloplaca marina*. (b) This is a foliose lichen, *Flavoparmelia caperata*. (c) This is a fruticose lichen, *Letharia vulpina*, which is sufficiently poisonous that it was once used to make arrowheads. (credit b and credit c: modification of work by Jason Hollinger)



### Check Your Understanding

What are the three growth forms of lichens?

### Clinical Focus

#### Resolution

Sarah's mother asks the doctor what she should do if the cream prescribed for Sarah's ringworm does not work. The doctor explains that ringworm is a general term for a condition caused by multiple species. The first step is to take a scraping for examination under the microscope, which the doctor has already done. He explains that he has identified the infection as a fungus, and that the antifungal cream works against the most common fungi associated with ringworm. However, the cream may not work against some species of fungus. If the cream is not working after a couple of weeks, Sarah should come in for another visit, at which time the doctor will take steps to identify the species of the fungus.

Positive identification of dermatophytes requires culturing. For this purpose, Sabouraud's agar may be used. In the case of Sarah's infection, which cleared up within 2 weeks of treatment, the culture would have a granular texture and would appear pale pink on top and red underneath. These features suggest that the fungus is *Trichophyton rubrum*, a common cause of ringworm.

Go back to the **previous** Clinical Focus box.



## Summary

---

### 5.1 Unicellular Eukaryotic Parasites

- **Protists** are a diverse, **polyphyletic** group of eukaryotic organisms.
- Protists may be unicellular or multicellular. They vary in how they get their nutrition, morphology, method of locomotion, and mode of reproduction.
- Important structures of protists include **contractile vacuoles**, cilia, flagella, **pellicles**, and pseudopodia; some lack organelles such as mitochondria.
- Taxonomy of protists is changing rapidly as relationships are reassessed using newer techniques.
- The protists include important pathogens and parasites.

### 5.2 The Parasitic Helminths

- Helminth parasites are included within the study of microbiology because they are often identified by looking for microscopic eggs and larvae.
- The two major groups of helminth parasites are the roundworms (Nematoda) and the flatworms (Platyhelminthes).
- Nematodes are common intestinal parasites often transmitted through undercooked foods, although they are also found in other environments.
- Platyhelminths include **tapeworms** and **flukes**, which are often transmitted through undercooked meat.

### 5.3 Fungi

- The fungi include diverse saprotrophic eukaryotic organisms with chitin cell walls
- Fungi can be unicellular or multicellular; some (like yeast) and fungal spores are microscopic, whereas some are large and conspicuous
- Reproductive types are important in distinguishing fungal groups
- Medically important species exist in the four fungal groups Zygomycota, Ascomycota, Basidiomycota, and Microsporidia
- Members of Zygomycota, Ascomycota, and Basidiomycota produce deadly toxins
- Important differences in fungal cells, such as ergosterols in fungal membranes, can be targets for antifungal medications, but similarities between human and fungal cells make it difficult to find targets for medications and these medications often have toxic adverse effects

### 5.4 Algae

- Algae are a diverse group of photosynthetic eukaryotic protists
- Algae may be unicellular or multicellular
- Large, multicellular algae are called seaweeds but are not plants and lack plant-like tissues and organs
- Although algae have little pathogenicity, they may be associated with toxic **algal blooms** that can and aquatic wildlife and contaminate seafood with toxins that cause paralysis
- Algae are important for producing **agar**, which is used as a solidifying agent in microbiological media, and **carrageenan**, which is used as a solidifying agent

### 5.5 Lichens

- **Lichens** are a symbiotic association between a fungus and an algae or a cyanobacterium
- The symbiotic association found in lichens is currently considered to be a controlled **parasitism**, in which the fungus benefits and the algae or cyanobacterium is harmed
- Lichens are slow growing and can live for centuries in a variety of habitats
- Lichens are environmentally important, helping to create soil, providing food, and acting as indicators of air pollution

## Review Questions

---

**Multiple Choice**

1. Which genus includes the causative agent for malaria?
  - a. *Euglena*
  - b. *Paramecium*
  - c. *Plasmodium*
  - d. *Trypanosoma*
2. Which protist is a concern because of its ability to contaminate water supplies and cause diarrheal illness?
  - a. *Plasmodium vivax*
  - b. *Toxoplasma gondii*
  - c. *Giardia lamblia*
  - d. *Trichomonas vaginalis*
3. A fluke is classified within which of the following?
  - a. Nematoda
  - b. Rotifera
  - c. Platyhelminthes
  - d. Annelida
4. A nonsegmented worm is found during a routine colonoscopy of an individual who reported having abdominal cramps, nausea, and vomiting. This worm is likely which of the following?
  - a. nematode
  - b. fluke
  - c. trematode
  - d. annelid
5. A segmented worm has male and female reproductive organs in each segment. Some use hooks to attach to the intestinal wall. Which type of worm is this?
  - a. fluke
  - b. nematode
  - c. cestode
  - d. annelid
6. Mushrooms are a type of which of the following?
  - a. conidia
  - b. ascus
  - c. polar tubule
  - d. basidiocarp
7. Which of the following is the most common cause of human yeast infections?
  - a. *Candida albicans*
  - b. *Blastomyces dermatitidis*
  - c. *Cryptococcus neoformans*
  - d. *Aspergillus fumigatus*
8. Which of the following is an ascomycete fungus associated with bat droppings that can cause a respiratory infection if inhaled?
  - a. *Candida albicans*
  - b. *Histoplasma capsulatum*
  - c. *Rhizopus stolonifera*
  - d. *Trichophyton rubrum*
9. Which polysaccharide found in red algal cell walls is a useful solidifying agent?
  - a. chitin
  - b. cellulose
  - c. phycoerythrin
  - d. agar
10. Which is the term for the hard outer covering of some dinoflagellates?
  - a. theca
  - b. thallus
  - c. mycelium
  - d. shell
11. Which protists are associated with red tides?
  - a. red algae
  - b. brown algae
  - c. dinoflagellates
  - d. green algae
12. You encounter a lichen with leafy structures. Which term describes this lichen?
  - a. crustose
  - b. foliose
  - c. fruticose
  - d. agarose
13. Which of the following is the term for the outer layer of a lichen?
  - a. the cortex
  - b. the medulla
  - c. the thallus
  - d. the theca
14. The fungus in a lichen is which of the following?
  - a. a basidiomycete
  - b. an ascomycete
  - c. a zygomycete
  - d. an apicomplexan

**Fill in the Blank**

15. The plasma membrane of a protist is called the \_\_\_\_\_.


16. Animals belong to the same supergroup as the kingdom \_\_\_\_\_.
17. Flukes are in class \_\_\_\_\_.
18. A species of worm in which there are distinct male and female individuals is described as \_\_\_\_\_.
19. Nonseptate hyphae are also called \_\_\_\_\_.
20. Unicellular fungi are called \_\_\_\_\_.
21. Some fungi have proven medically useful because they can be used to produce \_\_\_\_\_.
22. Structures in chloroplasts used to synthesize and store starch are called \_\_\_\_\_.
23. Algae with chloroplasts with three or four membranes are a result of \_\_\_\_\_.

### Short Answer

24. What are kinetoplastids?
25. Aside from a risk of birth defects, what other effect might a toxoplasmosis infection have?
26. What is the function of the ciliate macronucleus?
27. What is the best defense against tapeworm infection?
28. Which genera of fungi are common dermatophytes (fungi that cause skin infections)?
29. What is a dikaryotic cell?
30. What is a distinctive feature of diatoms?
31. Why are algae not considered parasitic?
32. Which groups contain the multicellular algae?
33. What are three ways that lichens are environmentally valuable?


### Critical Thinking

34. The protist shown in the picture has which of the following?
  - a. pseudopodia
  - b. flagella
  - c. a shell
  - d. cilia




**Figure 5.40** (credit: modification of work by Richard Robinson)

35. Protist taxonomy has changed greatly in recent years as relationships have been re-examined using newer approaches. How do newer approaches differ from older approaches?
36. What characteristics might make you think a protist could be pathogenic? Are certain nutritional characteristics, methods of locomotion, or morphological differences likely to be associated with the ability to cause disease?
37. Given the life cycle of the *Schistosoma* parasite, suggest a method of prevention of the disease.
38. Which of the drawings shows septate hyphae?



A




B

39. Explain the benefit of research into the pathways involved in the synthesis of chitin in fungi.

## Chapter 6

# Acellular Pathogens



**Figure 6.1** The year 2014 saw the first large-scale outbreak of Ebola virus (electron micrograph, left) in human populations in West Africa (right). Such epidemics are now widely reported and documented, but viral epidemics are sure to have plagued human populations since the origin of our species. (credit left: modification of work by Thomas W. Geisbert)

### Chapter Outline

- 6.1 Viruses
- 6.2 The Viral Life Cycle
- 6.3 Isolation, Culture, and Identification of Viruses
- 6.4 Viroids, Virusoids, and Prions

## Introduction

Public health measures in the developed world have dramatically reduced mortality from viral epidemics. But when epidemics do occur, they can spread quickly with global air travel. In 2009, an outbreak of H1N1 influenza spread across various continents. In early 2014, cases of Ebola in Guinea led to a massive epidemic in western Africa. This included the case of an infected man who traveled to the United States, sparking fears the epidemic might spread beyond Africa.

Until the late 1930s and the advent of the electron microscope, no one had seen a virus. Yet treatments for preventing or curing viral infections were used and developed long before that. Historical records suggest that by the 17th century, and perhaps earlier, inoculation (also known as variolation) was being used to prevent the viral disease smallpox in various parts of the world. By the late 18th century, Englishman Edward Jenner was inoculating patients with cowpox to prevent smallpox, a technique he coined *vaccination*.<sup>[1]</sup>

Today, the structure and genetics of viruses are well defined, yet new discoveries continue to reveal their complexities. In this chapter, we will learn about the structure, classification, and cultivation of viruses, and how they impact their hosts. In addition, we will learn about other infective particles such as viroids and prions.

1. S. Riedel "Edward Jenner and the History of Smallpox and Vaccination." *Baylor University Medical Center Proceedings* 18, no. 1 (January 2005): 21–25.



## 6.1 Viruses

### Learning Objectives

- Describe the general characteristics of viruses as pathogens
- Describe viral genomes
- Describe the general characteristics of viral life cycles
- Differentiate among bacteriophages, plant viruses, and animal viruses
- Describe the characteristics used to identify viruses as obligate intracellular parasites

Despite their small size, which prevented them from being seen with light microscopes, the discovery of a filterable component smaller than a bacterium that causes tobacco mosaic disease (TMD) dates back to 1892.<sup>[2]</sup> At that time, Dmitri Ivanovski, a Russian botanist, discovered the source of TMD by using a porcelain filtering device first invented by Charles Chamberland and Louis Pasteur in Paris in 1884. Porcelain Chamberland filters have a pore size of 0.1  $\mu\text{m}$ , which is small enough to remove all bacteria  $\geq 0.2 \mu\text{m}$  from any liquids passed through the device. An extract obtained from TMD-infected tobacco plants was made to determine the cause of the disease. Initially, the source of the disease was thought to be bacterial. It was surprising to everyone when Ivanovski, using a Chamberland filter, found that the cause of TMD was not removed after passing the extract through the porcelain filter. So if a bacterium was not the cause of TMD, what could be causing the disease? Ivanovski concluded the cause of TMD must be an extremely small bacterium or bacterial spore. Other scientists, including Martinus Beijerinck, continued investigating the cause of TMD. It was Beijerinck, in 1899, who eventually concluded the causative agent was not a bacterium but, instead, possibly a chemical, like a biological poison we would describe today as a toxin. As a result, the word *virus*, Latin for poison, was used to describe the cause of TMD a few years after Ivanovski's initial discovery. Even though he was not able to see the virus that caused TMD, and did not realize the cause was not a bacterium, Ivanovski is credited as the original discoverer of viruses and a founder of the field of virology.

Today, we can see viruses using electron microscopes (**Figure 6.2**) and we know much more about them. Viruses are distinct biological entities; however, their evolutionary origin is still a matter of speculation. In terms of taxonomy, they are not included in the tree of life because they are **acellular** (not consisting of cells). In order to survive and reproduce, viruses must infect a cellular host, making them obligate intracellular parasites. The genome of a virus

### Clinical Focus

#### Part 1

David, a 45-year-old journalist, has just returned to the U.S. from travels in Russia, China, and Africa. He is not feeling well, so he goes to his general practitioner complaining of weakness in his arms and legs, fever, headache, noticeable agitation, and minor discomfort. He thinks it may be related to a dog bite he suffered while interviewing a Chinese farmer. He is experiencing some prickling and itching sensations at the site of the bite wound, but he tells the doctor that the dog seemed healthy and that he had not been concerned until now. The doctor ordered a culture and sensitivity test to rule out bacterial infection of the wound, and the results came back negative for any possible pathogenic bacteria.

- Based on this information, what additional tests should be performed on the patient?
- What type of treatment should the doctor recommend?


Jump to the **next** Clinical Focus box.

2. H. Lecoq. "[Discovery of the First Virus, the Tobacco Mosaic Virus: 1892 or 1898?]." *Comptes Rendus de l'Academie des Sciences – Serie III – Sciences de la Vie* 324, no. 10 (2001): 929–933.

enters a host cell and directs the production of the viral components, proteins and nucleic acids, needed to form new virus particles called **virions**. New virions are made in the host cell by assembly of viral components. The new virions transport the viral genome to another host cell to carry out another round of infection. **Table 6.1** summarizes the properties of viruses.

Characteristics of Viruses
Infectious, acellular pathogens
Obligate intracellular parasites with host and cell-type specificity
DNA or RNA genome (never both)
Genome is surrounded by a protein capsid and, in some cases, a phospholipid membrane studded with viral glycoproteins
Lack genes for many products needed for successful reproduction, requiring exploitation of host-cell genomes to reproduce

**Table 6.1**



**Figure 6.2** (a) Tobacco mosaic virus (TMV) viewed with transmission electron microscope. (b) Plants infected with tobacco mosaic disease (TMD), caused by TMV. (credit a: modification of work by USDA Agricultural Research Service—scale-bar data from Matt Russell; credit b: modification of work by USDA Forest Service, Department of Plant Pathology Archive North Carolina State University)



### Check Your Understanding

- Why was the first virus investigated mistaken for a toxin?

## Hosts and Viral Transmission

Viruses can infect every type of host cell, including those of plants, animals, fungi, protists, bacteria, and archaea. Most viruses will only be able to infect the cells of one or a few species of organism. This is called the **host range**. However, having a wide host range is not common and viruses will typically only infect specific hosts and only specific cell types within those hosts. The viruses that infect bacteria are called **bacteriophages**, or simply phages. The word *phage* comes from the Greek word for devour. Other viruses are just identified by their host group, such as animal or plant viruses. Once a cell is infected, the effects of the virus can vary depending on the type of virus.

Viruses may cause abnormal growth of the cell or cell death, alter the cell's genome, or cause little noticeable effect in the cell.

Viruses can be transmitted through direct contact, indirect contact with fomites, or through a **vector**: an animal that transmits a pathogen from one host to another. Arthropods such as mosquitoes, ticks, and flies, are typical vectors for viral diseases, and they may act as **mechanical vectors** or **biological vectors**. Mechanical transmission occurs when the arthropod carries a viral pathogen on the outside of its body and transmits it to a new host by physical contact. Biological transmission occurs when the arthropod carries the viral pathogen inside its body and transmits it to the new host through biting.

In humans, a wide variety of viruses are capable of causing various infections and diseases. Some of the deadliest emerging pathogens in humans are viruses, yet we have few treatments or drugs to deal with viral infections, making them difficult to eradicate.

Viruses that can be transmitted from an animal host to a human host can cause zoonoses. For example, the avian influenza virus originates in birds, but can cause disease in humans. Reverse zoonoses are caused by infection of an animal by a virus that originated in a human.

## Micro Connections

### Fighting Bacteria with Viruses

The emergence of superbugs, or multidrug resistant bacteria, has become a major challenge for pharmaceutical companies and a serious health-care problem. According to a 2013 report by the US Centers for Disease Control and Prevention (CDC), more than 2 million people are infected with drug-resistant bacteria in the US annually, resulting in at least 23,000 deaths.<sup>[3]</sup> The continued use and overuse of antibiotics will likely lead to the evolution of even more drug-resistant strains.

One potential solution is the use of phage therapy, a procedure that uses bacteria-killing viruses (bacteriophages) to treat bacterial infections. Phage therapy is not a new idea. The discovery of bacteriophages dates back to the early 20th century, and phage therapy was first used in Europe in 1915 by the English bacteriologist Frederick Twort.<sup>[4]</sup> However, the subsequent discovery of penicillin and other antibiotics led to the near abandonment of this form of therapy, except in the former Soviet Union and a few countries in Eastern Europe. Interest in phage therapy outside of the countries of the former Soviet Union is only recently re-emerging because of the rise in antibiotic-resistant bacteria.<sup>[5]</sup>

Phage therapy has some advantages over antibiotics in that phages kill only one specific bacterium, whereas antibiotics kill not only the pathogen but also beneficial bacteria of the normal microbiota. Development of new antibiotics is also expensive for drug companies and for patients, especially for those who live in countries with high poverty rates.

Phages have also been used to prevent food spoilage. In 2006, the US Food and Drug Administration approved the use of a solution containing six bacteriophages that can be sprayed on lunch meats such as bologna, ham, and turkey to kill *Listeria monocytogenes*, a bacterium responsible for listeriosis, a form of food poisoning. Some consumers have concerns about the use of phages on foods, however, especially given the rising popularity of organic products. Foods that have been treated with phages must declare "bacteriophage preparation" in the list of ingredients or include a label declaring that the meat has been "treated with antimicrobial solution to reduce microorganisms."<sup>[6]</sup>

3. US Department of Health and Human Services, Centers for Disease Control and Prevention. "Antibiotic Resistance Threats in the United States, 2013." <http://www.cdc.gov/drugresistance/pdf/ar-threats-2013-508.pdf> (accessed September 22, 2015).

4. M. Clokie et al. "Phages in Nature." *Bacteriophage* 1, no. 1 (2011): 31–45.

5. A. Sulakvelidze et al. "Bacteriophage Therapy." *Antimicrobial Agents and Chemotherapy* 45, no. 3 (2001): 649–659.

6. US Food and Drug Administration. "FDA Approval of *Listeria*-specific Bacteriophage Preparation on Ready-to-Eat (RTE) Meat and Poultry Products." <http://www.fda.gov/food/ingredientspackaginglabeling/ucm083572.htm> (accessed September 22, 2015).




## Check Your Understanding

- Why do humans not have to be concerned about the presence of bacteriophages in their food?
- What are three ways that viruses can be transmitted between hosts?


## Viral Structures

In general, virions (viral particles) are small and cannot be observed using a regular light microscope. They are much smaller than prokaryotic and eukaryotic cells; this is an adaptation allowing viruses to infect these larger cells (see **Figure 6.3**). The size of a virion can range from 20 nm for small viruses up to 900 nm for typical, large viruses (see **Figure 6.4**). Recent discoveries, however, have identified new giant viral species, such as *Pandoravirus salinus* and *Pithovirus sibericum*, with sizes approaching that of a bacterial cell.<sup>[7]</sup>



**Figure 6.3** (a) In this transmission electron micrograph, a bacteriophage (a virus that infects bacteria) is dwarfed by the bacterial cell it infects. (b) An illustration of the bacteriophage in the micrograph. (credit a: modification of work by J.P. Nataro and S. Sears, Centers for Disease Control and Prevention—scale-bar data from Matt Russell)

7. N. Philippe et al. "Pandoraviruses: Amoeba Viruses with Genomes up to 2.5 Mb Reaching that of Parasitic Eukaryotes." *Science* 341, no. 6143 (2013): 281–286.



**Figure 6.4** The size of a virus is small relative to the size of most bacterial and eukaryotic cells and their organelles.

In 1935, after the development of the electron microscope, Wendell Stanley was the first scientist to crystallize the structure of the tobacco mosaic virus and discovered that it is composed of RNA and protein. In 1943, he isolated *Influenza B virus*, which contributed to the development of an influenza (flu) vaccine. Stanley's discoveries unlocked the mystery of the nature of viruses that had been puzzling scientists for over 40 years and his contributions to the field of virology led to him being awarded the Nobel Prize in 1946.




As a result of continuing research into the nature of viruses, we now know they consist of a nucleic acid (either RNA or DNA, but never both) surrounded by a protein coat called a **capsid** (see **Figure 6.5**). The interior of the capsid is not filled with cytosol, as in a cell, but instead it contains the bare necessities in terms of genome and enzymes needed to direct the synthesis of new virions. Each capsid is composed of protein subunits called **capsomeres** made of one or more different types of capsomere proteins that interlock to form the closely packed capsid.

There are two categories of viruses based on general composition. Viruses formed from only a nucleic acid and capsid are called **naked viruses** or **nonenveloped viruses**. Viruses formed with a nucleic-acid packed capsid surrounded by a lipid layer are called **enveloped viruses** (see **Figure 6.5**). The **viral envelope** is a small portion of phospholipid membrane obtained as the virion buds from a host cell. The viral envelope may either be intracellular or cytoplasmic in origin.

Extending outward and away from the capsid on some naked viruses and enveloped viruses are protein structures called **spikes**. At the tips of these spikes are structures that allow the virus to attach and enter a cell, like the influenza virus hemagglutinin spikes (H) or enzymes like the neuraminidase (N) influenza virus spikes that allow the virus to detach from the cell surface during release of new virions. Influenza viruses are often identified by their H and N spikes. For example, H1N1 influenza viruses were responsible for the pandemics in 1918 and 2009,<sup>[8]</sup> H2N2 for the pandemic in 1957, and H3N2 for the pandemic in 1968.


8. J. Cohen. "What's Old Is New: 1918 Virus Matches 2009 H1N1 Strain. *Science* 327, no. 5973 (2010): 1563–1564.





**Figure 6.5** (a) The naked adenovirus uses spikes made of glycoproteins from its capsid to bind to host cells. (b) The enveloped human immunodeficiency virus uses spikes made of glycoproteins embedded in its envelope to bind to host cells (credit b: modification of work by Centers for Disease Control and Prevention)

Viruses vary in the shape of their capsids, which can be either **helical**, **polyhedral**, or **complex**. A helical capsid forms the shape of tobacco mosaic virus (TMV), a naked helical virus, and Ebola virus, an enveloped helical virus. The capsid is cylindrical or rod shaped, with the genome fitting just inside the length of the capsid. Polyhedral capsids form the shapes of poliovirus and rhinovirus, and consist of a nucleic acid surrounded by a polyhedral (many-sided) capsid in the form of an icosahedron. An **icosahedral** capsid is a three-dimensional, 20-sided structure with 12 vertices. These capsids somewhat resemble a soccer ball. Both helical and polyhedral viruses can have envelopes. Viral shapes seen in certain types of bacteriophages, such as T4 phage, and poxviruses, like vaccinia virus, may have features of both polyhedral and helical viruses so they are described as a complex viral shape (see **Figure 6.6**). In the bacteriophage complex form, the genome is located within the polyhedral head and the **sheath** connects the head to the **tail fibers** and **tail pins** that help the virus attach to receptors on the host cell's surface. Poxviruses that have complex shapes are often brick shaped, with intricate surface characteristics not seen in the other categories of capsid.



**Figure 6.6** Viral capsids can be (a) helical, (b) polyhedral, or (c) have a complex shape. (credit a: modification of work by USDA ARS—scale-bar data from Matt Russell; credit b: modification of work by United States Department of Energy)



### Check Your Understanding

- Which types of viruses have spikes?

## Classification and Taxonomy of Viruses

Although viruses are not classified in the three domains of life, their numbers are great enough to require classification. Since 1971, the International Union of Microbiological Societies Virology Division has given the task of developing, refining, and maintaining a universal virus taxonomy to the International Committee on Taxonomy of Viruses (ICTV). Since viruses can mutate so quickly, it can be difficult to classify them into a genus and a species epithet using the binomial nomenclature system. Thus, the ICTV's viral nomenclature system classifies viruses into families and genera based on viral genetics, chemistry, morphology, and mechanism of multiplication. To date, the ICTV has classified known viruses in seven orders, 96 families, and 350 genera. Viral family names end in *-viridae* (e.g., *Parvoviridae*) and genus names end in *-virus* (e.g., *Parvovirus*). The names of viral orders, families, and genera are all italicized. When referring to a viral species, we often use a genus and species epithet such as *Pandoravirus dulcis* or *Pandoravirus salinus*.

The Baltimore classification system is an alternative to ICTV nomenclature. The Baltimore system classifies viruses according to their genomes (DNA or RNA, single versus double stranded, and mode of replication). This system thus creates seven groups of viruses that have common genetics and biology.

### Link to Learning



Explore the latest virus **taxonomy** (<http://www.openstaxcollege.org//22virustaxon>) at the ICTV website.

Aside from formal systems of nomenclature, viruses are often informally grouped into categories based on chemistry, morphology, or other characteristics they share in common. Categories may include naked or enveloped structure, single-stranded (ss) or double-stranded (ds) DNA or ss or ds RNA genomes, segmented or nonsegmented genomes, and positive-strand (+) or negative-strand (-) RNA. For example, herpes viruses can be classified as a dsDNA enveloped virus; human immunodeficiency virus (HIV) is a +ssRNA enveloped virus, and tobacco mosaic virus is a +ssRNA virus. Other characteristics such as host specificity, tissue specificity, capsid shape, and special genes or enzymes may also be used to describe groups of similar viruses. **Table 6.2** lists some of the most common viruses that are human pathogens by genome type.

**Common Pathogenic Viruses**

Genome	Family	Example Virus	Clinical Features
dsDNA, enveloped	<i>Poxviridae</i>	<i>Orthopoxvirus</i>	Skin papules, pustules, lesions
	<i>Poxviridae</i>	<i>Parapoxvirus</i>	Skin lesions
	<i>Herpesviridae</i>	<i>Simplexvirus</i>	Cold sores, genital herpes, sexually transmitted disease
dsDNA, naked	<i>Adenoviridae</i>	<i>Adenovirus</i>	Respiratory infection (common cold)
	<i>Papillomaviridae</i>	<i>Papillomavirus</i>	Genital warts, cervical, vulvar, or vaginal cancer
	<i>Reoviridae</i>	<i>Reovirus</i>	Gastroenteritis severe diarrhea (stomach flu)
ssDNA, naked	<i>Parvoviridae</i>	<i>Adeno-associated dependoparvovirus A</i>	Respiratory tract infection
	<i>Parvoviridae</i>	<i>Adeno-associated dependoparvovirus B</i>	Respiratory tract infection
dsRNA, naked	<i>Reoviridae</i>	<i>Rotavirus</i>	Gastroenteritis
+ssRNA, naked	<i>Picornaviridae</i>	<i>Enterovirus C</i>	Poliomyelitis
	<i>Picornaviridae</i>	<i>Rhinovirus</i>	Upper respiratory tract infection (common cold)
	<i>Picornaviridae</i>	<i>Hepatovirus</i>	Hepatitis

**Table 6.2**

### Common Pathogenic Viruses

Genome	Family	Example Virus	Clinical Features
+ssRNA, enveloped	<i>Togaviridae</i>	<i>Alphavirus</i>	Encephalitis, hemorrhagic fever
	<i>Togaviridae</i>	<i>Rubivirus</i>	Rubella
	<i>Retroviridae</i>	<i>Lentivirus</i>	Acquired immune deficiency syndrome (AIDS)
-ssRNA, enveloped	<i>Filoviridae</i>	<i>Zaire Ebolavirus</i>	Hemorrhagic fever
	<i>Orthomyxoviridae</i>	<i>Influenzavirus A, B, C</i>	Flu
	<i>Rhabdoviridae</i>	<i>Lyssavirus</i>	Rabies

Table 6.2



### Check Your Understanding

- What are the types of virus genomes?

### Classification of Viral Diseases

While the ICTV has been tasked with the biological classification of viruses, it has also played an important role in the classification of diseases caused by viruses. To facilitate the tracking of virus-related human diseases, the ICTV has created classifications that link to the International Classification of Diseases (ICD), the standard taxonomy of disease that is maintained and updated by the World Health Organization (WHO). The ICD assigns an alphanumeric code of up to six characters to every type of viral infection, as well as all other types of diseases, medical conditions, and causes of death. This ICD code is used in conjunction with two other coding systems (the Current Procedural Terminology, and the Healthcare Common Procedure Coding System) to categorize patient conditions for treatment and insurance reimbursement.

For example, when a patient seeks treatment for a viral infection, ICD codes are routinely used by clinicians to order laboratory tests and prescribe treatments specific to the virus suspected of causing the illness. This ICD code is then used by medical laboratories to identify tests that must be performed to confirm the diagnosis. The ICD code is used by the health-care management system to verify that all treatments and laboratory work performed are appropriate for the given virus. Medical coders use ICD codes to assign the proper code for procedures performed, and medical billers, in turn, use this information to process claims for reimbursement by insurance companies. Vital-records keepers use ICD codes to record cause of death on death certificates, and epidemiologists used ICD codes to calculate morbidity and mortality statistics.



### Check Your Understanding

- Identify two locations where you would likely find an ICD code.

## Clinical Focus

### Part 2

David's doctor was concerned that his symptoms included prickling and itching at the site of the dog bite; these sensations could be early symptoms of rabies. Several tests are available to diagnose rabies in live patients, but no single antemortem test is adequate. The doctor decided to take samples of David's blood, saliva, and skin for testing. The skin sample was taken from the nape of the neck (posterior side of the neck near the hairline). It was about 6-mm long and contained at least 10 hair follicles, including the superficial cutaneous nerve. An immunofluorescent staining technique was used on the skin biopsy specimen to detect rabies antibodies in the cutaneous nerves at the base of the hair follicles. A test was also performed on a serum sample from David's blood to determine whether any antibodies for the rabies virus had been produced.

Meanwhile, the saliva sample was used for reverse transcriptase-polymerase chain reaction (RT-PCR) analysis, a test that can detect the presence of viral nucleic acid (RNA). The blood tests came back positive for the presence of rabies virus antigen, prompting David's doctor to prescribe prophylactic treatment. David is given a series of intramuscular injections of human rabies immunoglobulin along with a series of rabies vaccines.

- Why does the immunofluorescent technique look for rabies antibodies rather than the rabies virus itself?
- If David has contracted rabies, what is his prognosis?

Jump to the **next** Clinical Focus box. Go back to the **previous** Clinical Focus box.

## 6.2 The Viral Life Cycle

### Learning Objectives

- Describe the lytic and lysogenic life cycles
- Describe the replication process of animal viruses
- Describe unique characteristics of retroviruses and latent viruses
- Discuss human viruses and their virus-host cell interactions
- Explain the process of transduction
- Describe the replication process of plant viruses

All viruses depend on cells for reproduction and metabolic processes. By themselves, viruses do not encode for all of the enzymes necessary for viral replication. But within a host cell, a virus can commandeer cellular machinery to produce more viral particles. Bacteriophages replicate only in the cytoplasm, since prokaryotic cells do not have a nucleus or organelles. In eukaryotic cells, most DNA viruses can replicate inside the nucleus, with an exception observed in the large DNA viruses, such as the poxviruses, that can replicate in the cytoplasm. RNA viruses that infect animal cells often replicate in the cytoplasm.


### The Life Cycle of Viruses with Prokaryote Hosts

The life cycle of bacteriophages has been a good model for understanding how viruses affect the cells they infect, since similar processes have been observed for eukaryotic viruses, which can cause immediate death of the cell or establish a latent or chronic infection. **Virulent phages** typically lead to the death of the cell through cell lysis. **Temperate phages**, on the other hand, can become part of a host chromosome and are replicated with the cell genome until such time as they are induced to make newly assembled viruses, or **progeny viruses**.



## The Lytic Cycle

During the **lytic cycle** of virulent phage, the bacteriophage takes over the cell, reproduces new phages, and destroys the cell. T-even phage is a good example of a well-characterized class of virulent phages. There are five stages in the bacteriophage lytic cycle (see **Figure 6.7**). **Attachment** is the first stage in the infection process in which the phage interacts with specific bacterial surface receptors (e.g., lipopolysaccharides and OmpC protein on host surfaces). Most phages have a narrow host range and may infect one species of bacteria or one strain within a species. This unique recognition can be exploited for targeted treatment of bacterial infection by phage therapy or for phage typing to identify unique bacterial subspecies or strains. The second stage of infection is entry or **penetration**. This occurs through contraction of the tail sheath, which acts like a hypodermic needle to inject the viral genome through the cell wall and membrane. The phage head and remaining components remain outside the bacteria.




**Figure 6.7** A virulent phage shows only the lytic cycle pictured here. In the lytic cycle, the phage replicates and lyses the host cell.

The third stage of infection is **biosynthesis** of new viral components. After entering the host cell, the virus synthesizes virus-encoded endonucleases to degrade the bacterial chromosome. It then hijacks the host cell to replicate, transcribe, and translate the necessary viral components (capsomeres, sheath, base plates, tail fibers, and viral enzymes) for the assembly of new viruses. Polymerase genes are usually expressed early in the cycle, while capsid and tail proteins are expressed later. During the **maturation** phase, new virions are created. To liberate free phages, the bacterial cell wall is disrupted by phage proteins such as holin or lysozyme. The final stage is release. Mature viruses burst out of the host cell in a process called **lysis** and the progeny viruses are liberated into the environment to infect new cells.

## The Lysogenic Cycle

In a **lysogenic cycle**, the phage genome also enters the cell through attachment and penetration. A prime example of a phage with this type of life cycle is the lambda phage. During the lysogenic cycle, instead of killing the host, the phage genome integrates into the bacterial chromosome and becomes part of the host. The integrated phage genome is called a **prophage**. A bacterial host with a prophage is called a **lysogen**. The process in which a bacterium is infected by a temperate phage is called **lysogeny**. It is typical of temperate phages to be latent or inactive within the cell. As the bacterium replicates its chromosome, it also replicates the phage's DNA and passes it on to new daughter cells during reproduction. The presence of the phage may alter the phenotype of the bacterium, since it can bring in extra genes (e.g., toxin genes that can increase bacterial virulence). This change in the host phenotype is called **lysogenic conversion** or **phage conversion**. Some bacteria, such as *Vibrio cholerae* and *Clostridium botulinum*, are less virulent in the absence of the prophage. The phages infecting these bacteria carry the toxin genes in their genome and enhance the virulence of the host when the toxin genes are expressed. In the case of *V. cholera*, phage encoded toxin can cause severe diarrhea; in *C. botulinum*, the toxin can cause paralysis. During lysogeny, the prophage will persist in the host chromosome until **induction**, which results in the excision of the viral genome from the host chromosome. After

induction has occurred the temperate phage can proceed through a lytic cycle and then undergo lysogeny in a newly infected cell (see **Figure 6.8**).



**Figure 6.8** A temperate bacteriophage has both lytic and lysogenic cycles. In the lysogenic cycle, phage DNA is incorporated into the host genome, forming a prophage, which is passed on to subsequent generations of cells. Environmental stressors such as starvation or exposure to toxic chemicals may cause the prophage to be excised and enter the lytic cycle.

### Link to Learning



This **video** (<http://www.openstaxcollege.org/l/22lysogeniclife>) illustrates the stages of the lysogenic life cycle of a bacteriophage and the transition to a lytic phase.




### Check Your Understanding

- Is a latent phage undetectable in a bacterium?

## Transduction

Transduction occurs when a bacteriophage transfers bacterial DNA from one bacterium to another during sequential infections. There are two types of transduction: generalized and specialized transduction. During the lytic cycle of viral replication, the virus hijacks the host cell, degrades the host chromosome, and makes more viral genomes. As it assembles and packages DNA into the phage head, packaging occasionally makes a mistake. Instead of packaging viral DNA, it takes a random piece of host DNA and inserts it into the capsid. Once released, this virion will then

inject the former host's DNA into a newly infected host. The asexual transfer of genetic information can allow for DNA recombination to occur, thus providing the new host with new genes (e.g., an antibiotic-resistance gene, or a sugar-metabolizing gene). **Generalized transduction** occurs when a random piece of bacterial chromosomal DNA is transferred by the phage during the lytic cycle. **Specialized transduction** occurs at the end of the lysogenic cycle, when the prophage is excised and the bacteriophage enters the lytic cycle. Since the phage is integrated into the host genome, the prophage can replicate as part of the host. However, some conditions (e.g., ultraviolet light exposure or chemical exposure) stimulate the prophage to undergo induction, causing the phage to excise from the genome, enter the lytic cycle, and produce new phages to leave host cells. During the process of excision from the host chromosome, a phage may occasionally remove some bacterial DNA near the site of viral integration. The phage and host DNA from one end or both ends of the integration site are packaged within the capsid and are transferred to the new, infected host. Since the DNA transferred by the phage is not randomly packaged but is instead a specific piece of DNA near the site of integration, this mechanism of gene transfer is referred to as specialized transduction (see **Figure 6.9**). The DNA can then recombine with host chromosome, giving the latter new characteristics. Transduction seems to play an important role in the evolutionary process of bacteria, giving them a mechanism for asexual exchange of genetic information.



**Figure 6.9** This flowchart illustrates the mechanism of specialized transduction. An integrated phage excises, bringing with it a piece of the DNA adjacent to its insertion point. On reinfection of a new bacterium, the phage DNA integrates along with the genetic material acquired from the previous host.




### Check Your Understanding

- Which phage life cycle is associated with which forms of transduction?

## Life Cycle of Viruses with Animal Hosts

Lytic animal viruses follow similar infection stages to bacteriophages: attachment, penetration, biosynthesis, maturation, and release (see **Figure 6.10**). However, the mechanisms of penetration, nucleic-acid biosynthesis, and release differ between bacterial and animal viruses. After binding to host receptors, animal viruses enter through endocytosis (engulfment by the host cell) or through membrane fusion (viral envelope with the host cell membrane). Many viruses are host specific, meaning they only infect a certain type of host; and most viruses only infect certain types of cells within tissues. This specificity is called a **tissue tropism**. Examples of this are demonstrated by the poliovirus, which exhibits tropism for the tissues of the brain and spinal cord, or the influenza virus, which has a primary tropism for the respiratory tract.






**Figure 6.10** In influenza virus infection, viral glycoproteins attach the virus to a host epithelial cell. As a result, the virus is engulfed. Viral RNA and viral proteins are made and assembled into new virions that are released by budding.

Animal viruses do not always express their genes using the normal flow of genetic information—from DNA to RNA to protein. Some viruses have a dsDNA genome like cellular organisms and can follow the normal flow. However, others may have ssDNA, dsRNA, or ssRNA genomes. The nature of the genome determines how the genome is replicated and expressed as viral proteins. If a genome is ssDNA, host enzymes will be used to synthesize a second strand that is complementary to the genome strand, thus producing dsDNA. The dsDNA can now be replicated, transcribed, and translated similar to host DNA.


If the viral genome is RNA, a different mechanism must be used. There are three types of RNA genome: dsRNA, **positive (+) single-strand (+ssRNA)** or **negative (-) single-strand RNA (-ssRNA)**. If a virus has a +ssRNA genome, it can be translated directly to make viral proteins. Viral genomic +ssRNA acts like cellular mRNA. However, if a virus contains a -ssRNA genome, the host ribosomes cannot translate it until the -ssRNA is replicated into +ssRNA by viral RNA-dependent RNA polymerase (RdRP) (see **Figure 6.11**). The RdRP is brought in by the virus and can be used to make +ssRNA from the original -ssRNA genome. The RdRP is also an important enzyme for the replication of dsRNA viruses, because it uses the negative strand of the double-stranded genome as a template to create +ssRNA. The newly synthesized +ssRNA copies can then be translated by cellular ribosomes.



**Figure 6.11** RNA viruses can contain +ssRNA that can be directly read by the ribosomes to synthesize viral proteins. Viruses containing -ssRNA must first use the -ssRNA as a template for the synthesis of +ssRNA before viral proteins can be synthesized.

An alternative mechanism for viral nucleic acid synthesis is observed in the **retroviruses**, which are +ssRNA viruses (see **Figure 6.12**). Single-stranded RNA viruses such as HIV carry a special enzyme called **reverse transcriptase** within the capsid that synthesizes a complementary ssDNA (cDNA) copy using the +ssRNA genome as a template. The ssDNA is then made into dsDNA, which can integrate into the host chromosome and become a permanent part of the host. The integrated viral genome is called a **provirus**. The virus now can remain in the host for a long time to establish a chronic infection. The provirus stage is similar to the prophage stage in a bacterial infection during the lysogenic cycle. However, unlike prophage, the provirus does not undergo excision after splicing into the genome.





**Figure 6.12** HIV, an enveloped, icosahedral retrovirus, attaches to a cell surface receptor of an immune cell and fuses with the cell membrane. Viral contents are released into the cell, where viral enzymes convert the single-stranded RNA genome into DNA and incorporate it into the host genome.



## Check Your Understanding

- Is RNA-dependent RNA polymerase made from a viral gene or a host gene?


## Persistent Infections

Persistent infection occurs when a virus is not completely cleared from the system of the host but stays in certain tissues or organs of the infected person. The virus may remain silent or undergo productive infection without seriously harming or killing the host. Mechanisms of persistent infection may involve the regulation of the viral or host gene expressions or the alteration of the host immune response. The two primary categories of persistent infections are latent infection and chronic infection. Examples of viruses that cause latent infections include herpes simplex virus (oral and genital herpes), varicella-zoster virus (chicken pox and shingles), and Epstein-Barr virus (mononucleosis). Hepatitis C virus and HIV are two examples of viruses that cause long-term chronic infections.

### Latent Infection

Not all animal viruses undergo replication by the lytic cycle. There are viruses that are capable of remaining hidden or dormant inside the cell in a process called latency. These types of viruses are known as **latent viruses** and may cause latent infections. Viruses capable of latency may initially cause an acute infection before becoming dormant.

For example, the varicella-zoster virus infects many cells throughout the body and causes chicken pox, characterized by a rash of blisters covering the skin. About 10 to 12 days postinfection, the disease resolves and the virus goes dormant, living within nerve-cell ganglia for years. During this time, the virus does not kill the nerve cells or continue replicating. It is not clear why the virus stops replicating within the nerve cells and expresses few viral proteins but, in some cases, typically after many years of dormancy, the virus is reactivated and causes a new disease called shingles (**Figure 6.13**). Whereas chicken pox affects many areas throughout the body, shingles is a nerve cell-specific disease emerging from the ganglia in which the virus was dormant.



**Figure 6.13** (a) Varicella-zoster, the virus that causes chicken pox, has an enveloped icosahedral capsid visible in this transmission electron micrograph. Its double-stranded DNA genome becomes incorporated in the host DNA. (b) After a period of latency, the virus can reactivate in the form of shingles, usually manifesting as a painful, localized rash on one side of the body. (credit a: modification of work by Erskine Palmer and B.G. Partin—scale-bar data from Matt Russell; credit b: modification of work by Rosmarie Voegtli)

Latent viruses may remain dormant by existing as circular viral genome molecules outside of the host chromosome. Others become proviruses by integrating into the host genome. During dormancy, viruses do not cause any symptoms of disease and may be difficult to detect. A patient may be unaware that he or she is carrying the virus unless a viral diagnostic test has been performed.

## Chronic Infection

A chronic infection is a disease with symptoms that are recurrent or persistent over a long time. Some viral infections can be chronic if the body is unable to eliminate the virus. HIV is an example of a virus that produces a chronic infection, often after a long period of latency. Once a person becomes infected with HIV, the virus can be detected in tissues continuously thereafter, but untreated patients often experience no symptoms for years. However, the virus maintains chronic persistence through several mechanisms that interfere with immune function, including preventing expression of viral antigens on the surface of infected cells, altering immune cells themselves, restricting expression of viral genes, and rapidly changing viral antigens through mutation. Eventually, the damage to the immune system results in progression of the disease leading to acquired immunodeficiency syndrome (AIDS). The various mechanisms that HIV uses to avoid being cleared by the immune system are also used by other chronically infecting viruses, including the hepatitis C virus.



### Check Your Understanding

- In what two ways can a virus manage to maintain a persistent infection?

## Life Cycle of Viruses with Plant Hosts

Plant viruses are more similar to animal viruses than they are to bacteriophages. Plant viruses may be enveloped or non-enveloped. Like many animal viruses, plant viruses can have either a DNA or RNA genome and be single stranded or double stranded. However, most plant viruses do not have a DNA genome; the majority have a +ssRNA genome, which acts like messenger RNA (mRNA). Only a minority of plant viruses have other types of genomes.

Plant viruses may have a narrow or broad host range. For example, the citrus tristeza virus infects only a few plants of the *Citrus* genus, whereas the cucumber mosaic virus infects thousands of plants of various plant families. Most plant viruses are transmitted by contact between plants, or by fungi, nematodes, insects, or other arthropods that act as mechanical vectors. However, some viruses can only be transferred by a specific type of insect vector; for example, a particular virus might be transmitted by aphids but not whiteflies. In some cases, viruses may also enter healthy plants through wounds, as might occur due to pruning or weather damage.

Viruses that infect plants are considered biotrophic parasites, which means that they can establish an infection without killing the host, similar to what is observed in the lysogenic life cycles of bacteriophages. Viral infection can be asymptomatic (latent) or can lead to cell death (lytic infection). The life cycle begins with the penetration of the virus into the host cell. Next, the virus is uncoated within the cytoplasm of the cell when the capsid is removed. Depending on the type of nucleic acid, cellular components are used to replicate the viral genome and synthesize viral proteins for assembly of new virions. To establish a systemic infection, the virus must enter a part of the vascular system of the plant, such as the phloem. The time required for systemic infection may vary from a few days to a few weeks depending on the virus, the plant species, and the environmental conditions. The virus life cycle is complete when it is transmitted from an infected plant to a healthy plant.




### Check Your Understanding

- What is the structure and genome of a typical plant virus?

## Viral Growth Curve

Unlike the growth curve for a bacterial population, the growth curve for a virus population over its life cycle does not follow a sigmoidal curve. During the initial stage, an inoculum of virus causes infection. In the **eclipse phase**, viruses bind and penetrate the cells with no virions detected in the medium. The chief difference that next appears in the viral

growth curve compared to a bacterial growth curve occurs when virions are released from the lysed host cell at the same time. Such an occurrence is called a **burst**, and the number of virions per bacterium released is described as the **burst size**. In a one-step multiplication curve for bacteriophage, the host cells lyse, releasing many viral particles to the medium, which leads to a very steep rise in **viral titer** (the number of virions per unit volume). If no viable host cells remain, the viral particles begin to degrade during the decline of the culture (see **Figure 6.14**).



**Figure 6.14** The one-step multiplication curve for a bacteriophage population follows three steps: 1) inoculation, during which the virions attach to host cells; 2) eclipse, during which entry of the viral genome occurs; and 3) burst, when sufficient numbers of new virions are produced and emerge from the host cell. The burst size is the maximum number of virions produced per bacterium.



### Check Your Understanding

- What aspect of the life cycle of a virus leads to the sudden increase in the growth curve?

### Eye on Ethics



#### Unregistered Treatments

Ebola is incurable and deadly. The outbreak in West Africa in 2014 was unprecedented, dwarfing other human Ebola epidemics in the level of mortality. Of 24,666 suspected or confirmed cases reported, 10,179 people died.<sup>[9]</sup>

No approved treatments or vaccines for Ebola are available. While some drugs have shown potential in laboratory studies and animal models, they have not been tested in humans for safety and effectiveness. Not only are these drugs untested or unregistered but they are also in short supply.

Given the great suffering and high mortality rates, it is fair to ask whether unregistered and untested medications are better than none at all. Should such drugs be dispensed and, if so, who should receive them,



in light of their extremely limited supplies? Is it ethical to treat untested drugs on patients with Ebola? On the other hand, is it ethical to withhold potentially life-saving drugs from dying patients? Or should the drugs perhaps be reserved for health-care providers working to contain the disease?

In August 2014, two infected US aid workers and a Spanish priest were treated with ZMapp, an unregistered drug that had been tested in monkeys but not in humans. The two American aid workers recovered, but the priest died. Later that month, the WHO released a report on the ethics of treating patients with the drug. Since Ebola is often fatal, the panel reasoned that it is ethical to give the unregistered drugs and unethical to withhold them for safety concerns. This situation is an example of “compassionate use” outside the well-established system of regulation and governance of therapies.

## Case in Point

### Ebola in the US

On September 24, 2014, Thomas Eric Duncan arrived at the Texas Health Presbyterian Hospital in Dallas complaining of a fever, headache, vomiting, and diarrhea—symptoms commonly observed in patients with the cold or the flu. After examination, an emergency department doctor diagnosed him with sinusitis, prescribed some antibiotics, and sent him home. Two days later, Duncan returned to the hospital by ambulance. His condition had deteriorated and additional blood tests confirmed that he has been infected with the Ebola virus.

Further investigations revealed that Duncan had just returned from Liberia, one of the countries in the midst of a severe Ebola epidemic. On September 15, nine days before he showed up at the hospital in Dallas, Duncan had helped transport an Ebola-stricken neighbor to a hospital in Liberia. The hospital continued to treat Duncan, but he died several days after being admitted.

The timeline of the Duncan case is indicative of the life cycle of the Ebola virus. The incubation time for Ebola ranges from 2 days to 21 days. Nine days passed between Duncan’s exposure to the virus infection and the appearance of his symptoms. This corresponds, in part, to the eclipse period in the growth of the virus population. During the eclipse phase, Duncan would have been unable to transmit the disease to others. However, once an infected individual begins exhibiting symptoms, the disease becomes very contagious. Ebola virus is transmitted through direct contact with droplets of bodily fluids such as saliva, blood, and vomit. Duncan could conceivably have transmitted the disease to others at any time after he began having symptoms, presumably some time before his arrival at the hospital in Dallas. Once a hospital realizes a patient like Duncan is infected with Ebola virus, the patient is immediately quarantined, and public health officials initiate a back trace to identify everyone with whom a patient like Duncan might have interacted during the period in which he was showing symptoms.

Public health officials were able to track down 10 high-risk individuals (family members of Duncan) and 50 low-risk individuals to monitor them for signs of infection. None contracted the disease. However, one of the nurses charged with Duncan’s care did become infected. This, along with Duncan’s initial misdiagnosis, made it clear that US hospitals needed to provide additional training to medical personnel to prevent a possible Ebola outbreak in the US.

- What types of training can prepare health professionals to contain emerging epidemics like the Ebola outbreak of 2014?
- What is the difference between a contagious pathogen and an infectious pathogen?





**Figure 6.15** Researchers working with Ebola virus use layers of defenses against accidental infection, including protective clothing, breathing systems, and negative air-pressure cabinets for bench work. (credit: Randal J. Schoepp)

### Link to Learning



For additional information about Ebola, please visit the **CDC** (<http://www.openstaxcollege.org/l/22ebolaccdc>) website.

## 6.3 Isolation, Culture, and Identification of Viruses


### Learning Objectives

- Discuss why viruses were originally described as filterable agents
- Describe the cultivation of viruses and specimen collection and handling
- Compare in vivo and in vitro techniques used to cultivate viruses

At the beginning of this chapter, we described how porcelain Chamberland filters with pores small enough to allow viruses to pass through were used to discover TMV. Today, porcelain filters have been replaced with membrane filters and other devices used to isolate and identify viruses.

### Isolation of Viruses

Unlike bacteria, many of which can be grown on an artificial nutrient medium, viruses require a living host cell for replication. Infected host cells (eukaryotic or prokaryotic) can be cultured and grown, and then the growth medium can be harvested as a source of virus. Virions in the liquid medium can be separated from the host cells by either centrifugation or filtration. Filters can physically remove anything present in the solution that is larger than the virions; the viruses can then be collected in the filtrate (see **Figure 6.16**).



**Figure 6.16** Membrane filters can be used to remove cells or viruses from a solution. (a) This scanning electron micrograph shows rod-shaped bacterial cells captured on the surface of a membrane filter. Note differences in the comparative size of the membrane pores and bacteria. Viruses will pass through this filter. (b) The size of the pores in the filter determines what is captured on the surface of the filter (animal [red] and bacteria [blue]) and removed from liquid passing through. Note the viruses (green) pass through the finer filter.




### Check Your Understanding

- What size filter pore is needed to collect a virus?

## Cultivation of Viruses


Viruses can be grown **in vivo** (within a whole living organism, plant, or animal) or **in vitro** (outside a living organism in cells in an artificial environment, such as a test tube, cell culture flask, or agar plate). Bacteriophages can be grown in the presence of a dense layer of bacteria (also called a **bacterial lawn**) grown in a 0.7% soft agar in a Petri dish or flat (horizontal) flask (see **Figure 6.17**). The agar concentration is decreased from the 1.5% usually used in culturing bacteria. The soft 0.7% agar allows the bacteriophages to easily diffuse through the medium. For lytic bacteriophages, lysing of the bacterial hosts can then be readily observed when a clear zone called a **plaque** is detected (see **Figure 6.17**). As the phage kills the bacteria, many plaques are observed among the cloudy bacterial lawn.



**Figure 6.17** (a) Flasks like this may be used to culture human or animal cells for viral culturing. (b) These plates contain bacteriophage T4 grown on an *Escherichia coli* lawn. Clear plaques are visible where host bacterial cells have been lysed. Viral titers increase on the plates to the left. (credit a: modification of work by National Institutes of Health; credit b: modification of work by American Society for Microbiology)

Animal viruses require cells within a host animal or tissue-culture cells derived from an animal. Animal virus cultivation is important for 1) identification and diagnosis of pathogenic viruses in clinical specimens, 2) production of vaccines, and 3) basic research studies. In vivo host sources can be a developing embryo in an embryonated bird's egg (e.g., chicken, turkey) or a whole animal. For example, most of the influenza vaccine manufactured for annual flu vaccination programs is cultured in hens' eggs.


The embryo or host animal serves as an incubator for viral replication (see **Figure 6.18**). Location within the embryo or host animal is important. Many viruses have a tissue tropism, and must therefore be introduced into a specific site for growth. Within an embryo, target sites include the amniotic cavity, the chorioallantoic membrane, or the yolk sac. Viral infection may damage tissue membranes, producing lesions called pox; disrupt embryonic development; or cause the death of the embryo.



**Figure 6.18** (a) The cells within chicken eggs are used to culture different types of viruses. (b) Viruses can be replicated in various locations within the egg, including the chorioallantoic membrane, the amniotic cavity, and the yolk sac. (credit a: modification of work by Hoang Chung)

For in vitro studies, various types of cells can be used to support the growth of viruses. A primary cell culture is freshly prepared from animal organs or tissues. Cells are extracted from tissues by mechanical scraping or mincing to release cells or by an enzymatic method using trypsin or collagenase to break up tissue and release single cells into suspension. Because of anchorage-dependence requirements, primary cell cultures require a liquid culture medium in a Petri dish or tissue-culture flask so cells have a solid surface such as glass or plastic for attachment and growth. Primary cultures usually have a limited life span. When cells in a primary culture undergo mitosis and a sufficient density of cells is produced, cells come in contact with other cells. When this cell-to-cell-contact occurs, mitosis is

triggered to stop. This is called contact inhibition and it prevents the density of the cells from becoming too high. To prevent contact inhibition, cells from the primary cell culture must be transferred to another vessel with fresh growth medium. This is called a secondary cell culture. Periodically, cell density must be reduced by pouring off some cells and adding fresh medium to provide space and nutrients to maintain cell growth. In contrast to primary cell cultures, continuous cell lines, usually derived from transformed cells or tumors, are often able to be subcultured many times or even grown indefinitely (in which case they are called immortal). Continuous cell lines may not exhibit anchorage dependency (they will grow in suspension) and may have lost their contact inhibition. As a result, continuous cell lines can grow in piles or lumps resembling small tumor growths (see **Figure 6.19**).



**Figure 6.19** Cells for culture are prepared by separating them from their tissue matrix. (a) Primary cell cultures grow attached to the surface of the culture container. Contact inhibition slows the growth of the cells once they become too dense and begin touching each other. At this point, growth can only be sustained by making a secondary culture. (b) Continuous cell cultures are not affected by contact inhibition. They continue to grow regardless of cell density.

An example of an immortal cell line is the HeLa cell line, which was originally cultivated from tumor cells obtained from Henrietta Lacks, a patient who died of cervical cancer in 1951. HeLa cells were the first continuous tissue-culture cell line and were used to establish tissue culture as an important technology for research in cell biology, virology, and medicine. Prior to the discovery of HeLa cells, scientists were not able to establish tissue cultures with any reliability or stability. More than six decades later, this cell line is still alive and being used for medical research. See this section's **Eye on Ethics** box to read more about this important cell line and the controversial means by which it was obtained.



### Check Your Understanding

- What property of cells makes periodic dilutions of primary cell cultures necessary?

## Eye on Ethics




### The Immortal Cell Line of Henrietta Lacks

In January 1951, Henrietta Lacks, a 30-year-old African American woman from Baltimore, was diagnosed with cervical cancer at John Hopkins Hospital. We now know her cancer was caused by the human papillomavirus (HPV). Cytopathic effects of the virus altered the characteristics of her cells in a process called transformation, which gives the cells the ability to divide continuously. This ability, of course, resulted in a cancerous tumor that eventually killed Mrs. Lacks in October at age 31. Before her death, samples of her cancerous cells were taken without her knowledge or permission. The samples eventually ended up in the possession of Dr. George Gey, a biomedical researcher at Johns Hopkins University. Gey was able to grow some of the cells from Lacks's sample, creating what is known today as the immortal HeLa cell line. These cells have the ability to live and grow indefinitely and, even today, are still widely used in many areas of research.

According to Lacks's husband, neither Henrietta nor the family gave the hospital permission to collect her tissue specimen. Indeed, the family was not aware until 20 years after Lacks's death that her cells were still alive and actively being used for commercial and research purposes. Yet HeLa cells have been pivotal in numerous research discoveries related to polio, cancer, and AIDS, among other diseases. The cells have also been commercialized, although they have never themselves been patented. Despite this, Henrietta Lacks's estate has never benefited from the use of the cells, although, in 2013, the Lacks family was given control over the publication of the genetic sequence of her cells.

This case raises several bioethical issues surrounding patients' informed consent and the right to know. At the time Lacks's tissues were taken, there were no laws or guidelines about informed consent. Does that mean she was treated fairly at the time? Certainly by today's standards, the answer would be no. Harvesting tissue or organs from a dying patient without consent is not only considered unethical but illegal, regardless of whether such an act could save other patients' lives. Is it ethical, then, for scientists to continue to use Lacks's tissues for research, even though they were obtained illegally by today's standards?

Ethical or not, Lacks's cells are widely used today for so many applications that it is impossible to list them all. Is this a case in which the ends justify the means? Would Lacks be pleased to know about her contribution to science and the millions of people who have benefited? Would she want her family to be compensated for the commercial products that have been developed using her cells? Or would she feel violated and exploited by the researchers who took part of her body without her consent? Because she was never asked, we will never know.







**Figure 6.20** A multiphoton fluorescence image of HeLa cells in culture. Various fluorescent stains have been used to show the DNA (cyan), microtubules (green), and Golgi apparatus (orange).



## Detection of a Virus

Regardless of the method of cultivation, once a virus has been introduced into a whole host organism, embryo, or tissue-culture cell, a sample can be prepared from the infected host, embryo, or cell line for further analysis under a brightfield, electron, or fluorescent microscope. **Cytopathic effects (CPEs)** are distinct observable cell abnormalities due to viral infection. CPEs can include loss of adherence to the surface of the container, changes in cell shape from flat to round, shrinkage of the nucleus, vacuoles in the cytoplasm, fusion of cytoplasmic membranes and the formation of multinucleated syncytia, inclusion bodies in the nucleus or cytoplasm, and complete cell lysis (see **Figure 6.21**).

Further pathological changes include viral disruption of the host genome and altering normal cells into transformed cells, which are the types of cells associated with carcinomas and sarcomas. The type or severity of the CPE depends on the type of virus involved. **Figure 6.21** lists CPEs for specific viruses.

Cytopathic Effects of Specific Viruses		
Virus	Cytopathic Effect	Example
<i>Paramyxovirus</i>	Syncytium and faint basophilic cytoplasmic inclusion bodies	 Micrograph showing a syncytium of cells with faint basophilic cytoplasmic inclusion bodies. Arrows point to the inclusion bodies. ASM MicrobeLibrary.org © Suchman and Blair
<i>Poxvirus</i>	Pink eosinophilic cytoplasmic inclusion bodies (arrows) and cell swelling	 Micrograph showing pink eosinophilic cytoplasmic inclusion bodies (arrows) and cell swelling. ASM MicrobeLibrary.org © Suchman and Blair
<i>Herpesvirus</i>	Cytoplasmic stranding (arrow) and nuclear inclusion bodies (dashed arrow)	 Micrograph showing cytoplasmic stranding (arrow) and nuclear inclusion bodies (dashed arrow). ASM MicrobeLibrary.org © Suchman and Blair
<i>Adenovirus</i>	Cell enlargement, rounding, and distinctive “grape-like” clusters	 Micrograph showing cell enlargement, rounding, and distinctive “grape-like” clusters. ASM MicrobeLibrary.org © Suchman and Blair

**Figure 6.21** (credit: American Society for Microbiology)

## Link to Learning









Watch this [video \(http://www.openstaxcollege.org/l/22virusesoncell\)](http://www.openstaxcollege.org/l/22virusesoncell) to learn about the effects of viruses on cells.

## Hemagglutination Assay

A serological assay is used to detect the presence of certain types of viruses in patient serum. Serum is the straw-colored liquid fraction of blood plasma from which clotting factors have been removed. Serum can be used in a direct assay called a hemagglutination assay to detect specific types of viruses in the patient's sample. Hemagglutination is the agglutination (clumping) together of erythrocytes (red blood cells). Many viruses produce surface proteins or spikes called hemagglutinins that can bind to receptors on the membranes of erythrocytes and cause the cells to agglutinate. Hemagglutination is observable without using the microscope, but this method does not always differentiate between infectious and noninfectious viral particles, since both can agglutinate erythrocytes.

To identify a specific pathogenic virus using hemagglutination, we must use an indirect approach. Proteins called antibodies, generated by the patient's immune system to fight a specific virus, can be used to bind to components such as hemagglutinins that are uniquely associated with specific types of viruses. The binding of the antibodies with the hemagglutinins found on the virus subsequently prevent erythrocytes from directly interacting with the virus. So when erythrocytes are added to the antibody-coated viruses, there is no appearance of agglutination; agglutination has been inhibited. We call these types of indirect assays for virus-specific antibodies hemagglutination inhibition (HAI) assays. HAI can be used to detect the presence of antibodies specific to many types of viruses that may be causing or have caused an infection in a patient even months or years after infection (see [Figure 6.22](#)). This assay is described in greater detail in [Agglutination Assays \(http://cnx.org/content/m58902/latest/\)](http://cnx.org/content/m58902/latest/).

	Components	Interaction	Microtiter Results
A	RBCs		No reaction 
B	Virus + RBCs		Hemagglutination 
C	Virus + Antibody + RBCs		Hemagglutination inhibition 

**Figure 6.22** This chart shows the possible outcomes of a hemagglutination test. Row A: Erythrocytes do not bind together and will sink to the bottom of the well plate; this becomes visible as a red dot in the center of the well. Row B: Many viruses have hemagglutinins that causes agglutination of erythrocytes; the resulting hemagglutination forms a lattice structure that results in red color throughout the well. Row C: Virus-specific antibody, the viruses, and the erythrocytes are added to the well plate. The virus-specific antibodies inhibit agglutination, as can be seen as a red dot in the bottom of the well.



### Check Your Understanding

- What is the outcome of a positive HIA test?

### Nucleic Acid Amplification Test

Nucleic acid amplification tests (NAAT) are used in molecular biology to detect unique nucleic acid sequences of viruses in patient samples. Polymerase chain reaction (PCR) is an NAAT used to detect the presence of viral DNA in a patient's tissue or body fluid sample. PCR is a technique that amplifies (i.e., synthesizes many copies) of a viral DNA segment of interest. Using PCR, short nucleotide sequences called primers bind to specific sequences of viral DNA, enabling identification of the virus.

Reverse transcriptase-polymerase chain reaction (RT-PCR) is an NAAT used to detect the presence of RNA viruses. RT-PCR differs from PCR in that the enzyme reverse transcriptase (RT) is used to make a cDNA from the small amount of viral RNA in the specimen. The cDNA can then be amplified by PCR. Both PCR and RT-PCR are used to detect and confirm the presence of the viral nucleic acid in patient specimens.

## Case in Point

### HPV Scare

Michelle, a 21-year-old nursing student, came to the university clinic worried that she might have been exposed to a sexually transmitted disease (STD). Her sexual partner had recently developed several bumps on the base of his penis. He had put off going to the doctor, but Michelle suspects they are genital warts caused by HPV. She is especially concerned because she knows that HPV not only causes warts but is a prominent cause of cervical cancer. She and her partner always use condoms for contraception, but she is not confident that this precaution will protect her from HPV.


Michelle's physician finds no physical signs of genital warts or any other STDs, but recommends that Michelle get a Pap smear along with an HPV test. The Pap smear will screen for abnormal cervical cells and the CPEs associated with HPV; the HPV test will test for the presence of the virus. If both tests are negative, Michelle can be more assured that she most likely has not become infected with HPV. However, her doctor suggests it might be wise for Michelle to get vaccinated against HPV to protect herself from possible future exposure.

- Why does Michelle's physician order two different tests instead of relying on one or the other?

## Enzyme Immunoassay

Enzyme immunoassays (EIAs) rely on the ability of antibodies to detect and attach to specific biomolecules called antigens. The detecting antibody attaches to the target antigen with a high degree of specificity in what might be a complex mixture of biomolecules. Also included in this type of assay is a colorless enzyme attached to the detecting antibody. The enzyme acts as a tag on the detecting antibody and can interact with a colorless substrate, leading to the production of a colored end product. EIAs often rely on layers of antibodies to capture and react with antigens, all of which are attached to a membrane filter (see **Figure 6.23**). EIAs for viral antigens are often used as preliminary screening tests. If the results are positive, further confirmation will require tests with even greater sensitivity, such as a Western blot or an NAAT. EIAs are discussed in more detail in **Enzyme Immunoassays (EIA) and Enzyme-Linked Immunosorbent Assays (ELISA)** (<http://cnx.org/content/m58903/latest/>) .





**Figure 6.23** Similar to rapid, over-the-counter pregnancy tests, EIAs for viral antigens require a few drops of diluted patient serum or plasma applied to a membrane filter. The membrane filter has been previously modified and embedded with antibody to viral antigen and internal controls. Antibody conjugate is added to the filter, with the targeted antibody attached to the antigen (in the case of a positive test). Excess conjugate is washed off the filter. Substrate is added to activate the enzyme-mediated reaction to reveal the color change of a positive test.



## Check Your Understanding

- What typically indicates a positive EIA test?

### Clinical Focus

#### Part 3

Along with the RT/PCR analysis, David's saliva was also collected for viral cultivation. In general, no single diagnostic test is sufficient for antemortem diagnosis, since the results will depend on the sensitivity of the assay, the quantity of virions present at the time of testing, and the timing of the assay, since release of virions in the saliva can vary. As it turns out, the result was negative for viral cultivation from the saliva. This is not surprising to David's doctor, because one negative result is not an absolute indication of the absence of infection. It may be that the number of virions in the saliva is low at the time of sampling. It is not unusual to repeat the test at intervals to enhance the chance of detecting higher virus loads.

- Should David's doctor modify his course of treatment based on these test results?

*Jump to the **next** Clinical Focus box. Go back to the **previous** Clinical Focus box.*

## 6.4 Viroids, Virusoids, and Prions

### Learning Objectives

- Describe viroids and their unique characteristics
- Describe virusoids and their unique characteristics
- Describe prions and their unique characteristics

Research attempts to discover the causative agents of previously uninvestigated diseases have led to the discovery of nonliving disease agents quite different from viruses. These include particles consisting only of RNA or only of protein that, nonetheless, are able to self-propagate at the expense of a host—a key similarity to viruses that allows them to cause disease conditions. To date, these discoveries include viroids, virusoids, and the proteinaceous prions.

### Viroids

In 1971, Theodor Diener, a pathologist working at the Agriculture Research Service, discovered an acellular particle that he named a viroid, meaning “virus-like.” **Viroids** consist only of a short strand of circular RNA capable of self-replication. The first viroid discovered was found to cause potato tuber spindle disease, which causes slower sprouting and various deformities in potato plants (see **Figure 6.24**). Like viruses, potato spindle tuber viroids (PSTVs) take control of the host machinery to replicate their RNA genome. Unlike viruses, viroids do not have a protein coat to protect their genetic information.



**Figure 6.24** These potatoes have been infected by the potato spindle tuber viroid (PSTV), which is typically spread when infected knives are used to cut healthy potatoes, which are then planted. (credit: Pamela Roberts)

Viroids can result in devastating losses of commercially important agricultural food crops grown in fields and orchards. Since the discovery of PSTV, other viroids have been discovered that cause diseases in plants. Tomato planta macho viroid (TPMVd) infects tomato plants, which causes loss of chlorophyll, disfigured and brittle leaves, and very small tomatoes, resulting in loss of productivity in this field crop. Avocado sunblotch viroid (ASBVd) results in lower yields and poorer-quality fruit. ASBVd is the smallest viroid discovered thus far that infects plants. Peach latent mosaic viroid (PLMVd) can cause necrosis of flower buds and branches, and wounding of ripened fruit, which leads to fungal and bacterial growth in the fruit. PLMVd can also cause similar pathological changes in plums, nectarines, apricots, and cherries, resulting in decreased productivity in these orchards, as well. Viroids, in general, can be dispersed mechanically during crop maintenance or harvesting, vegetative reproduction, and possibly via seeds and insects, resulting in a severe drop in food availability and devastating economic consequences.



### Check Your Understanding

- What is the genome of a viroid made of?

## Virusoids

A second type of pathogenic RNA that can infect commercially important agricultural crops are the **virusoids**, which are subviral particles best described as non-self-replicating ssRNAs. RNA replication of virusoids is similar to that of viroids but, unlike viroids, virusoids require that the cell also be infected with a specific “helper” virus. There are currently only five described types of virusoids and their associated helper viruses. The helper viruses are all from the family of Sobemoviruses. An example of a helper virus is the subterranean clover mottle virus, which has an associated virusoid packaged inside the viral capsid. Once the helper virus enters the host cell, the virusoids are released and can be found free in plant cell cytoplasm, where they possess ribozyme activity. The helper virus undergoes typical viral replication independent of the activity of the virusoid. The virusoid genomes are small, only 220 to 388 nucleotides long. A virusoid genome does not code for any proteins, but instead serves only to replicate virusoid RNA.

Virusoids belong to a larger group of infectious agents called satellite RNAs, which are similar pathogenic RNAs found in animals. Unlike the plant virusoids, satellite RNAs may encode for proteins; however, like plant virusoids, satellite RNAs must coinfect with a helper virus to replicate. One satellite RNA that infects humans and that has been described by some scientists as a virusoid is the hepatitis delta virus (HDV), which, by some reports, is also called hepatitis delta virusoid. Much larger than a plant virusoid, HDV has a circular, ssRNA genome of 1,700 nucleotides and can direct the biosynthesis of HDV-associated proteins. The HDV helper virus is the hepatitis B virus (HBV). Coinfection with HBV and HDV results in more severe pathological changes in the liver during infection, which is how HDV was first discovered.




## Check Your Understanding

- What is the main difference between a viroid and a virusoid?


## Prions

At one time, scientists believed that any infectious particle must contain DNA or RNA. Then, in 1982, Stanley Prusiner, a medical doctor studying scrapie (a fatal, degenerative disease in sheep) discovered that the disease was caused by proteinaceous infectious particles, or **prions**. Because proteins are acellular and do not contain DNA or RNA, Prusiner's findings were originally met with resistance and skepticism; however, his research was eventually validated, and he received the Nobel Prize in Physiology or Medicine in 1997.

A prion is a misfolded rogue form of a normal protein (PrP<sup>C</sup>) found in the cell. This rogue prion protein (PrP<sup>Sc</sup>), which may be caused by a genetic mutation or occur spontaneously, can be infectious, stimulating other endogenous normal proteins to become misfolded, forming plaques (see **Figure 6.25**). Today, prions are known to cause various forms of **transmissible spongiform encephalopathy (TSE)** in human and animals. TSE is a rare degenerative disorder that affects the brain and nervous system. The accumulation of rogue proteins causes the brain tissue to become sponge-like, killing brain cells and forming holes in the tissue, leading to brain damage, loss of motor coordination, and dementia (see **Figure 6.26**). Infected individuals are mentally impaired and become unable to move or speak. There is no cure, and the disease progresses rapidly, eventually leading to death within a few months or years.



**Figure 6.25** Endogenous normal prion protein (PrP<sup>C</sup>) is converted into the disease-causing form (PrP<sup>Sc</sup>) when it encounters this variant form of the protein. PrP<sup>Sc</sup> may arise spontaneously in brain tissue, especially if a mutant form of the protein is present, or it may originate from misfolded prions consumed in food that eventually find their way into brain tissue. (credit b: modification of work by University of Chicago—scale-bar data from Matt Russell)



**Figure 6.26** Creutzfeldt-Jakob disease (CJD) is a fatal disease that causes degeneration of neural tissue. (a) These brain scans compare a normal brain to one with CJD. (b) Compared to a normal brain, the brain tissue of a CJD patient is full of sponge-like lesions, which result from abnormal formations of prion protein. (credit a (left): modification of work by Dr. Laughlin Dawes; credit b (top): modification of work by Suzanne Wakim; credit b (bottom): modification of work by Centers for Disease Control and Prevention)

TSEs in humans include kuru, fatal familial insomnia, Gerstmann-Straussler-Scheinker disease, and Creutzfeldt-Jakob disease (see **Figure 6.26**). TSEs in animals include mad cow, scrapie (in sheep and goats), and chronic wasting disease (in elk and deer). TSEs can be transmitted between animals and from animals to humans by eating contaminated meat or animal feed. Transmission between humans can occur through heredity (as is often the case with GSS and CJD) or by contact with contaminated tissue, as might occur during a blood transfusion or organ transplant. There is no evidence for transmission via casual contact with an infected person. **Table 6.3** lists TSEs that affect humans and their modes of transmission.

**Transmissible Spongiform Encephalopathies (TSEs) in Humans**

Disease	Mechanism(s) of Transmission <sup>[10]</sup>
Sporadic CJD (sCJD)	Not known; possibly by alteration of normal prior protein (PrP) to rogue form due to somatic mutation
Variant CJD (vCJD)	Eating contaminated cattle products and by secondary bloodborne transmission
Familial CJD (fCJD)	Mutation in germline PrP gene
Iatrogenic CJD (iCJD)	Contaminated neurosurgical instruments, corneal graft, gonadotrophic hormone, and, secondarily, by blood transfusion
Kuru	Eating infected meat through ritualistic cannibalism
Gerstmann-Straussler-Scheinker disease (GSS)	Mutation in germline PrP gene

**Table 6.3**

10. National Institute of Neurological Disorders and Stroke. "Creutzfeldt-Jakob Disease Fact Sheet." [http://www.ninds.nih.gov/disorders/cjd/detail\\_cjd.htm](http://www.ninds.nih.gov/disorders/cjd/detail_cjd.htm) (accessed December 31, 2015).



### Transmissible Spongiform Encephalopathies (TSEs) in Humans

Disease	Mechanism(s) of Transmission
Fatal familial insomnia (FFI)	Mutation in germline PrP gene

**Table 6.3**

Prions are extremely difficult to destroy because they are resistant to heat, chemicals, and radiation. Even standard sterilization procedures do not ensure the destruction of these particles. Currently, there is no treatment or cure for TSE disease, and contaminated meats or infected animals must be handled according to federal guidelines to prevent transmission.



#### Check Your Understanding

- Does a prion have a genome?

#### Link to Learning



For more information on the handling of animals and prion-contaminated materials, visit the guidelines published on the **CDC** (<http://www.openstaxcollege.org/l/22cdcccontaminat>) and **WHO** (<http://www.openstaxcollege.org/l/22whocontaminat>) websites.

#### Clinical Focus

##### Resolution

A few days later, David's doctor receives the results of the immunofluorescence test on his skin sample. The test is negative for rabies antigen. A second viral antigen test on his saliva sample also comes back negative. Despite these results, the doctor decides to continue David's current course of treatment. Given the positive RT-PCR test, it is best not to rule out a possible rabies infection.

Near the site of the bite, David receives an injection of rabies immunoglobulin, which attaches to and inactivates any rabies virus that may be present in his tissues. Over the next 14 days, he receives a series of four rabies-specific vaccinations in the arm. These vaccines activate David's immune response and help his body recognize and fight the virus. Thankfully, with treatment, David symptoms improve and he makes a full recovery.

Not all rabies cases have such a fortunate outcome. In fact, rabies is usually fatal once the patient starts to exhibit symptoms, and postbite treatments are mainly palliative (i.e., sedation and pain management).

Go back to the *previous* Clinical Focus box.

## Summary

---

### 6.1 Viruses

- Viruses are generally ultramicroscopic, typically from 20 nm to 900 nm in length. Some large viruses have been found.
- **Virions** are acellular and consist of a nucleic acid, DNA or RNA, but not both, surrounded by a protein **capsid**. There may also be a phospholipid membrane surrounding the capsid.
- Viruses are obligate intracellular parasites.
- Viruses are known to infect various types of cells found in plants, animals, fungi, protists, bacteria, and archaea. Viruses typically have limited **host ranges** and infect specific cell types.
- Viruses may have **helical, polyhedral, or complex** shapes.
- Classification of viruses is based on morphology, type of nucleic acid, host range, cell specificity, and enzymes carried within the virion.
- Like other diseases, viral diseases are classified using ICD codes.

### 6.2 The Viral Life Cycle

- Many viruses target specific hosts or tissues. Some may have more than one host.
- Many viruses follow several stages to infect host cells. These stages include **attachment, penetration, uncoating, biosynthesis, maturation, and release**.
- Bacteriophages have a **lytic or lysogenic cycle**. The lytic cycle leads to the death of the host, whereas the lysogenic cycle leads to integration of phage into the host genome.
- Bacteriophages inject DNA into the host cell, whereas animal viruses enter by endocytosis or membrane fusion.
- Animal viruses can undergo **latency**, similar to lysogeny for a bacteriophage.
- The majority of plant viruses are positive-strand ssRNA and can undergo latency, chronic, or lytic infection, as observed for animal viruses.
- The growth curve of bacteriophage populations is a **one-step multiplication curve** and not a sigmoidal curve, as compared to the bacterial growth curve.
- Bacteriophages transfer genetic information between hosts using either **generalized or specialized transduction**.

### 6.3 Isolation, Culture, and Identification of Viruses

- Viral cultivation requires the presence of some form of host cell (whole organism, embryo, or cell culture).
- Viruses can be isolated from samples by filtration.
- Viral filtrate is a rich source of released virions.
- Bacteriophages are detected by presence of clear **plaques** on bacterial lawn.
- Animal and plant viruses are detected by **cytopathic effects**, molecular techniques (PCR, RT-PCR), enzyme immunoassays, and serological assays (hemagglutination assay, hemagglutination inhibition assay).

### 6.4 Viroids, Virusoids, and Prions

- Other acellular agents such as **viroids, virusoids, and prions** also cause diseases. Viroids consist of small, naked ssRNAs that cause diseases in plants. Virusoids are ssRNAs that require other helper viruses to establish an infection. Prions are proteinaceous infectious particles that cause **transmissible spongiform encephalopathies**.
- Prions are extremely resistant to chemicals, heat, and radiation.
- There are no treatments for prion infection.

## Review Questions

---

**Multiple Choice**

1. The component(s) of a virus that is/are extended from the envelope for attachment is/are the:
  - a. capsomeres
  - b. spikes
  - c. nucleic acid
  - d. viral whiskers
2. Which of the following does a virus lack? Select all that apply.
  - a. ribosomes
  - b. metabolic processes
  - c. nucleic acid
  - d. glycoprotein
3. The envelope of a virus is derived from the host's
  - a. nucleic acids
  - b. membrane structures
  - c. cytoplasm
  - d. genome
4. In naming viruses, the family name ends with \_\_\_\_\_ and genus name ends with \_\_\_\_\_.
  - a. *-virus*; *-viridae*
  - b. *-viridae*; *-virus*
  - c. *-virion*; *virus*
  - d. *-virus*; *virion*
5. What is another name for a nonenveloped virus?
  - a. enveloped virus
  - b. provirus
  - c. naked virus
  - d. latent virus
6. Which of the following leads to the destruction of the host cells?
  - a. lysogenic cycle
  - b. lytic cycle
  - c. prophage
  - d. temperate phage
7. A virus obtains its envelope during which of the following phases?
  - a. attachment
  - b. penetration
  - c. assembly
  - d. release
8. Which of the following components is brought into a cell by HIV?
  - a. a DNA-dependent DNA polymerase
  - b. RNA polymerase
  - c. ribosome
  - d. reverse transcriptase
9. A positive-strand RNA virus:
  - a. must first be converted to a mRNA before it can be translated.
  - b. can be used directly to translate viral proteins.
  - c. will be degraded by host enzymes.
  - d. is not recognized by host ribosomes.
10. What is the name for the transfer of genetic information from one bacterium to another bacterium by a phage?
  - a. transduction
  - b. penetration
  - c. excision
  - d. translation
11. Which of the followings cannot be used to culture viruses?
  - a. tissue culture
  - b. liquid medium only
  - c. embryo
  - d. animal host
12. Which of the following tests can be used to detect the presence of a specific virus?
  - a. EIA
  - b. RT-PCR
  - c. PCR
  - d. all of the above
13. Which of the following is NOT a cytopathic effect?
  - a. transformation
  - b. cell fusion
  - c. mononucleated cell
  - d. inclusion bodies
14. Which of these infectious agents do not have nucleic acid?
  - a. viroids
  - b. viruses
  - c. bacteria
  - d. prions
15. Which of the following is true of prions?
  - a. They can be inactivated by boiling at 100 °C.
  - b. They contain a capsid.
  - c. They are a rogue form of protein, PrP.
  - d. They can be reliably inactivated by an autoclave.

**True/False**

16. True or False: Scientists have identified viruses that are able to infect fungal cells.

**Fill in the Blank**

17. A virus that infects a bacterium is called a/an \_\_\_\_\_.

18. A/an \_\_\_\_\_ virus possesses characteristics of both a polyhedral and helical virus.

19. A virus containing only nucleic acid and a capsid is called a/an \_\_\_\_\_ virus or \_\_\_\_\_ virus.

20. The \_\_\_\_\_ on the bacteriophage allow for binding to the bacterial cell.

21. An enzyme from HIV that can make a copy of DNA from RNA is called \_\_\_\_\_.

22. For lytic viruses, \_\_\_\_\_ is a phase during a viral growth curve when the virus is not detected.

23. Viruses can be diagnosed and observed using a(n) \_\_\_\_\_ microscope.

24. Cell abnormalities resulting from a viral infection are called \_\_\_\_\_.

25. Both viroids and virusoids have a(n) \_\_\_\_\_ genome, but virusoids require a(n) \_\_\_\_\_ to reproduce.

**Short Answer**

26. Discuss the geometric differences among helical, polyhedral, and complex viruses.

27. What was the meaning of the word “virus” in the 1880s and why was it used to describe the cause of tobacco mosaic disease?

28. Briefly explain the difference between the mechanism of entry of a T-even bacteriophage and an animal virus.

29. Discuss the difference between generalized and specialized transduction.


30. Differentiate between lytic and lysogenic cycles.

31. Briefly explain the various methods of culturing viruses.

32. Describe the disease symptoms observed in animals infected with prions.

**Critical Thinking**


33. Name each labeled part of the illustrated bacteriophage.



34. In terms of evolution, which do you think arises first? The virus or the host? Explain your answer.

35. Do you think it is possible to create a virus in the lab? Imagine that you are a mad scientist. Describe how you would go about creating a new virus.

36. Label the five stages of a bacteriophage infection in the figure:




37. Bacteriophages have lytic and lysogenic cycles. Discuss the advantages and disadvantages for the phage.

38. How does reverse transcriptase aid a retrovirus in establishing a chronic infection?

39. Discuss some methods by which plant viruses are transmitted from a diseased plant to a healthy one.

40. Label the components indicated by arrows.



41. What are some characteristics of the viruses that are similar to a computer virus?

42. Does a prion replicate? Explain.





## Answer Key

---

### Chapter 1

1. D 2. D 3. A 4. B 5. C 6. C 7. D 8. A 9. B 10. D 11. C 12. A 13. D 14. B 15. scientific history 16. Lyme 17. fermentation 18. genus, species 19. Protista and Monera 20. Prokaryotes 21. Viruses 22. pathogen 23. helminths 24. virology 25. nucleus

### Chapter 2

1. C 2. A 3. D 4. B 5. C 6. D 7. B 8. E 9. D 10. D 11. B 12. refraction 13. compound 14. fluorochromes 15. atomic force microscope 16. 400× 17. acid-fast 18. Gram stain

### Chapter 3

1. C 2. D 3. C 4. A 5. C 6. B 7. C 8. C, D 9. A 10. D 11. D 12. D 13. B 14. C 15. D 16. B 17. A 18. False 19. False 20. True 21. spontaneous generation 22. epidemiology 23. miasma 24. Robert Hooke 25. bacilli 26. volutin (or metachromatic granule) 27. hydrogen peroxide 28. actin 29. Pasteur designed a flask in which media were sterilized but still exposed to air through the swan neck of the flask. The media remained sterile despite exposure to air because the airborne bacteria got stuck in the neck and could not get to the media to contaminate it. This refuted the theory of spontaneous generation because proponents of that theory believed that air contained a “life force” that, when exposed to the media, chemically produced life from nonlife. 30. Needham and Spallanzani boiled broths for different lengths of time. Needham used only brief periods of boiling, whereas Spallanzani boiled his broths for as long as an hour. Thus, Needham likely did not boil his broths long enough to kill all of the microbes, some of which may have been heat resistant or survived by forming endospores. Another possibility is that the two scientists may have used different types of broths, which would have supported the growth of different types and numbers of microbes. 31. Schleiden and Schwann believed that cells arose as a result of a crystallization process, whereas Virchow and Remak believed that cells arose as a result of cell division. 32. Both mitochondria and chloroplasts have their own DNA that is genetically and structurally similar to bacterial DNA. Mitochondria and chloroplasts divide by binary fission independent of the host cell chromosome. We can see examples of endosymbiosis in modern-day organisms as well, including the photosynthetic bacteria-like organelles of some protists and the symbiotic bacteria found within the cells of some insects (e.g., cockroaches). 33. Semmelweis observed a higher incidence of puerperal fever in the maternity ward staffed by physicians and medical students than in the ward staffed by midwives. He observed that it was common practice for physicians and medical students to examine maternity patients after performing autopsies without washing their hands in between these procedures. He established the practice of having the physicians and medical students wash their hands in chlorinated lime water before examining maternity patients, and this practice greatly reduced the incidence of puerperal fever in this ward. 34. In these environments, water flows into cells. Without cell walls, the osmotic pressure would be too great for the bacterial cell membrane to withstand, and cells would quickly lyse. Cell walls allow bacterial cells living in hypotonic environments to withstand minor changes in osmotic pressure. 35. The flagella will rotate in a counterclockwise direction (run) more frequently and reduce the length of tumbles (rotation of flagella in a clockwise direction), resulting in an overall movement toward the attractant. 36. They have 70S (prokaryotic) ribosomes, a DNA genome most similar to that of prokaryotic cells, and two membranes. Also, mitochondria and chloroplasts divide in a way similar to binary fission. 37. Because eukaryotic cells are typically much larger than prokaryotic cells, they cannot rely exclusively on diffusion and active transport within the cell for the transport of materials across the cell; they use the endomembrane system to traffic some of these materials. 38. Possible answers: Bacterial flagella are composed of thin, stiff protein tubes of flagellin subunits and move in a propeller-like fashion. Eukaryotic flagella are composed of microtubules arranged in a 9+2 array. These microtubules slide along each other, allowing each flagellum to bend, producing movement with a whip-like motion. 39. Results of his experiment that would have supported the theory of spontaneous generation would have been the growth of

microbes in the flask media after sterilization, despite the presence of the swan necks. **40.** So many of the genes originally contained in the chromosomes of the bacteria from which these organelles arose have been transferred to the host nuclei, so these organelles no longer have a full complement of genes available to code for themselves independent of the host cell. The modern-day organelles need nuclear genes to function. **41.** Snow was the first to use epidemiology, tracing the incidence of disease and mapping it to locate its source. In doing so, he was able to determine that cholera was caused by fecally contaminated water. This was an important step in the development of the germ theory and clearly refuted the prevailing miasma theory of that time. His work also resulted in the first public health response to an epidemic, which was successful. **42.** (a) fimbriae, (b) cytoplasm, (c) nucleoid, (d) flagellum, (e) cell wall, (f) plasma membrane **43.** d **44.** Any component found only in bacteria would be a good target. Common antibiotic targets include peptidoglycan, found in bacterial cell walls, and the bacterial ribosome, because it is structurally distinct from the eukaryotic cytoplasmic ribosome. **45.** The bacterium would be difficult to kill because it would form endospores to survive harsh conditions, such as the high temperatures of cooking or boiling, dehydration, ultraviolet radiation, or chemical exposure. For this reason, the food industry must follow carefully designed protocols to process preserved foods that may contain these bacteria or their endospores. **46.** A. plasma membrane; B. nucleus; C. nucleolus; D. mitochondria; E. centrioles; F. endoplasmic reticulum; G. Golgi apparatus; H. flagellum **47.** Peroxisomes grow through the incorporation of peroxin proteins synthesized by free ribosomes in the cytoplasm. When large enough, they divide by binary fission, similar to mitochondria. They differ from mitochondria in their function and, unlike mitochondria, they contain neither their own DNA nor their own ribosomes. **48.** Both of these organelles contain components that are used for the degradation of molecules. Lysosomes contain digestive enzymes that degrade engulfed foodstuffs, infectious agents, and damaged cellular constituents. Peroxisomes contain hydrogen peroxide, which is a strong oxidizing agent that aids in the breakdown of uric acid, amino acids, and fatty acids. If the contents of these organelles were not contained within membranes, they would damage cytoplasmic components within cells.

#### Chapter 4

**1.** B **2.** C **3.** B **4.** A **5.** A **6.** D **7.** C **8.** B **9.** D **10.** A **11.** A **12.** B **13.** B **14.** D **15.** B **16.** True **17.** parasitism  
**18.** Eukarya **19.** antibiotic **20.** ammonia **21.** obligate **22.** *Helicobacter pylori* **23.** Gammaproteobacteria  
**24.** *Treponema pallidum pallidum* **25.** purple nonsulfur **26.** genus **27.** pyogenic **28.** high **29.** distance **30.** common ancestor  
**31.** vents **32.** ionizing radiation **33.** *Sulfolobus* **34.** *Methanobrevibacter oralis* **35.** Commensalism is a relationship between two organisms in which one benefits and the other is unaffected. Amensalism is a relationship between two organisms in which one is unaffected and the other is harmed. **36.** Long-term treatment with antibiotics can change the vaginal microbiota. If acid-producing bacteria (e.g., *Lactobacillus*) are eliminated by antibiotics, the less-acidic vagina becomes populated by yeasts. **37.** Coliforms ferment lactose completely, producing acid and gas. Noncoliforms either can't ferment lactose or ferment incompletely. Several species of intestinal pathogens are part of the noncoliform group. **38.** They are unable to produce ATP and are metabolically inactive outside of host cells. They are only able to replicate inside host cells, where they can use the intracellular ATP. **39.** CFB group is a large nontaxonomic group of gram-negative nonproteobacteria that includes the genera *Cytophaga*, *Fusobacterium*, and *Bacteroides*. **40.** *Borrelia burgdorferi*, a gram-negative nonproteobacterium in the Spirochaetae, infects arachnoids (ticks) and can be transmitted to humans by tick bites. It is a fastidious bacterium, which is extremely difficult to culture. Microscopically, it looks like a very thin spiral. **41.** The phylum Cyanobacteria is a large group of aquatic bacteria known for their blue-green pigment and ability to produce large amounts of gaseous oxygen during photosynthesis. Cyanobacteria play a major role in the maintenance of oxygen content in our planet's atmosphere. **42.** methicillin-resistant *S. aureus* (MRSA) and vancomycin-resistant *S. aureus* (VRSA). These two types *S. aureus* are resistant to multiple classes of antibiotics, which makes treatment of infections challenging **43.** Deeply branching bacteria provide scientists with clues as to the nature of life on earth billions of years ago. In industry, some of them are used for molecular hydrogen production. **44.** the unique mechanisms of DNA repair  
**45.** photosynthesis that uses bacteriorhodopsin **46.** Samples of the Martian atmosphere contain the gases produced

by the terrestrial archaea. 47. *Helicobacter pylori* 48. *Staphylococcus* 49. The marsh gas associated with methane bogs is produced by archaean methanogens.

### Chapter 5

1. C 2. C 3. C 4. A 5. C 6. D 7. A 8. B 9. D 10. A 11. C 12. B 13. A 14. B 15. plasmalemma 16. Fungi 17. Trematoda 18. dioecious 19. coenocytic 20. yeasts 21. antibiotics 22. pyrenoids 23. secondary endosymbiosis 24. These are modified mitochondria found in some protozoa. 25. Toxoplasmosis may affect behavior and personality. 26. The macronucleus contains many copies of genes for metabolism. 27. The best defense is making sure meat is well cooked. 28. *Trichophyton*, *Microsporium*, *Epidermophyton* 29. It is a cell formed from the fusion of two cells without the nuclei being fused. 30. Diatoms have silica shells known as frustules. 31. Algae are photosynthetic and can produce their own food, so they do not need to get it from other organisms. 32. brown algae (Phaeophyta), green algae (Chlorophyta), red algae (Rhodophyta) 33. They create soil, are food for animals, and are indicators of air pollution (as well as other possible answers). 34. D 35. Students can compare older approaches that used similarities in morphology and locomotion with newer approaches using molecular genetics. 36. There are many possible answers to this question. For example, protists that are heterotrophic are more likely to cause disease than autotrophic protists. Protists that are missing certain organelles or that are nonmotile (such as apicomplexans) are more likely to be parasitic or pathogenic than those that have all of the usual organelles and those that are motile. 37. Infection occurs by contact with parasites in fresh water, so anything that prevents people from being in water would break the life cycle of the parasite. 38. A 39. Chitin is a component of the fungal cell wall and is not a compound found in humans or any other vertebrate. Understanding how chitin is made can mean finding methods for interfering with its production in fungal pathogens. This may lead to a cure for fungal diseases that has limited side effects on their hosts.

### Chapter 6

1. B 2. A and B 3. B 4. B 5. C 6. B 7. D 8. D 9. B 10. A 11. B 12. D 13. C 14. D 15. C 16. True 17. bacteriophage 18. complex 19. naked or nonenveloped 20. tail fibers 21. reverse transcriptase 22. eclipse 23. Electron 24. cytopathic effects 25. RNA, helper virus





## Index

### Symbols

(RT-PCR) analysis, **239**

+ssRNA, **247**

70S ribosome, **97, 0**

70S ribosomes, **125**

80s ribosome, **0**

80S ribosomes, **118**

### A

Absorbance, **34**

absorbance, **0**

*Acanthamoeba*, **191**

acellular, **20, 25, 0, 230, 0**

*Acetabularia*, **222**

*Acetothermus*, **172**

*Acetothermus paucivorans*, **173**

acid fuchsin, **63**

acid-fast bacteria, **66**

acid-fast stain, **66, 0**

acidic dye, **62, 0**

actin, **122, 0**

Actinobacteria, **163**

*Actinomyces*, **164**

active transport, **104, 128**

affitins, **175**

aflatoxin, **213, 0**

African sleeping sickness, **121, 197**

African trypanosomiasis, **197**

agar, **218, 222, 0**

*Agrobacterium*, **148**

*Alexandrium*, **219**

Algae, **22**

algae, **0, 185, 218**

algal bloom, **218, 0**

algal chloroplasts, **126**

Alphaproteobacteria, **147, 163, 0**

*Amblyomma*, **158**

amensalism, **143, 0**

amoebic dysentery, **191**

Amoebozoa, **187, 191**

amphitrichous, **111, 0**

amplitude, **34, 0**

*Anabaena cylindrica*, **99**

anchorage dependency, **253**

*Ancylostoma duodenale*, **206**

*Ancylostoma duodenale*, **201**

anoxygenic photosynthesis, **161**

anthrax, **68, 101, 146, 168**

antibiotics, **232**

antifungal medications, **217**

apical complex, **193**

apicomplexans, **193**

*Aquifex*, **173**

Aquificae, **173**

archaea, **17, 0**

Archaea, **21, 140, 174, 0**

archaeal cell walls, **107**

archaeal membrane, **102**

Archaeplastida, **218**

Archaeplastids, **221**

Aristotle, **80**

*Arthrobacter*, **165**

*Ascaris lumbricoides*, **200, 206**

ascocarps, **213, 0**

Ascomycota, **213**

ascospore, **0**

ascospores, **213**

ascus, **213, 0**

*Aspergillus flavus*, **213**

atomic force microscope, **57, 0**

Attachment, **240**

attachment, **0**

atypical bacteria, **145**

atypical pneumonia, **169**

Avocado sunblotch viroid, **262**

axial filament, **157**

### B

*B. anthracis*, **68**

Bacillariophyta, **219**

Bacilli, **163, 168**

bacilli, **167**

bacillus, **0**

*Bacillus*, **168, 170**

*Bacillus anthracis*, **143, 146**

*Bacillus cereus*, **168**

bacteria, **17, 0**

Bacteria, **21, 140, 147**

bacterial cell walls, **105**

bacterial lawn, **251, 0**

bacterial meningitis, **149**

bacteriochlorophyll, **161**

bacteriochlorophylls, **104, 161, 0**

bacteriology, **26, 0**

bacteriophage, **231, 0**

bacteriophage preparation, **232**

Bacteriophages, **251**

bacteriorhodopsin, **176**

*Bacteroides*, **144, 145, 159, 160**

*Balantidium coli*, **194**

Baltimore classification system, **237**

*Bartonella*, **149**

basal body, **110, 130, 0**

basic dye, **62, 0**

basic fuchsin, **62, 68**

basidia, **215**

basidia (basidium, sing.), **0**

basidiocarps, **215, 0**

Basidiomycota, **215**

basidiospores, **215, 0**

*Bdellovibrio*, **154**

*Beggiatoa*, **153**

Beijerinck, **230**

Benda, **125**

*Bergey's Manual*, **18, 145**

Betaproteobacteria, **149, 163, 0**

*Bifidobacterium*, **164**

binocular, **44, 0**

binomial nomenclature, **18, 0**

biofilms, **56, 107**

biological vector, **232, 0**

bioluminescence, **152**

biosynthesis, **240, 0**

*Blastomyces dermatitidis*, **213**

blastomycosis, **213**

*Bordetella*, **150**

*Bordetella pertussis*, **145, 150**

*Borrelia*, **158**

botulinum neurotoxin, **166**

botulism, **101, 166**

brightfield microscope, **44, 0**

Brock, **139**

Brown, **86**

- brown algae, **219**
- Brucella*, **149**
- budding yeasts, **208, 0**
- Burkholderia*, **150**
- Burkholderiales, **150**
- burst, **248, 0**
- burst size, **248, 0**
- C**
- C. difficile*, **68**
- Campylobacter*, **156**
- Candida albicans*, **144, 208, 213**
- capsid, **234, 0**
- capsomere, **234, 0**
- capsule, **107, 0**
- capsule staining, **67, 0**
- capsules, **67**
- carbolfuchsin, **66**
- carbon fixation, **141**
- carboxysome, **99, 0**
- carotenoids, **161**
- carrageenan, **218, 222, 0**
- Caulobacter*, **149**
- cell envelope, **102, 0**
- cell membrane, **102, 0**
- cell morphology, **0**
- cell theory, **84, 0**
- cell wall, **95, 102, 105, 129, 0**
- cellular slime molds, **191**
- cellulose, **129**
- centriole, **0**
- centrioles, **124**
- centrosome, **0**
- centrosomes, **124**
- cesarean section, **144**
- cestodes, **204**
- CFB group, **145, 157, 159, 0**
- Chagas disease, **121, 197**
- Chamberland, **230**
- Chamberland filters, **230**
- chancroid, **151**
- Charophyta, **221**
- Chemical fixatives, **61**
- chemotaxis, **111, 0**
- chicken pox, **246**
- chitin, **129, 209, 217, 0**
- Chlamydia*, **145, 147**
- Chlamydomonas*, **221**
- Chlorella*, **222**
- Chlorobi, **163**
- Chlorobium*, **161**
- Chloroflexi, **163**
- Chloroflexus*, **161**
- chlorophyll, **126, 0, 161**
- chlorophylls, **104**
- Chlorophyta, **221**
- chloroplast, **0**
- chloroplasts, **126**
- chlorosomes, **161**
- Chromalveolata, **187, 193, 218**
- Chromatium*, **161**
- chromophores, **45, 0**
- chronic infection, **246**
- chronic wasting disease, **264**
- Chrysophyta, **219**
- chytrids, **211**
- cilia, **130, 187, 194**
- cilia (singular: cilium), **0**
- ciliate, **186**
- ciliates, **193, 0**
- cisternae, **119, 0**
- citrus tristeza virus, **247**
- CJD, **28**
- climate change, **142, 161**
- Clonorchis*, **201**
- Clostridium*, **166, 170**
- Clostridium difficile*, **167**
- Clostridium tetani*, **146**
- coarse focusing knob, **44, 0**
- cocci, **167**
- Coccidioides immitis*, **213**
- coenocytic hyphae, **207**
- coenocyte, **0**
- coenocytes, **117**
- coenocytic hyphae, **0**
- Coliforms, **152**
- colitis, **101**
- collagen, **129**
- color, **38**
- commensalism, **143, 0**
- community, **142, 0**
- competitive interactions, **143, 0**
- complex, **235**
- complex virus, **0**
- compound microscope, **41, 0**
- Conan the Bacterium, **173**
- concave lenses, **36**
- condenser lens, **44, 0**
- confocal microscope, **51, 0**
- conidia, **213, 0**
- conjugation, **186, 194**
- constitutive secretory pathways, **120**
- contact inhibition, **253**
- continuous cell line, **253, 0**
- contractile vacuole, **187**
- contractile vacuoles, **0**
- contrast, **39, 0**
- convex lens, **36**
- cooperative interactions, **143, 0**
- cortex, **223, 0**
- Corynebacterium*, **164**
- Corynebacterium diphtheriae*, **98, 146**
- counterstain, **63, 0**
- Coxiella*, **149**
- Crenarchaeota, **174**
- crenation, **95, 0**
- Creutzfeldt-Jakob disease, **264**
- Creutzfeldt-Jakob disease (CJD), **28**
- crustose lichens, **224, 0**
- Cryptococcus neoformans*, **215**
- Cryptosporidium parvum*, **194**
- crystal violet, **62**
- cucumber mosaic virus, **247**
- cyanobacteria, **21, 104, 0, 222**
- Cyanobacteria, **162**
- cyanobacterial blooms, **162**
- cyanophycin, **162**
- cyst, **185**
- cysticercosis, **199**
- cysts, **0**
- cytopathic effect, **0**
- Cytopathic effects (CPEs), **255**
- Cytophaga*, **159, 160**
- Cytophaga-Flavobacterium-Bacteroides* (CFB), **145**
- cytoplasm, **92, 0**

cytoplasmic membrane, **102, 0**  
 cytoproct, **186, 194, 0**  
 cytoskeleton, **113, 121, 0**  
 cytosome, **194**  
 cytostome, **186, 0**  
 cytotoxicity, **0**

## D

darkfield microscope, **46, 0**  
*De Contagione*, **40**  
 de Duve, **120**  
 decolorizing agent, **63, 0**  
 deeply branching bacteria, **145, 172, 0**  
*Deinococcus radiodurans*, **173**  
 Deltaproteobacteria, **154, 0**  
 dermatophytes, **213, 224**  
*Desulfovibrio*, **155**  
 diaphragm, **45, 0**  
 diatoms, **193, 219, 0**  
*Dictyostelium discoideum*, **191**  
 Diener, **261**  
 Differential interference contrast (DIC) microscopes, **50**  
 differential interference-contrast microscope, **0**  
 differential staining, **63, 0**  
 diffraction, **35, 0**  
 dikaryotic, **209, 0**  
 dimorphic fungi, **208**  
 dimorphic fungus, **0**  
 dinoflagellates, **193, 219, 0**  
 dioecious, **200, 0**  
 diphtheria, **164**  
*Diphylobothrium latum*, **205**  
 direct immunofluorescence assay (DFA), **51**  
*Dirofilaria immitis*, **201**  
 dispersion, **38, 0**  
 Dracunculiasis, **206**  
*Dracunculus medinensis*, **206**  
 dsDNA, **243**  
 dsRNA, **243**  
 dynein, **130, 0**

## E

Ebola epidemics, **248**

*Echinococcus granulosus*, **205**  
 eclipse phase, **247, 0**  
 ectoplasm, **186, 0**  
*Ehrlichia*, **149**  
*Eimeria*, **185**  
 electromagnetic radiation (EMR), **37**  
 electron microscope, **53, 0**  
 elementary bodies, **148, 0**  
 embryonated bird's egg, **252**  
 encystment, **185, 0**  
 endocarditis, **164**  
 endocytosis, **128, 0, 242**  
 endomembrane system, **118, 0**  
 endoplasm, **186, 0**  
 endoplasmic reticulum, **0**  
 endoplasmic reticulum (ER), **119**  
 endospore, **0**  
 Endospore staining, **68**  
 endospore staining, **0**  
 endospores, **49, 68, 99, 143, 168**  
 endosymbiotic hypothesis, **86**  
 endosymbiotic theory, **87, 125, 0**  
*Entamoeba*, **191**  
 enteric, **152, 0**  
*Enterobacter*, **153**  
 Enterobacteriaceae, **152**  
*Enterobius vermicularis*, **200**  
*Enterococcus*, **167, 170**  
*Enterocystozoan bieneusi*, **216**  
 enterotoxin, **169**  
 enveloped virus, **234, 0**  
 Enzyme immunoassays (EIAs), **259**  
 eosin, **63**  
 epidemic typhus, **147**  
*Epidermophyton*, **213**  
 epiphyte, **0**  
 epiphytes, **223**  
 Epsilonproteobacteria, **156, 0**  
 Epstein-Barr virus, **246**  
 ergosterols, **209, 217**  
*Erwinia*, **153**  
 Escherich, **153**  
*Escherichia*, **153**  
*Escherichia coli*, **143, 152, 168**  
*Escherichia coli* O157:H7, **107**

ether linkages, **174**  
*Euglena*, **197**  
 Euglenozoa, **197**  
 Eukarya, **0, 140**  
 eukaryote, **15, 0**  
 eukaryotic cell, **0**  
 eukaryotic cells, **92, 113**  
 eukaryotic flagella, **130**  
 eukaryotic membranes, **127**  
 Eumycetozoa, **191**  
 Euryarchaeota, **174**  
 eutroph, **149**  
 eutrophs, **0**  
 Excavata, **187**  
 excystment, **185, 0**  
 exocytosis, **129, 0**  
 extracellular matrix, **129, 0**  
 extrapolymeric substance (EPS), **56**  
 extremophiles, **174**

## F

F pilus, **109**  
 F pilus (F pili), **0**  
 facilitated diffusion, **103, 128**  
*Fasciolopsis buski*, **201**  
 fatal familial insomnia, **264**  
 fermentation, **6**  
 fimbriae, **109, 0**  
 fine focusing knob, **44, 0**  
 fixation, **61, 0**  
 Flagella, **68, 110**  
 flagella, **130, 0, 175, 187**  
 flagella staining, **68, 0**  
 flagellum (eukaryotic) (plural: flagella), **0**  
 flatworms, **201**  
 fluid mosaic model, **102, 0**  
 flukes, **201, 0**  
 fluorescence in situ hybridization (FISH), **56, 72**  
 fluorescence microscope, **50, 0**  
 fluorescent, **38, 0**  
 fluorochromes, **50, 0**  
 focal length, **36, 0**  
 focal point, **36, 0**  
 foliose lichens, **224, 0**  
 fomites, **232**

- Fornicata, **197**  
 Fox, **17**  
 Fracastoro, **40, 88**  
*Frankia*, **165**  
 free ribosome, **0**  
 free ribosomes, **118**  
 frequency, **34, 0**  
 fruiting bodies, **155**  
 frustule, **0**  
 frustules, **219**  
 fruticose lichens, **224, 0**  
*Fuligo septica*, **191**  
 Fungi, **23**  
 fungi, **0, 207, 211**  
 fungicides, **217**  
*Fusobacterium*, **144, 159, 160**
- G**
- Galileo, **41**  
 Gammaproteobacteria, **151, 161, 0**  
*Gardnerella*, **165**  
 gas gangrene, **101, 166**  
 Generalized transduction, **242**  
 generalized transduction, **0**  
 germ theory of disease, **88, 0**  
 germination, **101, 0**  
 Gerstmann-Straussler-Scheinker disease, **264**  
 giant viral species, **233**  
*Giardia lamblia*, **127, 197**  
 glycocalyx, **107, 0**  
 Golden Age of Microbiology, **11**  
 golden algae, **219**  
 Golgi, **120**  
 Golgi apparatus, **120, 0**  
 gonorrhea, **149**  
*Gonyaulax*, **219**  
 Gram, **63**  
 Gram stain procedure, **63, 0**  
 gram-negative bacteria, **145, 151**  
 gram-negative cells, **106**  
 gram-positive bacteria, **145**  
 gram-positive cells, **106**  
 Gram's iodine, **63**  
 granules, **164**  
 green algae, **221**  
 green nonsulfur bacteria, **161, 0**  
 green sulfur bacteria, **161, 0**  
 Group A strep, **109**  
 guinea worm, **25**  
 Guinea worm disease, **206**
- H**
- H1N1 influenza, **234**  
 H2N2, **234**  
 Haeckel, **14**  
*Haemophilus*, **144, 151**  
 Halobacteria, **176**  
*Halobacterium salinarum*, **176**  
*Haloferax volcanii*, **176**  
*Halothiobacillus neapolitanus*, **99**  
 Hämmerling, **222**  
 Hansen's disease, **164**  
 heartworm, **201**  
 heat fixing, **61**  
 HeLa cell line, **253**  
 helical, **235**  
 helical virus, **235, 0**  
*Helicobacter*, **156**  
*Helicobacter pylori*, **156**  
 helminth, **25, 0**  
 helper viruses, **262**  
 hemagglutination assay, **257**  
 hemagglutination inhibition (HAI) assays, **257**  
*Hemophilus*, **154**  
 hepatitis B virus (HBV), **262**  
 Hepatitis C virus, **246**  
 hepatitis delta virus (HDV), **262**  
 herpes simplex virus, **246**  
 heterocysts, **162**  
 high G+C gram-positive bacteria, **146, 163, 0**  
 Hippocrates, **9**  
*Histoplasma capsulatum*, **208, 213**  
 histoplasmosis, **208**  
 HIV, **246**  
 holdfast, **159**  
 holozoic, **187, 0**  
 Hooke, **41, 84**  
 Hooke's law of elasticity, **57**  
 hookworm, **201, 206**  
 host range, **231, 0**  
 human microbiome, **143**  
 Human Microbiome Project, **146**  
 hydrogenosomes, **127, 187**  
 hyperthermophiles, **173**  
 hypertonic medium, **95, 0**  
 hyphae, **207, 0**  
*Hyphomicrobium*, **149**  
 hypotonic medium, **95, 0**
- I**
- icosahedral, **235, 0**  
 illuminator, **44, 0**  
 image point (focus), **36, 0**  
 immunofluorescence, **50, 72, 72, 0**  
 immunofluorescent staining, **239**  
 Immunology, **26**  
 immunology, **0**  
 impetigo, **110, 167**  
 in vitro, **251, 0**  
 in vivo, **251, 0**  
 inclusions, **98, 0**  
 India ink, **67**  
 indirect immunofluorescence assay (IFA), **51**  
 induction, **240, 0**  
*Influenza B virus*, **234**  
 influenza vaccine, **252**  
 influenza virus, **242**  
 interference, **35, 0**  
 intermediate filament, **0**  
 intermediate filaments, **121**  
 International Classification of Diseases (ICD), **238**  
 International Committee on Taxonomy of Viruses (ICTV), **236**  
 International Union of Microbiological Societies, **236**  
 intestinal fluke, **201**  
 Irish potato famine, **196**  
 isoprene chains, **174**  
 isotonic medium, **95, 0**  
 Ivanovski, **230**  
*Ixodes*, **158**  
*Ixodes scapularis*, **194**
- J**
- Janssen, **42**

**K**

karyogamy, 209  
 keratitis, 191  
 kinetoplastids, 187  
 Kinyoun technique, 66, 0  
*Klebsiella*, 154  
 Koch, 11, 89  
 Korarchaeota, 174  
 kuru, 264

**L**

Lacks, 253  
 Lactobacillales, 167  
*Lactobacilli*, 144  
*Lactobacillus*, 167, 170  
*Laminaria*, 220  
 Lancefield, 109  
 Lancefield group, 167  
 last universal common ancestor (LUCA), 172  
 latent infection, 246  
 latent virus, 246, 0  
 Leeuwenhoek, 10  
*Legionella*, 152  
 Legionnaires disease, 152  
*Leishmania*, 199  
 lenses, 36  
 leprosy, 164  
*Leptospira*, 160  
*Leptothrix*, 150  
*Leuconostoc*, 167, 170  
 lichen, 223, 0  
 light microscopes, 43  
 Linnaean taxonomy, 13  
 Linnaeus, 13  
 lipopolysaccharide (LPS), 106, 0  
 Lippershey, 42  
 Lister, 89  
*Listeria monocytogenes*, 146  
 listeriosis, 146  
 liver flukes, 201  
 lophotrichous, 111, 0  
 low G+C gram-positive bacteria, 146, 163, 166, 166, 0  
 LUCA, 172  
 lumen, 119, 0

Lyme disease, 158  
 lymphogranuloma venereum, 148  
 lysis, 240, 0  
 lysogen, 240, 0  
 lysogenic conversion, 240  
 lysogenic conversion (phage conversion), 0  
 lysogenic cycle, 240, 0  
 lysogeny, 240, 0  
 lysosome, 0  
 lysosomes, 120  
 Lytic animal viruses, 242  
 lytic cycle, 240, 0

**M**

M protein, 110  
 macronucleus, 194, 0  
 mad cow, 264  
 magnetosomes, 99, 0  
*Magnetospirillum magnetotacticum*, 99  
 magnetotaxis, 111, 0  
 Magnification, 39  
 magnification, 0  
 malachite green, 63, 68  
 malaria, 183  
 Margulis, 87, 125  
 maturation, 240, 0  
 mechanical vector, 232, 0  
 medulla, 223, 0  
 membrane fusion, 242  
 membrane-bound ribosome, 0  
 membrane-bound ribosomes, 118  
 meningitis, 151  
 Mereschkowski, 86  
 mesophiles, 174  
 metachromatic granule, 0  
 metachromatic granules, 98  
 metagenomic analysis, 146  
 methanogen, 176, 0  
 methanogenesis, 174  
 methanogens, 176  
*Methanosarcina*, 176  
 methicillin-resistant *S. aureus*, 169  
 methylene blue, 63  
*Methylocystis*, 149  
*Methylomonas*, 154  
 miasma hypothesis, 8  
 miasma theory, 88  
 microbe, 6, 0  
 microbial ecology, 143, 0  
 Microbiology, 26  
 microbiology, 0  
 microbiome, 143, 0  
*Micrococcus*, 165  
*Microcystis*, 162  
 microfilament, 0  
 microfilaments, 121  
*Micrographia*, 41  
 micronucleus, 194, 0  
 microorganism, 6, 0  
 Microsporidia, 216  
 microsporidia, 0  
 microsporidiosis, 216  
*Microsporum*, 213  
 microtubule, 0  
 microtubules, 121, 187  
 mitochondria, 125  
 mitochondrial DNA, 125  
 mitochondrial matrix, 125, 0  
 mitochondrion (plural: mitochondria), 0  
 mitosomes, 127  
 mold, 24, 0  
 molds, 207  
 monocular, 44, 0  
 monoecious, 200, 0  
 monotrichous, 111, 0  
 mordant, 63, 0  
 morphologies, 115  
 morphology, 93  
 MRSA, 73, 169  
*Mucor*, 212  
 multidrug resistant bacteria, 232  
 murein, 105  
 mutualism, 143, 0, 223  
 mycelium, 207, 0  
*Mycobacterium*, 164  
*Mycobacterium tuberculosis*, 66, 146  
 mycolic acids, 105, 0  
 mycology, 26, 0  
*Mycoplasma*, 145, 169  
*Mycoplasma pneumoniae*, 132



mycorrhizal fungi, **211**  
 mycoses, **207**  
 mycoses (mycosis, sing.), **0**  
 myxobacteria, **155**  
*Myxobacterium*, **155**  
 myxospores, **155**

## N

NAAT, **259**  
*Naegleria fowleri*, **191**  
 naked virus, **234, 0**  
 Nanoarchaeota, **174**  
*Necator americanus*, **201**  
*Necatur americanus*, **206**  
 necrotizing fasciitis, **110, 167**  
 Needham, **81**  
 negative (–) single-strand RNA (–ssRNA), **243, 0**  
 negative stain, **62, 0**  
 neglected parasitic infections (NPIs), **199**  
*Neisseria*, **144, 149**  
 Nematoda, **200, 0**  
*Neurospora crassa*, **213**  
 neutralism, **143, 0**  
 nigrosin, **67**  
 nitrogen fixation, **141, 162**  
*Nocardia*, **165**  
 noncoliforms, **152**  
 nonenveloped virus, **234, 0**  
 nonproteobacteria, **157**  
 Nonproteobacteria, **160**  
 nonpyogenic streptococci, **167**  
 nonsulfur bacteria, **161**  
 nuclear envelope, **117, 0**  
 nuclear lamina, **117, 0**  
 nuclear membrane, **117**  
 Nucleic acid amplification tests (NAAT), **258**  
 nucleoid, **97, 0**  
 nucleoid-associated protein (NAP), **0**  
 nucleoid-associated proteins (NAPs), **97**  
 nucleolus, **117, 0**  
 nucleus, **116, 0**  
 numerical aperture, **39, 0**

## O

objective lenses, **44, 0**  
 obligate intracellular pathogen, **147, 0**  
 ocular lens, **44, 0**  
 oil immersion lens, **45, 0**  
 oligotroph, **147, 0**  
 one-step multiplication curve for bacteriophage, **248**  
 opacity, **35, 0**  
 Opisthokonta, **187**  
*Opistorchis*, **201**  
 oriental lung fluke, **201**  
 osmosis, **95, 0**  
 osmotic pressure, **95, 0**  
 Ötzi the Iceman, **8**  
 outer membrane, **106, 0**

## P

*palisades*, **164**  
 Pap smear, **259**  
 Parabasalia, **197**  
*Paragonimus westermani*, **201**  
*Paramecium*, **117, 194**  
 pararosaniline, **68**  
 parasitic helminths, **200**  
 parasitism, **143, 0, 223**  
 parasitology, **26, 0**  
 passive transport, **103**  
 Pasteur, **11, 82, 89, 230**  
*Pasteurellaceae*, **151**  
 pathogen, **21, 0**  
 Peach latent mosaic viroid, **262**  
 pellicle, **186, 197, 0**  
 penetration, **240, 0**  
 penicillin, **207**  
*Penicillium*, **207, 213**  
 peptidoglycan, **105**  
 peptidoglycan (murein), **0**  
 periodontitis, **164**  
 periplasmic space, **106, 0**  
 peritonitis, **153**  
 peritrichous, **111, 0**  
 peroxisome, **0**  
 peroxisomes, **121**  
 Persistent infection, **246**  
 pertussis, **150**  
*Pfiesteria piscicida*, **219**  
 Phaeophyta, **219**  
 phage conversion, **240**  
 phage therapy, **232**  
 phagocytosis, **128, 0**  
 phase-contrast microscope, **0**  
 Phase-contrast microscopes, **48**  
 phospholipid fatty acid analysis (PLFA), **103**  
 phosphorescence, **38, 0**  
 photosynthesis, **160, 176**  
 photosynthetic bacteria, **104**  
 phototaxis, **111, 0**  
 phototrophic bacteria, **160, 0**  
 phycobilisomes, **162**  
 phycocyanin, **162**  
 phylogenetic tree, **172**  
 phylogenies, **14**  
 phylogeny, **0**  
*Phytophthora*, **196**  
 phytoplankton, **185, 0**  
 pili, **109, 0**  
 pinocytosis, **128, 0**  
 pinworm, **200**  
 Planctomycetes, **157, 159**  
 plankton, **185, 0**  
 Plant viruses, **247**  
 plaque, **251, 0**  
 plasma membrane, **102, 0**  
 plasmalemma, **186, 0**  
 plasmid, **97, 0**  
 plasmodial slime molds, **191**  
*Plasmodium*, **183, 193**  
 plasmolysis, **95, 0**  
 Platyhelminthes, **201, 0**  
 pleomorphic, **169, 0**  
 pneumonia, **90, 168, 169**  
 polar tubule, **216, 0**  
 poliovirus, **242**  
 polyhedral, **235**  
 polyhedral virus, **0**  
 Polyhydroxybutyrate (PHB), **98**  
 polyhydroxybutyrate (PHB), **0**  
 Polymerase chain reaction, **258**  
 polyphyletic, **187, 0**

population, **142**  
 positive (+) single-strand (+ssRNA), **243**  
 positive (+) strand, **0**  
 positive stain, **62, 0**  
 potato tuber spindle disease, **261**  
 primary amoebic meningoencephalitis (PAM), **191**  
 primary cell culture, **252, 0**  
 primary stain, **63, 0**  
 prion, **263, 0**  
 prions, **28**  
 progeny virus, **239, 0**  
 proglottid, **204, 0**  
 prokaryote, **15, 0**  
 prokaryotes, **139, 140, 163**  
 prokaryotic cell, **0**  
 prokaryotic cells, **92**  
 properties of light, **34**  
 prophage, **240, 0**  
*Propionibacterium*, **165**  
 Proteobacteria, **145, 147, 154, 156, 160, 0**  
*Proteus*, **154**  
 Protist, **22**  
 protist, **0**  
 protist conjugation, **186**  
 protists, **185**  
 Protozoa, **23**  
 protozoan, **0**  
 protozoans, **185**  
 protozoology, **26, 0**  
 provirus, **244, 0**  
 Prusiner, **263**  
*Pseudo-nitzschia*, **220**  
 pseudohypha, **208**  
 pseudohyphae, **0**  
*Pseudomonas*, **151**  
*Pseudomonas aeruginosa*, **108, 111**  
 pseudomurein, **174**  
 pseudopeptidoglycan, **174**  
 pseudopodia, **122, 0, 187, 191**  
 purple nonsulfur bacteria, **161, 0**  
 purple sulfur bacteria, **161, 0**  
 pyrenoids, **218, 0**

## Q

quantum tunneling, **57**

## R

*R. rickettsii*, **147**  
 rat-bite fever, **111**  
 Receptor-mediated endocytosis, **128**  
 receptor-mediated endocytosis, **0**  
 red algae, **221**  
 red tide, **219, 0**  
 Redi, **80**  
 Reflection, **34**  
 reflection, **0**  
 refraction, **35, 0**  
 refractive index, **36, 0**  
 regulated secretory pathways, **120**  
 Remak, **85**  
 resident microbiota, **143, 0**  
 resolution, **39, 0**  
 respiration, **125**  
 retrovirus, **244, 0**  
 reverse transcriptase, **244, 0**  
 reverse transcriptase-polymerase chain reaction, **239**  
 Reverse transcriptase-polymerase chain reaction, **258**  
 Reverse zoonoses, **232**  
 rheostat, **45, 0**  
 rheumatic fever, **110**  
 Rhizaria, **187**  
 rhizines, **223, 0**  
*Rhizobium*, **141, 149**  
*Rhizopus stolonifer*, **212**  
*Rhodococcus*, **165**  
*Rhodocyclus*, **163**  
 Rhodophyta, **221**  
*Rhodospirillum*, **161**  
 ribosome, **0**  
 ribosomes, **92, 117**  
*Rickettsia*, **145, 147**  
 ringworm, **184, 213, 224**  
 RNA-dependent RNA polymerase (RdRP), **243**  
 Rocky Mountain spotted fever, **147**  
 rogue form, **0**

rose bengal, **63**  
 rough endoplasmic reticulum, **0**  
 rough endoplasmic reticulum (RER), **119**  
 roundworms, **200**  
 runs, **111**  
 runs (running), **0**  
 rusts, **211**

## S

S-layer, **108, 0**  
 Sabouraud's agar, **224**  
*Saccharomyces*, **213**  
 safranin, **63, 68**  
*Salmonella*, **153**  
 salmonellosis, **153**  
 saprozoic, **187, 0**  
 satellite RNA, **262**  
 scanning electron microscope (SEM), **53, 0**  
 scanning probe microscope, **57, 0**  
 scanning tunneling microscope, **57, 0**  
 scarlet fever, **110**  
 Schaeffer-Fulton method, **68**  
 Schimper, **86**  
*Schistosoma*, **201**  
 Schistosomiasis, **201**  
 schizogony, **186, 193, 0**  
 Schleiden, **84**  
 Schwann, **85**  
 scolex, **204, 0**  
 scrapie, **264**  
 secretory vesicle, **0**  
 secretory vesicles, **120**  
 Semmelweis, **88**  
 septate hyphae, **207, 0**  
 serotype, **0**  
 serotypes, **153**  
 serovars, **153**  
*Serratia*, **154**  
 sex pilus, **109, 0**  
 sheath, **235, 0**  
*Shewanella*, **139**  
 Shiga toxin, **153**  
*Shigella*, **154**  
 shingles, **246**

- simple diffusion, **103, 128**
  - simple microscope, **41, 0**
  - simple staining, **63, 0**
  - slime layer, **107, 0**
  - slime molds, **191**
  - smear, **61, 0**
  - smooth endoplasmic reticulum, **0**
  - smooth endoplasmic reticulum (SER), **119**
  - smuts, **211**
  - Snow, **89**
  - Sobemoviruses, **262**
  - Spallanzani, **81**
  - Specialized transduction, **242**
  - specialized transduction, **0**
  - Sphingobacterium*, **160**
  - spike, **234, 0**
  - Spirillum minus*, **111**
  - spirochetes, **145, 157, 0**
  - spontaneous generation, **80, 0**
  - spores, **209, 0**
  - sporulation, **100, 0**
  - ssDNA, **243**
  - ssRNA, **243**
  - stage, **44, 0**
  - staining, **62, 0**
  - staining patterns, **145**
  - Stanley, **234**
  - Staphylococcus*, **168, 170**
  - Staphylococcus aureus*, **73, 143, 169**
  - Staphylococcus epidermidis*, **143**
  - Stentor*, **194**
  - sterols, **127**
  - stigma, **197, 222, 0**
  - stramenopiles, **219, 0**
  - strep throat, **110, 167**
  - streptococci, **167**
  - Streptococcus*, **144, 146, 167, 170**
  - Streptococcus pneumoniae*, **108**
  - Streptococcus pyogenes*, **130**
  - Streptomyces*, **165**
  - stroma, **126, 0**
  - sulfate-reducing bacteria, **154**
  - Sulfolobus*, **175**
  - sulfur bacteria, **161**
  - superbugs, **74**
  - supergroups, **187**
  - symbiosis, **143, 0**
  - syngamy, **186, 0**
  - syphilis, **71, 158**
  - SYTOX Green, **73**
- ## T
- Taenia saginata*, **204**
  - Taenia solium*, **199**
  - tail fiber, **235, 0**
  - tail pins, **235, 0**
  - tapeworms, **204, 0**
  - Taxonomy, **13**
  - taxonomy, **0**
  - temperate phage, **0**
  - Temperate phages, **239**
  - tetanus, **101, 146, 166**
  - thallus, **207, 0**
  - Thaumarchaeota, **174**
  - theca, **219, 0**
  - Theileria (Babesia) microti*, **194**
  - Thermoproteus*, **175**
  - Thermotoga*, **173**
  - Thermotogae, **173**
  - thin sections, **71, 0**
  - Thiobacillus*, **150**
  - Thucydides, **10**
  - thylakoid, **126**
  - thylakoids, **0, 162**
  - ticks, **158**
  - tissue tropism, **242, 0**
  - tobacco mosaic disease, **230**
  - tobacco mosaic virus, **234**
  - toga, **173**
  - Tomato planta macho viroid, **262**
  - total magnification, **44, 0**
  - toxic shock syndrome, **110, 169**
  - toxicariasis, **199**
  - Toxocara canis*, **200**
  - toxocariasis, **201**
  - Toxoplasma gondii*, **194**
  - toxoplasmosis, **194, 199**
  - trachoma, **148**
  - transient microbiota, **143, 0**
  - transmissible spongiform encephalopathy, **263, 0**
  - transmission electron microscope (TEM), **53, 0**
  - transmittance, **34, 0**
  - transparency, **35, 0**
  - transport vesicle, **0**
  - transport vesicles, **119**
  - traveler's diarrhea, **153**
  - trematodes, **201**
  - Treponema*, **158, 160**
  - Treponema pallidum*, **47, 71, 145**
  - Trichinella spiralis*, **201**
  - trichinosis, **201**
  - Trichomonas vaginalis*, **127, 197**
  - trichomoniasis, **197**
  - Trichophyton*, **213**
  - Trichophyton rubrum*, **217, 224**
  - trophozoite, **185, 0**
  - Trypanosoma*, **197**
  - Trypanosoma brucei*, **198**
  - tuberculosis, **66, 146, 164**
  - tumbles, **111**
  - tumbles (tumbling), **0**
  - two-photon microscope, **52, 0**
  - Twort, **232**
  - typhoid fever, **153**
- ## U
- ultramicrotome, **71, 0**
  - Ureaplasma*, **170**
- ## V
- Valley fever, **213**
  - van Helmont, **80**
  - van Leeuwenhoek, **40**
  - vancomycin-resistant *S. aureus*, **169**
  - varicella-zoster virus, **246**
  - Varro, **10**
  - vector, **232, 0**
  - vegetative cell, **0**
  - vegetative cells, **99**
  - Vibrio*, **154**
  - Vibrio cholerae*, **151**
  - Vibrionales, **151**
  - viral envelope, **234, 0**
  - viral titer, **248, 0**
  - Virchow, **85**

virion, **231, 0**  
viroid, **0**  
Viroids, **261**  
virology, **26, 0**  
virulent phage, **0**  
Virulent phages, **239**  
Virus, **25**  
virus, **0, 230**  
virusoid, **262, 0**  
visible light, **37**  
Volutin, **98**  
volutin, **0**  
VRSA, **169**

## **W**

walking pneumonia, **169**  
Wallin, **86**  
water molds, **185, 196**  
wavelength, **34, 0**  
Western blot, **259**  
wet mount, **61, 0**  
Whittaker, **15**  
whooping cough, **150**  
Woese, **17, 147**  
Woods lamp, **184**

## **X**

x-y mechanical stage knobs, **44, 0**  
xenobiotics, **151**

## **Y**

yeast, **0**  
yeasts, **208**  
*Yersinia*, **154**  
*Yersinia pestis*, **153**

## **Z**

Ziehl-Neelsen technique, **66, 0**  
ZMapp, **249**  
zoonoses, **232**  
zooplankton, **185, 0**  
Zygomycota, **212**  
zygospores, **212, 0**